

Impaired Sensitivity to Thyroid Hormones is Associated with Mild Cognitive Impairment in Euthyroid Patients with Type 2 Diabetes [Response to Letter]

Zi-Wei Yu¹, Sheng-Dan Pu², Xiao-Tong Sun², Xi-Chang Wang¹, Xin-Yuan Gao², Zhong-Yan Shan¹

¹Department of Endocrinology and Metabolism and the Institute of Endocrinology, The NHC Key Laboratory of Diagnosis and Treatment of Thyroid Diseases, First Hospital of China Medical University, Shenyang, 110001, People's Republic of China; ²Department of Endocrinology, First Affiliated Hospital of Harbin Medical University, Harbin, 150001, People's Republic of China

Correspondence: Zhong-Yan Shan, Department of Endocrinology and Metabolism and the Institute of Endocrinology, The NHC Key Laboratory of Diagnosis and Treatment of Thyroid Diseases, First Hospital of China Medical University, Shenyang, 110001, People's Republic of China, Tel +8613804905018, Email cmushanzhongyan@163.com; Xin-Yuan Gao, Department of Endocrinology, First Affiliated Hospital of Harbin Medical University, Harbin, 150001, People's Republic of China, Tel +8613904517001; +86 451 8555 5060, Email m15124768260@163.com

Dear editor

We appreciate Sun et al for their interest and comments about our recent paper¹ in *Clinical Interventions in Ageing* regarding a relationship between thyroid hormone sensitivity and mild cognitive impairment (MCI) in euthyroid patients with type 2 diabetes (T2D). In terms of the exclusion criteria, Sun et al raised four points to be clarified or discussed, and we appreciate the opportunity to provide supplementary insights.

The first point concerns the lack of vitamin D data. We agree that vitamin D levels are generally low in the Chinese population with T2D and that reduced vitamin D concentrations may be associated with cognitive impairment. However, recent large-scale population studies have yielded conflicting results, indicating that vitamin D status may not be significantly associated with cognitive function.^{2,3} Given the existing contradictions in the current findings, it may be premature to definitively conclude that vitamin D plays a pivotal role in cognitive function. Furthermore, we observed that the previously published literature on cognitive impairment related to T2D^{4,5} or the association between thyroid function and cognitive function^{6,7} did not exclude individuals with vitamin D deficiency. Therefore, it is plausible that vitamin D deficiency may not be an obligatory exclusion criterion.

The second point is about the lack of data on bone mineral density (BMD) and bone turn over markers (BTMS). While we acknowledge the potential link between bone loss, osteoporosis, and cognitive impairment, it is noteworthy that there exists no consensus within the literature on this subject.⁸ Moreover, it appears that bone loss and osteoporosis are not established as recognized exclusion criteria in studies of cognitive impairment related to T2D^{4,5} or in studies examining the association between thyroid function and cognitive function.^{6,7}

The third point pertains to the lack of nutrition status data. We concur that major diseases, such as poor nutritional status, cancer, and severe liver or renal dysfunction, are closely related to cognitive impairment. Consequently, we excluded patients with these aforementioned conditions.

The fourth point regards the lack of estimated glomerular filtration rate (eGFR) data. First, as mentioned above, we have excluded patients with severe renal dysfunction. Secondly, it is regrettable that we did not collect eGFR data; however, we did obtain creatinine, which is another crucial indicator for assessing renal function. Our findings showed no significant difference in creatinine between the normal cognitive function group and the MCI group.

In conclusion, as confirmed by our study,¹ impaired sensitivity to thyroid hormones is associated with MCI in euthyroid patients with T2D. We actively recommend thyroid function examination for the early identification of MCI in

patients with T2D. It's crucial to emphasize that the focus on thyroid function should not solely revolve around single hormone alterations but should also encompass a consideration of thyroid hormone sensitivity.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Yu ZW, Pu SD, Sun XT, Wang XC, Gao XY, Shan ZY. Impaired sensitivity to thyroid hormones is associated with mild cognitive impairment in euthyroid patients with type 2 diabetes. *Clin Interv Aging*. 2023;18:1263–1274. doi:10.2147/CIA.S413584
2. Barmaki O, Mansour A, Moodi M, et al. Serum vitamin D status and cognitive function in Iranian older adults: evidence from the Birjand Longitudinal Aging Study. *J Nutr*. 2023;153(8):2312–2319. doi:10.1016/j.tjn.2023.06.033
3. Lee DH, Chon J, Kim Y, et al. Association between vitamin D deficiency and cognitive function in the elderly Korean population: a Korean frailty and aging cohort study. *Medicine*. 2020;99(8):e19293. doi:10.1097/MD.00000000000019293
4. Zheng T, Qin L, Chen B, et al. Association of plasma DPP4 activity with mild cognitive impairment in elderly patients with type 2 diabetes: results from the GDMD Study in China. *Diabetes Care*. 2016;39(9):1594–1601. doi:10.2337/dc16-0316
5. Zhang Z, Zhang B, Wang X, et al. Olfactory dysfunction mediates adiposity in cognitive impairment of type 2 diabetes: insights from Clinical and Functional Neuroimaging Studies. *Diabetes Care*. 2019;42(7):1274–1283. doi:10.2337/dc18-2584
6. Szlejf C, Suemoto CK, Santos IS, et al. Thyrotropin level and cognitive performance: baseline results from the ELSA-Brasil Study. *Psychoneuroendocrinology*. 2018;87:152–158. doi:10.1016/j.psyneuen.2017.10.017
7. Zhang H, Yang S, Zhu W, et al. Free triiodothyronine levels are related to executive function and scene memory in type 2 diabetes mellitus patients without diagnosed thyroid diseases. *Diabetes Metab Syndr Obes*. 2022;15:1041–1050. doi:10.2147/DMSO.S355656
8. Patel A, Jameson KA, Edwards MH, et al. Mild cognitive impairment is associated with poor physical function but not bone structure or density in late adulthood: findings from the Hertfordshire cohort study. *Arch Osteoporos*. 2018;13(1):44. doi:10.1007/s11657-018-0455-3

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>

<https://doi.org/10.2147/CIA.S439290>