

# Treatment Outcomes of Severe Acute Malnutrition and Its Determinants Among Paediatric Patients in Quetta City, Pakistan

Nazir Ahmed<sup>1</sup>, Fehmida Umar<sup>2</sup>, Fahad Saleem<sup>1</sup>, Qaiser Iqbal<sup>1</sup>, Sajjad Haider<sup>1</sup>, Mohammad Bashaar<sup>3</sup>

<sup>1</sup>Faculty of Pharmacy & Health Sciences, University of Balochistan, Quetta, Baluchistan, Pakistan; <sup>2</sup>Gynecology & Obstetrics Unit I, Sandeman Provincial Hospital Quetta, Quetta, Baluchistan, Pakistan; <sup>3</sup>Research Services Division, SMART Afghan International Trainings & Consultancy, Kabul, Afghanistan

Correspondence: Mohammad Bashaar, SMART Afghan International Trainings & Consultancy, Kabul, Afghanistan, Tel +93 788233865, Email dr.mbashar@gmail.com

**Purpose:** Severe acute malnutrition (SAM) is the most prevalent reason for admission to a paediatric unit, and it is a leading cause of mortality in many countries, including Pakistan. This study aimed to assess treatment outcomes and associated factors among children aged 6–59 months with severe acute malnutrition.

**Patients and Methods:** A retrospective cohort study was conducted at the Outpatient Therapeutic Feeding Program Centre established at the Sheikh Khalifa bin Zayed Al Nahyan Medical Complex Quetta. Out of 225 patients' records, data from 182 (80.8%) records were analysed based on the inclusion criteria. The SAM logbook was used as a source of data. Predictors of treatment outcomes were identified by applying a regression model with  $p < 0.05$  taken as significant.

**Results:** One hundred and twenty (65.9%) of the children were diagnosed with SAM, while the remaining 34.1% had Moderate Acute Malnutrition. Ninety-five (52.2%) children were included in the marasmus, while 47.8% were included in the Kwashiorkor cohort. The recovery rate was 68.6%; 22.5% were non-responsive, 11% defaulted on the program, and 3.5% died during management. The multivariate logistic regression identified the presence of diarrhea and the use of amoxicillin as significant prognosticators of treatment outcomes. Consequently, the odds of recovery on SAM among children with diarrhea [AOR = 0.60, 95% CI: (0.35–0.75)] were lower than those without diarrhea. Likewise, children on PO amoxicillin had higher chances of recovery [AOR = 2.45, 95% CI: (2.21–4.68)].

**Conclusion:** This study found that the recovery rate among children treated for SAM was poor based on the established Sphere Standard recommendation. In addition to community-based educational campaigns, capacity enhancement of OTP and frequent monitoring of services as well as program evaluation based on the management protocol is recommended to reduce the frequency of SAM among children.

**Keywords:** treatment outcomes, severe acute malnutrition, determinants, paediatric patients

## Introduction

Malnutrition (MN) refers to the imbalance of nutrients in the human body. Defined by the World Health Organization as deficiencies or excesses in nutrient intake, imbalance of essential nutrients, or impaired nutrient utilization,<sup>1</sup> Katoch et al argued that the definition of MN is not universally accepted.<sup>2</sup> Generally, the communal ideology reflects that MN results due to undernutrition or poor intake of nutrients by an individual. On the contrary, high nutrient intake and an erroneous amount of nutrient consumption are also described as MN.<sup>3</sup> Subsequently, MN is caused by a variety of interconnected factors<sup>4</sup> and is divided into two broad groups of conditions: undernutrition (including stunting, wasting, underweight and micronutrient deficiencies) and overweight, obesity, and diet-related non-communicable diseases.<sup>5</sup> Affecting the global population irrespective of location, age, wealth, or gender, MN is a universal crisis and explains about 45% of deaths among children under five. Furthermore, MN elevates the risk of non-communicable diseases contributing to global deaths and disabilities.<sup>5</sup>

In line with what is being discussed, The United Nations International Children's Emergency Fund (UNICEF), the World Health Organization (WHO), and The World Bank (WB) Group estimated that 149.2 million children had stunting, 45.4 million had wasting, 38.9 million were categorized as overweight and out of those 3.1 million children died because of MN in 2020.<sup>6</sup> Within this context, the WHO has emphasized addressing issues related to MN at the earliest. Especially among infants, Severe Acute Malnutrition (SAM), including its variants with or without medical consequences, must be prioritized.<sup>7</sup> Severe Acute Malnutrition is diagnosed in children with extremely low weight for height (below  $-3$  z scores of the median WHO growth criteria), by obvious severe wasting, or by the presence of oedema in both feet and the mid-upper arm circumference (MUAC) of 115 mm.<sup>8</sup> Severe Acute Malnutrition is categorized as kwashiorkor and marasmus, and both of these can commonly coexist.<sup>9</sup> Treatment for SAM is provided by using the Community-based Management of Acute Malnutrition model, one of which is an Outpatient Therapeutic Program (OTP) to treat cases without medical complications. This model was made possible by developing ready-to-use therapeutic food (RUTF), allowing most of the treatment in the patient's home.<sup>10</sup> To summarize, MN (severe acute or acute) has placed a persistent economic, social, and medical impact on communities and countries,<sup>11</sup> and is a constant global interest to researchers, social scientists, healthcare professionals, and policymakers.

As discussed earlier, half of the global children's deaths result from hunger and MN. A systematic review published in 2022 acknowledged child MN as a significant public health problem whereby 149 million children <5 years of age were estimated to be stunted, and 45 million were wasted.<sup>12</sup> Within this context, the bulk of instances (55 and 39%, respectively) occur in low- to middle-income Asian and African nations.<sup>13</sup> According to Hosen et al, stunting and underweight affect 37.9% and 33.6% of children in South Asia, and undernutrition is responsible for about 45% of fatalities in children below the age of five.<sup>14</sup> Given what has been said, the subcontinent alone is home to 58 million stunted children or 39% of all stunted children worldwide.<sup>15,16</sup> Shifting our concern to SAM in Pakistan, the country's frightening malnutrition rate results in an estimated three million deaths per year in children under the age of five, or 65.2 deaths per 1000 live births.<sup>17</sup> Pakistan has the third-highest proportion of stunted children globally, behind Nigeria and India. Stunting prevalence was projected to be 45%, with wasting at 10.5% and underweight at 31.6% in Pakistan.<sup>18</sup> Additionally, the Government of Pakistan reported an estimated 5.9% of Pakistani children under five with both wasted and stunted growth. This prevalence of concurrent wasting and stunting was highest in rural areas (6.8%), among children of mothers with no formal education (7.7%), and in households with the lowest quintiles of wealth (10.7%). Geographically speaking, Sindh province has the most significant frequency of concurrent stunting and wasting (10.0%), followed by Balochistan (6.5%), Punjab (4.3%), and Khyber Pakhtunkhwa (3.7%).<sup>19</sup>

Although recognized as a continuous crisis for decades, MN is still a silent threat to Pakistan's health, development, and future. Even though there is evidence of progress in various healthcare domains, MN is alarmingly prevalent and hinders the nation's growth and prosperity. Therefore, it is high time to prioritize this crisis and take immediate action to address its root causes. Investment in sustainable solutions, raising awareness, and fostering collaboration are needed.<sup>20</sup> Within this frame of reference and based on the broad strategic plan of the Pakistan Multi-Sectoral Nutrition Strategy (2018–2025), Pakistan has created plans for stunting reduction (as well as wasting management, maternal and adolescent nutrition) at the federal and provincial levels. By increasing the Average Annual Reduction Rate for stunting from 0.5% to 1%, this approach aims to lower the prevalence of stunting from 40% in 2018 to 33% in 2025. According to this approach, a multi-sectoral response has been designed and is being implemented to some extent in all of Pakistan's provinces.<sup>21</sup> One of the core components is the OTP, which uses ready-to-use therapeutic meals, community outreach, and mobilization, making treatments for the management of SAM accessible at decentralized treatment locations within primary healthcare settings. Although the OTP services are offered for free at healthcare centers, there is limited evidence of research-based program evaluation from Pakistan. Moreover, little is known about the outcomes level and factors determining the recovery rate of children from SAM. This scarcity of information regarding the evaluation of the OTP services in Quetta was a contributing reason for this research and is the study's unique contribution to the literature for local and international investigators. Hence, the current study aimed to fill these gaps by assessing the treatment outcomes of the OTP offered in Quetta City, Pakistan, regarding recovery, default, non-respondent, and mortality rates. Furthermore, the study was also focused on highlighting the factors associated with SAM among children aged 6–59 months.

## Materials and Methods

### Setting and Study Design

The study was conducted at Sheikh Khalifa Bin Zayed Al Nahyan Medical Complex (SKBZMC). Located at the periphery of Quetta city, SKBZMC is a general hospital fully equipped with the requirements of a modern hospital. The hospital started its operations in March 2003 and provides care to most of the local population at a subsidized cost that has been incurred by the Government of Balochistan since 2012.<sup>22</sup>

The OTP is offered to malnourished children under the administrative management of the Provincial Nutrition Directorate, Health Department, Government of Balochistan. Specialized healthcare workers, including physicians, pharmacists, and nurses, are trained to manage children with SAM with or without medical complications. Based on the protocol for the Community-based management of SAM,<sup>23</sup> nurses, in consultation with the physicians screen the children with SAM for admission into the program. Once admitted, the children are provided routine medications (where needed), monitor the child's progress, and provide referrals to inpatients when children fail appetite tests or develop complications. All services including screening, admission cost, RUTF, medications, and inpatient facilities are provided free of cost to the children.

We conducted a retrospective cohort study. The research design is suitable for indicating the temporal sequence between exposure and outcome. Furthermore, a cohort study allows examining multiple effects of a single exposure and avoids selection bias. However, differential losses to follow-up can bias retrospective cohort studies. To avoid loss to follow-up, every single possibility was adopted by the research team and data was extracted from every available source. Using the data of SAM children admitted to SKBZMC between 1st January 2022 to 31st December 2022. The data was collected from January 2023 and was finalized on 28 February 2023 (two months).

### Study Participants and Eligibility Criteria

This was a retrospective review of the eligible record of children treated for SAM in the SKBZMC. Children of age between 6–59 months, receiving treatment between January 1, 2022, and 31 December 2022, having MUAC values < 11.0 cm<sup>24</sup> or with bilateral pitting nutritional oedema,<sup>25</sup> and with no medical complications were selected for the study. We also categorized patients without oedema as marasmus and patients with oedema as the kwashiorkor group. On the contrary, patients referred to the OTP from other feeding centers, and being stable on arrival were excluded.

### Sample Size and Procedure

Based on the study objectives and eligibility criteria, we screened all children who visited the hospital for the management and treatment of SAM during the year 2022. Two hundred and twenty-five records were screened, and one hundred eighty-two children fulfilled the eligibility criteria. Hence, 43 records were discarded (13 patients aged > 6 months, 30 incomplete records) and an analysis of 182 children was carried out.

### Data Collection and Quality Assurance

The first author collected the data. A pre-validated structured questionnaire based on an extensive literature review<sup>26–29</sup> was used to extract information from the patient's records. The first author was trained by a research team and two physicians to ensure data accuracy and quality.

The first section of the questionnaire monitored the socio-demographic characteristics of each child including age, weight, locality, chaperone, gender, mother's living status, and travel time to the malnutrition center. The second section of the questionnaire recorded admission criteria presence/absence of bilateral pitting oedema, breastfeeding status, MUAC values, and the type of MN identified among the respondents. The third and fourth sections of the questionnaire offered clinical presentations, appetite test findings, and routine medications prescribed to the patients at the time of admission. The research team cleaned the data, ensuring completeness and consistency with continuous feedback from the physicians of the OTP unit.

## Operational Definitions

**Severe Acute Malnutrition:** Weight-for-height ratio of below 70% of the median NCHS reference, presence of nutritional or bilateral pitting oedema, and or MUAC < 11.5 cm.

**Malnutrition type:** Grouped as marasmus (non-oedematous) and kwashiorkor (oedematous) based on the diagnosis.

**Treatment outcome:** The treatment outcome was categorized into the “recovered group” and “non-recovered group” from SAM at the OTP unit in the current study.

**Recovered:** A child attaining 15% body weight and without oedema (two successive visits or weeks) declared cured by the healthcare professionals were categorized as recovered.

**Non-recovered:** If a child failed to recover at the end of the eighth week (2 months), he/she was rated as non-responsive and hence was referred to inpatient settings.

**Defaulter:** In case of absence of three consecutive follow-up visits, the child was declared as a defaulter.

**Death:** A patient in OTP who died during the management of SAM.

**Appetite test:** A child was said to pass the appetite test when he/she could consume the amount of RUTF recommended for his/her body weight.

**Admission type:** Admission to OTP refers to new admission [admitted for the first time or after 2 months of recovery] and readmission (admitted within 2 months).

**Ready to Use Therapeutic Food:** According to the protocol, children receive different numbers of RUTF [F-75 (75 kcal and 0.9 g protein per 100 mL) and F-100 (100 kcal and 2.9g protein per 100 mL)] sachets depending on body weight (as soon as the child stabilizes on 75 kcal, he/she is shifted to “catch-up formulation” of 100 kcal until wasted tissues are rebuilt).

**Medications:** Admission medication included amoxicillin, vitamin A, mebendazole, and metoclopramide.

**Rate of MUAC and weight gain:** The rate of MUAC gain in mm/day and weight gain in g/kg/day was calculated for the marasmus group. However, for the kwashiorkor group, the rate of MUAC and weight gain was assessed once the oedema resolved.

## Measurement of Variables

**Body weight:** Weight was measured by using the UNICEF infant, spring-type scale.<sup>30</sup>

**MUAC:** MUAC was measured with the help of UNICEF MUAC, 11.5 Red, PAC-50. The MUAC tape was placed on the left upper arm of a child with the arm hanging down the side of the body in a relaxed position.<sup>31</sup>

**Fever:** Fever was measured by taking the temperature rectally as recommended by the American Academy of Paediatrics.<sup>32</sup>

**Oedema:** Normal thumb pressure was applied on both feet for three seconds and edema was confirmed by shallow print on both feet.<sup>33</sup>

**Diarrhea, nausea, and vomiting:** These parameters were assessed by the physicians based on their observations and apparent signs and symptoms of the children.

## Statistical Analysis

Data were entered into and analysed by SPSS v 26.0. Descriptive analysis was initially conducted for the entire data. Treatment outcomes were dichotomized into recovered and non-recovered. The Kolmogorov–Smirnov test was used for normality assessment. Bivariate and Multivariate regression was also carried out to determine the association between dependent and independent variables. All statistical tests in this study were declared significant at <0.05.

## Ethical Approval and Consent to Participate

This study was carried out in accordance with the Declaration of Helsinki. Ethical approval was obtained from the Institutional Review Board of the Faculty of Pharmacy & Health Sciences, University of Balochistan, Quetta (FoP&HS/IRB/74/22). Permission from the Executive Director of SKBZMC was also acquired. All patients’ records were fully

anonymized before data collection and the authors had no access to information that could identify individual participants during or after data collection. Being a retrospective record review, the requirement for informed consent was waived by the Institutional Review Board of the Faculty of Pharmacy & Health Sciences, University of Balochistan, Quetta.

## Results

### Socio-Demographic Characteristics of the Malnourished Children

The study included 182 children who were managed for SAM under the OTP from January – December 2022. The gender distribution was almost equal ie, 96 (52.7%) males and 86 (47.3%) females. Most children (136, 74.7%) were in the age group of 6–15 months followed by 30 (16.5%) in the age group of 16–25 months. The mean weight of the cohort was  $5.15 \pm 1.18$  kgs with a median of 5. Most children had urban residency (151, 83%) and were accompanied by their mothers (140, 76.9%). One hundred and sixty-seven were within one hour's distance of the OTP center's location as shown in Table 1.

**Table 1** Socio-Demographics and Related Characteristics of Children with SAM

Characteristics	Frequency	Percentage
<b>Age (months)</b>		
6–15	136	74.7
16–25	30	16.5
26–35	4	2.2
36–45	10	5.5
46–54	2	1.1
<b>Weight (kgs)</b>		
< 5	69	37.9
5–10	113	62.1
<b>Locality</b>		
Quetta Rural	151	83.0
Quetta Urban	20	11.0
Kalat	4	2.2
Noshki	2	1.1
Chaghi	2	1.1
Much	3	1.6
<b>Chaperone</b>		
Mother	140	76.9
Father	25	13.7
Blood relative	17	9.4

(Continued)

**Table 1** (Continued).

Characteristics	Frequency	Percentage
<b>Gender</b>		
Male	96	52.7
Female	86	47.3
<b>Mother's living status</b>		
Alive	168	92.3
Deceased	14	7.7
<b>Travel time to the malnutrition center</b>		
Within 1 Hour	167	91.8
Two Hours	12	6.6
Three Hours	3	1.6

## Treatment Outcomes of the Outpatient Therapeutic Feeding Program

As mentioned earlier, the Sphere Standard acceptable ranges were used as a benchmark for evaluating treatment outcomes. One hundred and fifteen (68.65) children were cured and discharged from the OTP program. Forty-one (22.5%) were non-responsive and were referred to inpatient settings. Twenty (10.9%) defaulted on the program and 6 (3.2%) died during the management process. The relapse, screening refusal data, and vaccination history were not recorded by the healthcare professionals at the OTP unit (Table 2).

## Assessment of Malnutrition and Its Types

The MUAC was used to diagnose malnutrition among all children. One hundred and twenty (65.9%) of the children had MUAC values of < 11 cm and hence were categorized as SAM. The remaining 34.1% had MUAC values ranging from 11–12.5 cm and were diagnosed with Moderate Acute Malnutrition (MAM). Fifty-four (29.7%) children were breast-feeding and data for thirty-two (17.6%) was not recorded. Ninety-five (52.2%) of the children had no oedema and were included in the marasmus while 47.8% were included in the Kwashiorkor cohort (Table 3).

## Co-Morbidities Among the Children at the Time of Admission to the OTP Program

Forty-one children had diarrhea at enrolment to the OTP program followed by nausea and vomiting in 24.7% and fever in 28.5% of the children. Cough was also reported among 26.3% while the appetite test was positive for 74.1%. Nearly 80% recorded stool frequency of 1–3 times a day and urine flow (defined in terms of the size and force of the patient's urine stream) was positive among 93.4% of the children (Table 4).

**Table 2** Treatment Outcomes of the Outpatient Therapeutic Feeding Program

Treatment Outcomes		
Cured and discharged (recovered)	115	68.6
Referred to inpatient (non-responsive)	41	22.5
Defaulter	20	10.9
Death	6	3.2

**Note:** Relapse, screening refusal, and vaccination data were not recorded.

**Table 3** Assessment of Malnutrition and Its Types

Characteristics	Frequency	Percentage
<b>Admission criteria</b>		
MUAC	182	100.0
<b>MUAC (cm)</b>		
<11	120	65.9
11–12.5	62	34.1
<b>Breastfeeding</b>		
Yes	54	29.7
No	96	52.7
Not Recorded	32	17.6
<b>Malnutrition type</b>		
Kwashiorkor (Bilateral Pitting Edema positive)	67	36.9
Marasmus (Bilateral Pitting Edema negative)	115	63.1

**Table 4** Co-Morbidities at the Time of Admission to the OTP Program

Characteristics	Frequency	Percentage
<b>Diarrhea</b>		
Present	41	22.5
Absent	141	77.5
<b>Nausea and vomiting</b>		
Present	45	24.7
Absent	137	75.3
<b>Fever</b>		
Present	52	28.5
Absent	120	71.5
<b>Cough</b>		
Present	48	26.3
Absent	134	73.6
<b>Appetite</b>		
Positive	135	74.1
Negative	47	25.9

(Continued)

**Table 4** (Continued).

Characteristics	Frequency	Percentage
<b>Stool frequency (per day)</b>		
1–3	146	80.2
4–5	28	15.4
> 5	8	4.3
<b>Urine flow</b>		
Positive	170	93.4
Negative	12	6.6

## Management and Medication for Malnutrition

The management and medication for malnutrition are presented in Table 5. The RUTF (plumpy nuts) was offered to all children. Admitted cases with severe to moderate acute malnutrition to OTP were managed following the Pakistan National Guidelines for Community-based Management of Acute Malnutrition.<sup>23</sup> Out of the 182 children whose

**Table 5** Management and Medication for Malnutrition and Co-Morbid

Characteristics	Frequency	Percentage
<b>RUTF (Plumpy nuts)</b>		
Prescribed	182	100.0
Not prescribed	0	0
<b>Retinol (Vitamin A, 5000IU/day)</b>		
Prescribed	77	42.3
Not prescribed	105	57.7
<b>Anthelmintic (Mebendazole, 100 mg, BID for 3 days)</b>		
Prescribed	44	24.2
Not prescribed	138	75.8
<b>Antiemetic (Metoclopramide, 0.5 mg/kg in 24 hours for 5 days)</b>		
Prescribed	9	4.9
Not prescribed	173	95.1
<b>Antibiotic (Amoxicillin, 70–100 mg/kg/day in two divided doses for 5 days)</b>		
Prescribed	69	37.9
Not prescribed	113	62.1
<b>Analgesic (Paracetamol, 10 mg/kg, 3 times per 24 hours)</b>		
Prescribed	25	13.7
Not prescribed	157	86.3



medication records were available for review, the most prescribed medication was Vitamin A (PO, 42.3%) followed by Amoxicillin (PO, 37.9%) and Mebendazole (PO, 23.6%).

## Treatment Outcomes and Associated Factors

In the bivariate logistic regression analysis of malnutrition, the dependent variable was assessed for a relationship against the independent variables. The significant variables with  $p < 0.05$  (adjusting confounders) were later entered into the multivariable logistic regression model. The multivariate logistic regression identified the presence of diarrhea and the use of amoxicillin as significant prognosticators of treatment outcomes. Consequently, the odds of recovery on SAM among children with diarrhea [AOR = 0.60, 95% CI: (0.35–0.75)] were lower than those without diarrhea. Likewise, children on PO amoxicillin had higher chances of recovery [AOR = 2.45, 95% CI: (2.21–4.68)].

## Discussion

Severe Acute Malnutrition is a growing concern worldwide. Although much attention is provided to SAM, the efforts to decrease the frequency and prevalence around the globe are not yet satisfactory. Kassaw et al estimated that 1–2 million children die every year due to SAM, majority of them live in South Asia and Sub-Saharan Africa.<sup>34</sup> Within this context, Asia is home to the largest number of children under 5 years of age with SAM<sup>35</sup> and hence the SAM-related complications are also expected to be on the higher side. Shifting our concerns to SAM in Pakistan, UNICEF reported that nearly half of all children in Pakistan are chronic while 11% are acutely malnourished. The country has the worst breastfeeding indicators in South Asia and hence SAM contributes to high mortality and morbidity rates among children under 5 years of age. Those who survive are faced with poor mental and physical growth that leads to a significant economic impact on the country's development and prosperity.<sup>36</sup> To overcome this problem, Pakistan with the help of UNICEF and other organizations offers an OTP program for SAM children, however, a detailed study of the effectiveness of these programs and treatment outcome measures is not available in the literature. We, therefore, conducted this study to highlight the treatment outcomes and the associated factors related to the OTP program offered to children with SAM in Quetta City, Pakistan.

The result revealed that 115 (68.6%) SAM children admitted to OTP recovered. The recovery rate of the current study is lower than the Sphere Standard acceptable ranges (mortality  $< 3\%$ , default  $< 15\%$ , and recovered  $> 75\%$ ),<sup>37</sup> and to what is reported by several studies of the same nature.<sup>38–42</sup> However, it is comparable to other studies reporting recovery rates from SAM.<sup>43–47</sup> The conflicts of the recovery rates are attributed to the difference in settings where the OTP program was offered which is not the same in various healthcare settings. Also, the variance might be due to differences in socioeconomic status, quality of care provided for children, health-seeking behaviour, availability, and accessibility of therapeutic foods and medications at the time of OTP enrolment. Promisingly, the recovery rate is higher than reported in a study in Karachi, Pakistan.<sup>48</sup> Even though the recovery rate is not appreciated as per the Sphere Standards, comparing the results with the published literature revealed that the OTP program effectively achieved good survival and recovery rates. Similarly, the increased efforts of the Government and the support of UNICEF to improve maternal and child nutrition through several community-based programs may have also played a key role in this improvement. Nevertheless, we see a huge room for improvement, and increasing the recovery rate should be a priority for healthcare professionals and policymakers dealing with SAM in Pakistan.

Marasmus (63.1%) was the predominant form of MN in the current study and the finding is parallel to studies from other parts of the world.<sup>43,45,47,49</sup> The finding is not overwhelming as marasmus is more common in the age group below two years<sup>50</sup> which is also the case in this study as  $>70\%$  of the study population was in the age category of  $< 24$  months. An encouraging finding of the study was the death rate whereby this study reported a lower proportion of deaths (3.2%). The proportion of death rate is also lower than the recommended minimum Sphere Standard which should be  $< 10\%$ .<sup>37</sup> The lower death rate is attributed to the appropriate management of the children at the OTP and children reaching the OTP without complications. Our hypothesis is supported by a study from Ethiopia whereby the authors also presented similar reasons for lower death rates.<sup>43</sup>

This hospital-based study found that rates of SAM were noticeably identical for both genders (52.7% for males and 47.3% for females) and 74.7% had an age range of 6–15 months. Although there is no scientific evidence connected to

this finding, the gender distribution is the same as reported by other studies.<sup>43,51,52</sup> Nevertheless, there is one observation that we would like to highlight and discuss here. In the developing world, socioeconomic parameters and gender-related preferences are important factors that result in malnutrition. Within this context, male babies are favoured more by their parents, and gender inequality (quantity and quality of food provision) is widely reported and observed. We agree with the assertions of Shafiq et al, whereby the authors described that even today a female child is considered less important than a male child.<sup>53</sup> The authors also reported that this gender-based discrimination is noticed in nutrition provision, education, and health. Hence, many female children suffer from SAM compared to their counterparts.<sup>53</sup> However, the current study while comparing the proportion found no gender-based favour and SAM was noticed in equal percentages among males and females and that is a positive sign of reduced gender-based discrimination.

Results of the current study highlighted that nearly thirty percent of the children were breastfed, while over 50% were not. The role of breastfeeding in the prevention of MN is well recognised in the literature.<sup>54,55</sup> Although the current study was not aimed to associate breastfeeding and MN, the lack of breastfeeding and development of MN is linked to several factors. Malnutrition negatively impacts school outcomes, cognitive development, and overall Gross Domestic Product which in the long term are disastrous to any nation. Our claims are in line with what is reported by Katoch et al in their two different systematic reviews.<sup>56,57</sup> The authors discussed the deleterious effects of MN and its negative effect on the health and overall well-being of society. The authors further concluded indirect consequences that must be dealt with a prompt necessity.<sup>58</sup> Consequently, based on the findings of the current study and upon the available evidence educating mothers exclusively about breastfeeding is recommended to avoid MN-related complications.

In the current study, the use of amoxicillin and the recovery from diarrhea were identified as predictors of treatment success. The use of PO amoxicillin has shown a positive impact on treatment outcomes specifically in the cases of SAM. Correlating our results, children receiving PO amoxicillin reported improved recovery from SAM as compared to those who did not take them with odds of 3.38 and 1.95 respectively.<sup>43,47</sup> The supportive role of amoxicillin in SAM is well documented in the literature. Trehan et al in their controlled trial concluded that adding amoxicillin to the management of SAM resulted in remarkable improvements in recovery rates. Significant improvements in weight and gain in the MUAC were also observed. The authors consequently established that adding amoxicillin to therapeutic regimens for SAM was associated with significantly improving retrieval and mortality rates.<sup>59</sup> Likewise, Williams and Berkley in their systematic review established the continued use of broad-spectrum oral amoxicillin for treating children with uncomplicated SAM as outpatients which again supports the results of the present study.<sup>60</sup>

Results of the current study reported an association of diarrhea with SAM. The odds of recovery on SAM among children with diarrhea were lower compared to those without diarrhea. It is a well-known fact that diarrhea is usually associated with infections that result in further nutritional consumption. This shares nutrients and may delay the recovery time among children with SAM. The relationship between diarrhea and physical growth is recognized whereby the presence of diarrhea and a delay in recovery time was associated in literature.<sup>61–63</sup> Each diarrheal episode makes MN worse as diarrhea causes cytokine-induced MN which results from the actions of proinflammatory cytokines.<sup>64</sup> Hence, diarrhea-related MN interaction represents a dangerous web that can be disentangled by promoting breastfeeding, improving hygiene conditions, and improving nutrition. However, concurrently using PO antibiotics (amoxicillin) among children with MN can increase antibiotic-associated diarrhea episodes (AAD). Although AAD is documented in the literature, the underlying mechanisms of AAD in children remain unclear. Kuehn et al, in their review, highlighted the increased incidence of AAD following the use of oral penicillin and therefore advised consideration when prescribing antibiotics, especially in children.<sup>65</sup> Consequently, we recommend closely monitoring the malnourished children on amoxicillin and taking immediate remedial actions if AAD develops.

## Limitations

The study does have certain limitations. The relapse, screening refusal, and vaccination data were not recorded at the OTP center. These are important indicators and could have provided additional information. The generalizability of the findings is always questionable in a single-centered study.

## Recommendations

We recommended a study with multiple OTP centers and a larger sample to get an in-depth view of MN and the accompanying factors. Relapse, screening refusal, and vaccination data must be included to get a clear picture of MN and its associated factors that will help in developing effective interventions for malnourished children.

## Conclusion

Recovery rates reported by the current study are below the cut-off points of the minimum standards, nevertheless, the death rate was lower. Diarrhea was a statistically significant factor that impeded the recovery rate of children with SAM. On the other hand, PO amoxicillin was a positive indicator of recovery. Consequently, healthcare providers should pay exclusive attention to SAM cases with diarrhea and ensure strict follow-up according to the protocol. Healthcare providers must be provided with continuous medical education specifically focused on managing conditions like diarrhea that will help them in the targeted management of MN. We also recommended providing community-based health education and counselling for mothers regarding breastfeeding, hygiene measures, and when to approach the OTP center to avoid complications among the SAM children. A well-informed and educated mother will be an excellent resource in reducing MN frequency hence reducing the healthcare burden and enhancing community well-being.

## Acknowledgments

We sincerely thank the administration of SKBZMC and healthcare professionals stationed at the OTP for their assistance during the data collection and curation process.

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. World Health Organization. Malnutrition. Available from: [https://www.who.int/health-topics/malnutrition#tab=tab\\_1](https://www.who.int/health-topics/malnutrition#tab=tab_1). Accessed August 20, 2023.
2. Katoch OR, Sharma A, Nawaz A, Shrikant A. Nutritional assessment using Composite Index of Anthropometric Failure (CIAF) among school-going Children in J&K, India. *J Tianjin Univ Sci*. 2021;54(8):404–420.
3. Kobylńska M, Antosik K, Decyk A, Kurowska K. Malnutrition in obesity: is it possible? *Obes Facts*. 2022;15(1):19–25. doi:10.1159/000519503
4. Katoch OR. Author's response to comment on "Determinants of malnutrition among children: a systematic review". *Nutrition*. 2022;102:111733. doi:10.1016/j.nut.2022.111733
5. Global Nutrition Report. About malnutrition. Available from: <https://globalnutritionreport.org/resources/about-malnutrition/>. Accessed June 9, 2023.
6. World Health Organization. UNICEF/WHO/The World Bank Group joint child malnutrition estimates: key findings of the 2021 edition. Available from: [https://www.who.int/data/gho/data/themes/topics/joint-child-malnutrition-estimates-unicef-who-wb#:~:text=In%202020%2C%20149.2%20million%20children,for%20their%20height%20\(overweight\)](https://www.who.int/data/gho/data/themes/topics/joint-child-malnutrition-estimates-unicef-who-wb#:~:text=In%202020%2C%20149.2%20million%20children,for%20their%20height%20(overweight)). Accessed June 1, 2023.
7. World Health Organization. Updates on the management of severe acute malnutrition in infants and children. Available from: [https://apps.who.int/iris/bitstream/handle/10665/95584/9789241506328\\_eng.pdf;jsessionid=800ECEF1FF1D0B67DE1EE1141980A19?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/95584/9789241506328_eng.pdf;jsessionid=800ECEF1FF1D0B67DE1EE1141980A19?sequence=1). Accessed May 25, 2023.
8. Kebede F, Kebede T, Negese B, Abera A, Fentaw G, Kasaw A. Incidence and predictors of severe acute malnutrition mortality in children aged 6–59 months admitted at Pawe general hospital, Northwest Ethiopia. *PlosOne*. 2022;17(2):e0263236. doi:10.1371/journal.pone.0263236
9. Alou MT, Golden MH, Million M, Raoult D, Raoult D. Difference between kwashiorkor and marasmus: comparative meta-analysis of pathogenic characteristics and implications for treatment. *Microb Pathog*. 2021;150:104702. doi:10.1016/j.micpath.2020.104702
10. Rogers E, Tappis H, Doocy S, et al. Costs and cost-effectiveness of three point-of-use water treatment technologies added to community-based treatment of severe acute malnutrition in Sindh Province, Pakistan. *Glob Health Action*. 2019;12(1):1568827. doi:10.1080/16549716.2019.1568827
11. Katoch OR. Is maternal higher education effective in preventing child malnutrition? Evidence from literature. *Public Health Ancient Sci*. 2023;3(1):3–13.
12. Katoch OR. Determinants of malnutrition among children: a systematic review. *Nutrition*. 2022;96:111565. doi:10.1016/j.nut.2021.111565
13. Black RE, Victora CG, Walker SP, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013;382(9890):427–451. doi:10.1016/S0140-6736(13)60937-X
14. Hosen MZ, Pulok MH, Hajizadeh M. Effects of maternal employment on child malnutrition in South Asia: an instrumental variable approach. *Nutrition*. 2023;105:111851. doi:10.1016/j.nut.2022.111851
15. World Health Organization. Levels and trends in child malnutrition: UNICEF/WHO/The World Bank Group joint child malnutrition estimates: key findings of the 2021 edition. Available from: <https://www.who.int/publications/i/item/9789240025257>. Accessed June 2, 2023.
16. Wali N, Agho KE, Renzaho AM. Factors associated with stunting among children under 5 years in five South Asian countries (2014–2018): analysis of demographic health surveys. *Nutrients*. 2020;12(12):3875. doi:10.3390/nu12123875
17. Global Nutrition Report. Nourishing the SDGs. Available from: [https://globalnutritionreport.org/documents/822/Global\\_Nutrition\\_Report\\_2017.pdf](https://globalnutritionreport.org/documents/822/Global_Nutrition_Report_2017.pdf). Accessed May 25, 2023.
18. Rebouças P, Falcão IR, Barreto ML. Social inequalities and their impact on children's health: a current and global perspective. *J Pediatr*. 2022;98:55–65. doi:10.1016/j.jpeds.2021.11.004

19. Shuja S, Ategbro EA, Ihtesham Y, Ahmed KM Wasting prevention and treatment - central to stunting reduction in Pakistan. Available from: <https://www.enonline.net/fex/63/wastingstuntingconnectionpakistan#:~:text=An%20estimated%205.9%25%20of%20children,households%20within%20the%20poorest%20wealth>. Accessed June 1, 2023.
20. Effendi A Malnutrition in Pakistan — urgent action needed. Available from: <https://tribune.com.pk/story/2420101/malnutrition-in-pakistan-urgent-action-needed>. Accessed August 21, 2023.
21. Emergency Nutrition Network. Wasting prevention and treatment—central to stunting reduction in Pakistan. Available from: <https://www.enonline.net/fexdigest/63/wastingstuntingconnectionpakistan>. Accessed June 2, 2023.
22. Ishaq R, Baloch NS, Iqbal Q, et al. Frequency and evaluation of the perceptions towards caesarean section among pregnant women attending public hospitals in Pakistan and the implications. *Hosp Pract*. 2017;45(3):104–110. doi:10.1080/21548331.2017.1328250
23. Ministry of National Health Services Regulations and Coordination. Pakistan national guidelines for community-based management of acute malnutrition. Available from: <https://www.unicef.org/pakistan/media/4161/file/CMAM%20Guidelines.pdf>. Accessed June 10, 2023.
24. Eaton-Evans J. Nutritional assessment: anthropometry. In: Caballero B, editor. *Encyclopedia of Human Nutrition*. Elsevier Inc; 2013:311–318.
25. Dominique R, Naïma H, Pankti M, et al. Management of oedematous malnutrition in infants and children aged >6 months: a systematic review of the evidence. Available from: [https://cdn.who.int/media/docs/default-source/nutritionlibrary/publications/malnutrition/guideline-updates-on-The-management-of-severe-acute-malnutrition-in-infants-and-children/updates-management-sam-infantandchildren-review2.pdf?sfvrsn=7362c6d6\\_4](https://cdn.who.int/media/docs/default-source/nutritionlibrary/publications/malnutrition/guideline-updates-on-The-management-of-severe-acute-malnutrition-in-infants-and-children/updates-management-sam-infantandchildren-review2.pdf?sfvrsn=7362c6d6_4). Accessed June 10, 2023.
26. Katoch OR, Sharma A, Nawaz A. Determinants of malnutrition (stunting) among rural farming households: evidences from rural areas of District Doda, Jammu & Kashmir. *Asian J Soc Sci Humanit*. 2017;7(7):166–176. doi:10.5958/2249-7315.2017.00375.6
27. Aguayo VM, Badgaiyan N, Qadir SS, et al. Community management of acute malnutrition (CMAM) programme in Pakistan effectively treats children with uncomplicated severe wasting. *Matern Child Nutr*. 2018;14:e12623. doi:10.1111/mcn.12623
28. Khan GN, Turab A, Khan MI, et al. Prevalence and associated factors of malnutrition among children under-five years in Sindh, Pakistan: a cross-sectional study. *BMC Nutr*. 2016;2(1):69. doi:10.1186/s40795-016-0112-4
29. Menghwar B, Laghari ZA, Memon SF, Warsi J, Shaikh SA, Baig NM. Prevalence of malnutrition in children under five years' age in District Tharparkar Sindh, Pakistan. *J Pak Med Assoc*. 2022;72(01):33–36. doi:10.47391/JPMMA.20-540
30. UNICEF. Scale, infant, springtype, 25kg x 100g. Available from: <https://supply.unicef.org/s0145555.html>. Accessed June 4, 2023.
31. UNICEF. Mid-upper arm circumference (MUAC) measuring tapes – technical bulletin. Available from: <https://www.unicef.org/supply/media/1421/file/mid-upper-arm-circumference-measuring-tapes-technical-bulletin.pdf>. Accessed June 5, 2023.
32. American Academy of Pediatrics. Infant fever. Available from: <https://www.aap.org/en/patient-care/infant-fever/>. Accessed June 2, 2023.
33. World Health Organization. WHO child growth standards and the identification of severe acute malnutrition in infants and children: joint statement by the World Health Organization and the United Nations Children's Fund. Available from: <https://apps.who.int/iris/bitstream/handle/10665/44129/9789?sequence=1>. Accessed June 4, 2023.
34. Kassaw A, Amare D, Birhanu M, et al. Survival and predictors of mortality among severe acute malnourished under-five children admitted at Felege-Hiwot comprehensive specialized hospital, northwest, Ethiopia: a retrospective cohort study. *BMC Pediatr*. 2021;21(1):1–10. doi:10.1186/s12887-021-02651-x
35. Ahmed T, Hossain M, Mahfuz M, et al. Severe acute malnutrition in Asia. *Food Nutr Bull*. 2014;35(2\_suppl1):S14–S26. doi:10.1177/15648265140352S103
36. Malik AS. Fighting malnutrition in Pakistan with a helping hand from children abroad. Available from: <https://www.unicef.org/stories/fighting-malnutrition-pakistan-helping-hand-children-abroad>. Accessed June 7, 2023.
37. Association S. The Sphere handbook: humanitarian charter and minimum standards in humanitarian response. Available from: <https://spherestandards.org/wp-content/uploads/Sphere-Handbook-2018-EN.pdf>. Accessed June 29, 2023.
38. Das K, Das S, Mohapatra S, Swain A, Mohakud NK. Risk and adverse outcome factors of severe acute malnutrition in children: a Hospital-Based Study in Odisha. *Cureus*. 2021;13(9):e18364. doi:10.7759/cureus.18364
39. Kangas ST, Marron B, Tausanovitch Z, et al. Effectiveness of acute malnutrition treatment at health center and community levels with a simplified, combined protocol in Mali: an Observational Cohort Study. *Nutrients*. 2022;14(22):4923. doi:10.3390/nu14224923
40. Mekuria G, Derese T, Hailu G. Treatment outcome and associated factors of severe acute malnutrition among 6–59 months old children in Debre Markos and Finote Selam hospitals, Northwest Ethiopia: a retrospective cohort study. *BMC Nutr*. 2017;3(1):1–8. doi:10.1186/s40795-017-0161-3
41. Misrak A. *Treatment Outcome of Severe Acute Malnutrition and Associated Factors Among Under five Children Admitted at Hospitals in Arsi Zone*. Oromia Regional State, South East Ethiopia: Addis Ababa University; 2018.
42. Simachew Y, Zerfu T, Alemu W. treatment outcomes and predictors of recovery from severe acute malnutrition among children aged 6–59 months attending an outpatient therapeutic program in Wenago District, Southern Ethiopia. *Nutr Diet Suppl*. 2020;Volume 12:189–200. doi:10.2147/NDS.S245070
43. Abate BB, Tilahun BD, Kassie AM, Kassaw MW, Gurgel RQ. Treatment outcome of severe acute malnutrition and associated factors among under-five children in outpatient therapeutics unit in Gubalafto Wereda, North Wollo Zone, Ethiopia, 2019. *PloSOne*. 2020;15(9):e0238231. doi:10.1371/journal.pone.0238231
44. Bizuneh F, Bekonjo N, Kebede T. Severe acute malnutrition treatment cure rate for under-five children admitted in stabilizing center: north west Ethiopia 2021. *J Nutr Med Diet Care*. 2021;7:1–9.
45. Kabalo MY, Seifu CN. Treatment outcomes of severe acute malnutrition in children treated within Outpatient Therapeutic Program (OTP) at Wolaita Zone, Southern Ethiopia: retrospective cross-sectional study. *J Health Popul Nutr*. 2017;36(1):1–8. doi:10.1186/s41043-017-0083-3
46. Nyeko R, Calbi V, Ssegujja BO, Ayot GF. Treatment outcome among children under-five years hospitalized with severe acute malnutrition in St. Mary's hospital Lacor, Northern Uganda. *BMC Nutr*. 2016;2(1):1–7. doi:10.1186/s40795-016-0058-6
47. Yebo HG, Kendall C, Nigusse D, Lemma W, Bhutta ZA. Outpatient therapeutic feeding program outcomes and determinants in treatment of severe acute malnutrition in Tigray, northern Ethiopia: a retrospective cohort study. *PloSOne*. 2013;8(6):e65840. doi:10.1371/journal.pone.0065840
48. Ahmed S, Ejaz K, Mehnaz A, Adil F. Implementing WHO feeding guidelines for inpatient management of malnourished children. *J Coll Physicians Surg Pak*. 2014;24(7):493–497.
49. Munthali T, Jacobs C, Sitali L, Dambe R, Michelo C. Mortality and morbidity patterns in under-five children with severe acute malnutrition (SAM) in Zambia: a five-year retrospective review of hospital-based records (2009–2013). *Arch Public Health*. 2015;73(1):1–9. doi:10.1186/s13690-015-0072-1

50. Bhagavan NV, Chung-Eun H. *Essentials of Medical Biochemistry*. Elsevier Inc; 2015.
51. Mena MB, Dedefo MG, Billoro BB. Treatment outcome of severe acute malnutrition and its determinants among pediatric patients in West Ethiopia. *Int J Pediatr*. 2018;2018:1–7. doi:10.1155/2018/8686501
52. Mezemer M, Girma M, Bekele D. Treatment outcome and associated factors of acute malnutrition among children in the therapeutic feeding center of Public Hospitals in Addis Ababa, Ethiopia: an Institutional-Based Cross-Sectional Study. *Pediatric Health Med Ther*. 2022;Volume 13:145–154. doi:10.2147/PHMT.S296979
53. Shafiq A, Hussain A, Asif M, Jameel A, Sadiq S, Kanwel S. Determinants of gender disparity in nutritional intake among children in Pakistan: evidence from PDHS. *Children*. 2021;9(1):7. doi:10.3390/children9010007
54. Scherbaum V, Srouf ML. The role of breastfeeding in the prevention of childhood malnutrition. *World Rev Nutr Diet*. 2016;115:82–97.
55. World Health Organization. Infant and young child feeding. Available from: <https://www.who.int/news-room/fact-sheets/detail/infant-and-young-child-feeding>. Accessed August 23, 2023.
56. Katoch OR, Sehgal S, Shrikant A, Nawaz A, Parihar S, Ahmed A. Analysis of the effects of child malnutrition on school outcomes, cognitive development, and gross domestic product (GDP): a systematic review. *Int J Food Nutri Sci*. 2020;11(S1):2932–2954.
57. Katoch OR, Sharma R, Parihar S, Nawaz A. Energy poverty and its impacts on health and education: a systematic review. *Int J Energy Sect Manag*. 2023. doi:10.1108/IJESM-10-2022-0007
58. Katoch OR. Exploring the relationship between expenditure on power and state finances: an empirical study in Jammu and Kashmir, India. *Environ Dev Sustain*. 2023. doi:10.1007/s10668-023-03734-8
59. Trehan I, Goldbach HS, LaGrone LN, et al. Antibiotics as part of the management of severe acute malnutrition. *N Engl J Med*. 2013;13(268):425–435. doi:10.1056/NEJMoa1202851
60. Williams PC, Berkley JA. Guidelines for the treatment of severe acute malnutrition: a systematic review of the evidence for antimicrobial therapy. *Paediatr Int Child Health*. 2018;38(sup1):S32–S49. doi:10.1080/20469047.2017.1409453
61. Amare D, Negesse A, Tsegaye B, Assefa B, Ayenye B. Prevalence of undernutrition and its associated factors among children below five years of age in Bure Town, West Gojjam Zone, Amhara National Regional State, Northwest Ethiopia. *Adv Public Health*. 2016;2016:7145708. doi:10.1155/2016/7145708
62. Fekadu Y, Mesfin A, Haile D, Stoecker BJ. Factors associated with nutritional status of infants and young children in Somali Region, Ethiopia: a cross-sectional study. *BMC Public Health*. 2015;15(1):1–9. doi:10.1186/s12889-015-2190-7
63. Soboksa NE, Gari SR, Hailu AB, Mengistie Alemu B. Childhood malnutrition and the association with diarrhea, water supply, sanitation, and hygiene practices in Kersa and Omo Nada Districts of Jimma Zone, Ethiopia. *Environ Health Insights*. 2021;15:1178630221999635. doi:10.1177/1178630221999635
64. Patwari A. Diarrhoea and malnutrition interaction. *Indian J Pediatr*. 1999;66(1 Suppl):S124–34.
65. Kuehn J, Ismael Z, Long PF, Barker CI, Sharland M. Reported rates of diarrhea following oral penicillin therapy in pediatric clinical trials. *J Pediatr Pharmacol Ther*. 2015;20(2):90–104. doi:10.5863/1551-6776-20.2.90

## Journal of Multidisciplinary Healthcare

Dovepress

### Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>