

The Causal Relationship Between Gastroesophageal Reflux Disease and Chronic Obstructive Pulmonary Disease: A Bidirectional Two-Sample Mendelian Randomization Study

Bo Liu^{1,2}, Mengling Chen¹, Junjie You¹, Silin Zheng³, Min Huang²

¹School of Nursing, Southwest Medical University, Luzhou, Sichuan Province, 646000, People's Republic of China; ²Department of Respiratory and Critical Care Medicine, The Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan Province, 646000, People's Republic of China; ³Nursing Department, The Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan Province, 646000, People's Republic of China

Correspondence: Silin Zheng, Nursing Department, The Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan, 646000, People's Republic of China, Email 1400223549@qq.com; Min Huang, Department of Respiratory and Critical Care Medicine, The Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan Province, 646000, People's Republic of China, Email hm20140313@163.com

Background: Gastroesophageal reflux disease (GERD) and Chronic Obstructive Pulmonary Disease (COPD) often coexist and have been associated in observational studies. However, the real potential causal relationship between GERD and COPD is unknown and not well established.

Methods: In this study, we conducted a bidirectional two-sample Mendelian randomization(MR) to estimate whether GERD and COPD are causal. The GERD genetic data is from summary level data of a genome-wide association (GWAS) meta-analysis (Ncases = 71,522, Ncontrol=26,079). The COPD GWAS are available from the FinnGen (Ncases=16,410, Ncontrol=283,589). MR-Egger regression, Weighted Median, and Inverse-variance weighted (IVW) were used for MR analysis from the R package "TwoSampleMR", and IVW was the dominant estimation method. Additionally, the MR pleiotropy residual sum and outlier (MR-PRESSO), Cochran Q statistic, and leave-one-out analysis were used to detect and correct for the effect of heterogeneity and horizontal pleiotropy.

Results: MR analysis indicated that GERD was causally associated with an increased risk of COPD (IVW odds ratio (OR): 1.3760, 95% confidence interval (CI): 1.1565–1.6371, P=0.0003), and vice versa (IVW OR: 1.1728, 95% CI: 1.0613–1.2961, P=0.0018). The analyses did not reveal any pleiotropy or heterogeneity.

Conclusion: Our study revealed possible evidence for a bidirectional causal relationship between GERD and COPD. Implementing screening and preventive strategies for GERD in individuals with COPD, and vice versa, will be crucial in future healthcare management. Further studies are needed to elucidate the mechanisms underlying the causal relationship between GERD and COPD.

Keywords: human genetics, Mendelian randomization, gastroesophageal reflux disease, causal relationship, chronic obstructive pulmonary disease

Introduction

Chronic obstructive pulmonary disease (COPD) and Gastroesophageal reflux disease (GERD) are two common and debilitating diseases that significantly impact global health. COPD is a progressive chronic respiratory disease (CRD) characterized by persistent airflow limitation, chronic systemic inflammation, respiratory symptoms such as coughing, wheezing, and shortness of breath, and pathological changes in the airways.^{1,2} COPD has become the third leading cause of death worldwide, with high morbidity, mortality, and disability rates, and exacerbating the burden on healthcare systems and economies.^{3,4} The incidence of COPD increases with age, with smoking being the major risk factor.^{2,5} In 2019, the global prevalence of COPD was estimated to be approximately 212.3 million cases, with an additional 16.2 million new cases reported. The disease claimed the lives of 3.3 million people, making it the deadliest of all CRD.⁵ Gastroesophageal reflux disease (GERD) is a common gastrointestinal

disorder that affects approximately 13% of the world’s population,^{6,7} and 20% of adults in Western countries.⁸ In recent years, an increasing number of observational studies have shown bidirectional associations between COPD and GERD.^{9–11} It has been reported that the incidence of GERD in patients with COPD ranges from 17% to 78%.¹² In addition, a systematic review showed that the incidence of mixed reflux in patients with GERD was much higher in individuals with COPD than in those without COPD,¹³ and exacerbations of COPD have also been associated with GERD.¹⁴ While GERD and COPD are distinct clinical entities, several studies suggested a potential association between these two conditions. However, the results are inconclusive and susceptible because bias due to confounding and reverse causation cannot be excluded in observational studies. Therefore, the direction of this association and the underlying mechanisms remain to be elucidated.

This is the first study to evaluate the causal relationship between COPD and GERD. We conducted a bidirectional two-sample Mendelian randomization (MR) study using summary-level genome-wide association study (GWAS) data to evaluate the causal relationship of COPD on the risk of GERD and vice versa. MR is a genetic epidemiological approach that uses genetic variants as instrumental variables (IVs) to evaluate the causal effect of an exposure on an outcome and overcomes many of the limitations of traditional observational studies, including confounding and reverse causation.^{15,16} The bidirectional MR will help to clarify whether GERD is a risk factor for the development of COPD or whether COPD contributes to the development of GERD. This study can advance our understanding of both diseases’ pathophysiology by detecting the causal relationship between COPD and GERD. In addition, understanding the nature of this relationship is critical as it may have important clinical implications for disease management and treatment strategies.

Materials and Methods

We conducted a bidirectional two-sample MR to explore the causal relationship between COPD and GERD. This study was approved by the Ethics Committee of the Affiliated Hospital of Southwest Medical University. Because of this study relied on the publicly available summary data from the GWAS study, the requirement for informed consent was waived.

Data Sources of GERD

The GWAS summary statistics from a large meta-analysis of GERD in UK Biobank and QSkin Sun and Health Study cohorts by An et al,¹⁷ including 332,601 participants of European-descent (71,522 GERD and 261,079 controls) and can be downloaded from URL (10.6084/m9.figshare.8986589). The data of patients diagnosed with GERD were collated by data fields code in the UKBB (68,535 GERD and 250,910 controls): self-report (field ID: 20002—Noncancer illness code, self-reported Medical conditions), ICD-10 (41202—main diagnoses, 41204—secondary diagnosis), ICD-9 (41203—main diagnoses, 41205—secondary diagnosis), the Office of Population Censuses and Surveys (41200—main operative procedures; 41210—secondary operative procedures) and treatment/medicine. And GRED cases were defined by self-reported heartburn and medical records of reflux medications taken in QSkin Sun and Health Study (2987 cases and 10,169 controls)¹⁷ (Table 1).

Data Sources of COPD

The COPD GWAS summary data were obtained from FinnGen (<https://www.finnngen.fi/en>), a large-scale genomic research project developed in Finland.¹⁸ The COPD GWAS data from FinnGen included 16,410 COPD cases and 283,589 control cases. In the FinnGen project, the diagnosis of COPD was based on the International Classification of Diseases, 10th edition (ICD-10), 9th edition (ICD-9), and 8th edition (ICD-8) codes. The definition of endpoints events in the study is determined by the presence of COPD ICD codes (ICD-10:J43/44; ICD-9:491.2/492; ICD-8:491.04/492). All of the participants were of European ancestry (Table 1).

Table 1 Details of the GWASs Data

Traits	Sample Size (Cases/Controls)	Population	Consortium	Web Source
GERD	71,522/261,079	European descent	A meta-analysis of GWAS	https://doi.org/10.1038/s41467-019-11968-2 https://r8.finnngen.fi/pheno/J10_COPD
COPD	16,410/283,589	European	FinnGen	

Abbreviations: GERD, Gastroesophageal Reflux Disease; COPD, Chronic Obstructive Pulmonary Disease.

Selection of IVs

Three basic principles of MR must be met: (a) the IVs are strongly linked with exposure; (b) the IVs are not strongly linked with potential confounders, and (c) the IVs do not directly affect the outcome.¹⁶

For IVs, We chose single nuclear polymorphisms (SNPs) for GERD/COPD with a genome-wide significant threshold of $P < 5 \times 10^{-8}$. Then, we clumped the selected SNPs to avoid linkage disequilibrium (LD) with each other.¹⁹ The clumping window with stringent threshold of $r^2 < 0.001$, window size = 10,000 kb. We searched the PhenoScanner (<http://www.phenoscaner.medschl.cam.ac.uk/>) to filter out the SNPs associated with known confounders to ensure that the SNPs are independent of the confounders. In the next step, we harmonized the exposure and outcome variation effect estimates and removed any SNPs with incompatible alleles or palindromic SNPs. We did not use proxies to replace missing outcome data.²⁰ In addition, we estimated the F-statistic ($F = \beta^2 / \text{se}^2$) for each SNP to assess its power, and SNPs were removed if the $F < 10$.²¹ We only used SNPs available for all traits analyzed as IVs to maintain consistency. Ultimately, 5 SNPs were available for COPD as exposures to analyses of the causal effect on GERD, and in the reverse direction, 18 SNPs were available for GERD (Table 2).

Statistical Analysis

In this study, we conducted a bidirectional MR study to estimate the potential causal associations between GERD and COPD, and the random-effects IVW method was used as the top-choice statistical method. Then, to ensure the robustness of the main analysis, we performed sensitivity analyses. Cochran's Q test was used to assess the heterogeneity among each IV. The horizontal pleiotropy of IVs was evaluated by the MR-Egger intercept test.²² To identify potential horizontal pleiotropic outliers and to address any potential bias caused by these outliers, we used the MR-PRESSO test.²³ This test was used as a corrective measure in our analysis. In addition, we performed a leave-one-SNP-out analysis to evaluate the effect of excluding single SNP of exposure on the results of the MR study. All statistical analyses mentioned above were

Table 2 The Detailed Information of the Instrumental Variables in Each Trait

Traits	SNP	Effect Allele	Other Allele	P-value	Beta	SE	F
COPD	rs143031266	G	A	2.74E-08	0.164683	0.0296348	30.88115523
	rs2273500	C	T	2.19E-11	0.0951338	0.0142143	44.79388539
	rs2904259	C	T	1.97E-08	0.0664499	0.0118358	31.52052393
	rs66694560	A	G	5.19E-10	0.0763852	0.0122937	38.60585883
	rs9270664	A	G	1.25E-08	0.067614	0.0118783	32.40146741
GERD	rs111472920	T	G	4.47E-08	0.0987	0.018	30.06694444
	rs12706746	A	G	4.24E-09	0.0369	0.0063	34.30612245
	rs12939066	T	C	3.13E-11	0.0415	0.0062	44.80359001
	rs13167137	T	G	2.82E-08	-0.0328	0.0059	30.90606148
	rs1473115	T	C	3.95E-10	0.0396	0.0063	39.51020408
	rs15071	T	C	1.01E-08	-0.0428	0.0075	32.56604444
	rs1858828	T	G	3.23E-08	0.0327	0.0059	30.71789716
	rs3072	T	C	1.86E-08	-0.0345	0.0061	31.98736899
	rs62046253	T	C	1.30E-08	-0.0352	0.0062	32.23309053
	rs62442944	T	G	3.49E-08	0.0418	0.0076	30.25
	rs6683411	A	G	4.82E-08	0.0324	0.0059	30.15685148
	rs6710685	T	C	1.96E-09	0.0391	0.0065	36.18485207
	rs6762606	T	C	4.18E-08	-0.0359	0.0065	30.5043787
	rs6809836	A	G	3.70E-09	0.0383	0.0065	34.71928994
	rs6991878	T	C	2.71E-08	-0.0338	0.0061	30.70249933
	rs72704785	A	G	6.81E-09	0.0459	0.0079	33.7575709
	rs7552188	T	C	1.07E-08	0.0413	0.0072	32.90297068
	rs769671	T	C	9.28E-10	-0.0385	0.0063	37.34567901

Abbreviations: SNP, single-nucleotide polymorphism; Beta, the per-allele effect on each trait; SE, standard error; F, F-statistic.




Figure 1 Estimated causal effect of GERD on COPD using different MR methods.

Abbreviations: SNP, single-nucleotide polymorphism; pval, p value; OR, Odds Ratio; CI, Confidence Interval.

two-sided and conducted using the TwoSampleMR packages in R 4.2.3. $P < 0.05$ were considered statistically significant.²⁴

Results

The Causal Effect of GERD on COPD

For the forward MR analysis, 1 SNP rs11171710 associated with emphysema was excluded. 4 SNPs (rs7590354, rs2271856, rs2927089, and rs1050458) were not found in FinnGen of COPD GWAS data. 2 SNPs (rs10228350; rs2108959) for being palindromic were excluded after the harmonization process. Finally, 18 eligible SNPs were selected as IVs of GERD as the exposure and COPD as the outcome, and the F statistic of all SNP was greater than 10 (Table 2). The study of the main result showed IVW (OR: 1.3760, 95% CI: 1.1565–1.6371, $P=0.0003$) and indicated that genetic predictions for GERD showed a significant association with the incidence of COPD, increasing the risk by 37.6%. MR-egger (OR: 2.2620, 95% CI: 0.7150–7.1565, $P=0.1839$), weight median (OR: 1.2800, 95% CI: 1.0117–1.6195, $P=0.0397$). (Figures 1 and S1A) There was no significant bias in the causal effect of GERD on COPD in the sensitivity analysis (all P values >0.05 for Cochran's Q test; MR-egger intercept test: Intercept = -0.0191 , $SE=0.0223$, $P=0.4047$; MR-PRESSO global test $P=0.3400$) (Table 3). The leave-one-SNP-out analysis showed that the effect estimates were not affected by the presence of a single SNP (Figure S1B).

The Causal Effect of COPD on GERD

For the reverse-direction MR analysis, 1 SNP rs708461 for being palindromic was excluded after the harmonization process. The trait of SNP rs8040868 and SNP rs113623975 were associated with smoking and were eliminated, which was the potential confounder of GERD.⁸ 1 SNP rs28929474 was the outlier MR-PRESSO detected and was excluded. Finally, 5 eligible SNPs remained as IVs of COPD as the exposure and GERD as the outcome. The F statistic of all remained SNP is greater than 10 (Table 2). In the IVW mode analysis, we found that COPD had a significant causal risk effect on GERD (OR: 1.1728, 95% CI: 1.0613–1.2961, $P=0.0018$). MR-egger (OR: 1.4337, 95% CI: 0.8934–2.3009, $P=0.2323$), weight median (OR: 1.1754, 95% CI: 1.0451–1.3219, $P=0.0070$) (Figures 2 and S2A). There was no significant bias in the causal effect of COPD on GERD in the sensitivity analysis (all P values >0.05 for Cochran's Q test; MR-egger intercept test: Intercept = -0.0160 , $SE=0.0188$, $P=0.4564$; MR-PRESSO global test $P=0.2673$) (Table 3). The leave-one-SNP-out analysis did not find any high-impact points (Figure S2B).

Table 3 Sensitivity Analysis of MR

Exposure	Outcome	nSNPs	Heterogeneity Test		MR-Egger Pleiotropy Test		MR-PRESSO Global Outlier Test		
			Q	P-value	Intercept	P-value	RSSobs	P-value	Outlier
GERD	COPD	18	19.275370	0.3129	-0.0191	0.4047	21.77467	0.3400	None
COPD	GERD	6	14.371421	0.0134	0.0141	0.2562	24.56575	0.0367	rs28929474
After remove the outlier	GERD	5	5.964387	0.2018	-0.0160	0.4564	9.919114	0.2673	None

Abbreviations: GERD, Gastroesophageal Reflux Disease; COPD, Chronic Obstructive Pulmonary Disease; SNP, single-nucleotide polymorphism; MR-PRESSO, MR pleiotropy residual sum and outlier.




Figure 2 Estimated causal effect of COPD on GERD using different MR methods.

Abbreviations: SNP, single-nucleotide polymorphism; pval, p value; OR, Odds Ratio; CI, Confidence Interval.

Discussion

To our knowledge, this is the first bidirectional MR study to evaluate the causal association between COPD and GERD. The results of our MR study may provide evidence supporting that genetically predicted COPD was positively associated with the risk of GERD and vice versa. The sensitivity analyses conducted in this study consistently and robustly supported the causal association between COPD and GERD. Although MR-egger results do not support a causal association between GERD and COPD, the results are in the same direction as IVW. Some researchers believe that if the IVW method produces a significant result, even if the results of the other methods are not significant and there is no evidence of pleiotropy and heterogeneity, this result can be considered positive as long as the beta values of the other methods point in the same direction.^{25–27}

For the forward MR analysis, our results showed a significant positive causal association of GERD with COPD risk, consistent with previous study findings. In this case, the presence of GERD-associated genetic variants may causally associated with an elevated risk of developing COPD. This finding implies that the genetic factors influencing GERD contribute, at least in part, to the development of COPD. García Rodríguez LA et al¹⁰ used the UK General Practice Research Database to perform a study that found that patients diagnosed with GERD had a 17% increased risk of developing COPD. Faulty anti-reflux barriers cause GERD, one of the primary mechanisms.²⁸ Current perspectives and some studies show that GERD may contribute to the progression of COPD by inhaling gastric contents, acid reflux, leading to bronchoconstriction and airway inflammation and potentially causing or exacerbating COPD.^{29–32} These findings could have significant implications for the clinical management of individuals with GERD, highlighting the importance of early diagnosis and treatment of GERD to potentially reduce the risk of subsequent development of COPD.

Conversely, in the reversal MR analysis, we also discovered that COPD has a potential causal association with GERD risk. Current extensive research indicates a relatively high prevalence of GERD in COPD patients.^{12,33,34} Furthermore, the coexistence of GERD in patients with COPD appears to be associated with worsened pulmonary symptoms, decreased quality of life, and increased frequency of COPD exacerbations.^{35–38} A comprehensive systematic review and meta-analysis showed that GERD was a significant risk factor for exacerbations of COPD (OR: 5.37; 95% CI 2.71–10.64).³⁹ According to Rascon-Aguilar et al,⁴⁰ patients with GERD-positive COPD were twice as likely to experience exacerbations as patients without GERD. Patients with COPD experience significant lung function impairment and reduced lung compliance, which can lead to dyspnea, breathing way changes, and respiratory muscle fatigue, which increases intrathoracic pressure.²⁶ Elevated intrathoracic pressure can increase intra-abdominal pressure, resulting in decreased pressure on the lower esophageal sphincter (LES), causing reflux of gastric acid and digestive fluid into the esophagus.^{29,41} In addition, patients with COPD require long-term use of respiratory medications such as inhaled bronchodilators, anticholinergics, and corticosteroids, which may increase the risk of developing GERD.^{31,42,43} Obviously, the mechanisms described in the previous paragraph are closely related to this section, often resulting in a vicious cycle. However, because GERD is often silent and imperceptible in patients with COPD, it is may always overlooked as a potential cause of respiratory illness and symptoms.⁴⁴ Therefore, screening for GERD in patients with COPD is necessary.

As mentioned above, the underlying mechanisms of the interaction between GERD and COPD remain unclear. Based on our research, we speculate on possible mechanisms underlying the association between these two diseases. rs15071 is located on chromosome 5, in the 3' UTR of the VCAN gene. The VCAN gene encodes a proteoglycan called versican in the extracellular matrix (ECM). A study conducted by Liao et al⁴⁵ found a significant association between the VCAN gene and lung function through dmGWAS network analysis and validation. It has been reported that versican expression

negatively correlates with FEV1 in human alveolar walls and rims.⁴⁶ Versican has been implicated in the tissue remodeling process in the lungs of COPD patients. A study conducted by Wu et al⁴⁷ demonstrated that inhibition of versican synthesis may be a potential therapeutic strategy to increase the deposition of insoluble elastic proteins and stimulate elastic fiber repair in the lungs of COPD patients. In addition, COPD patients are known to have persistent chronic inflammation. Versican plays a critical role in the regulation of immune and inflammatory responses.^{48,49} Versican can interact with inflammatory cells indirectly or directly; these interactions activate signaling pathways that promote the synthesis and secretion of inflammatory cytokines such as TNF- α , IL-6, and NF- κ B.^{49,50}

In addition, we identified rs6683411 located on chromosome 1 in the intergenic region of the hnRNPA1P46 gene. Heterogeneous nuclear ribonucleoprotein A/B (hnRNPA/B) is a core member of the hnRNP family of RNA-binding proteins (RBPs), which includes four major subtypes, A0, A1, A2/B1, and A3.⁵¹ Previous studies suggest that hnRNPs were involved in and induce epithelial-mesenchymal transition (EMT), a process closely associated with the development of COPD and a potential mechanism leading to small airway fibrosis.^{52–55} rs6762606 is located on chromosome 3, in the intronic region of the EPHB1 gene. As mentioned above, the study by Liao et al⁴⁵ also clearly shows a significant association between the EPHB1 gene and lung function. rs72704785 on chromosome 5 is located in the upstream promoter region of the miR-4456 gene. Existing research suggests that miR-4456 may potentially inhibit the tight junction(TJ) damage caused by tobacco exposure in COPD through its interaction with chemokine ligands.⁵⁶

Our study has some strengths. First, the MR design allowed us to assess causality by using genetic variants as IVs, thereby reducing the effects of confounding and reverse causation. Second, we used large-scale GWAS to obtain robust genetic instruments and summary statistics for COPD and GERD. Third, bidirectional MR analysis provided insight into the directionality of the causal relationship between COPD and GERD. Of course, there are some potential limitations. First, the prior observation studies have explored this relationship, so our research findings may have a limited impact on advancing the development of this field. Second, as with all MR studies, our results depend on certain assumptions, including that the genetic variants used as IVs are valid instruments and that there is no pleiotropy. Although we conducted sensitivity analyses to assess the robustness of our results, the possibility of unmeasured pleiotropy or violation of these assumptions cannot be completely ruled out. Third, the generalizability of our findings may be limited to populations of European ancestry, as the genetic tools were primarily derived from studies conducted in these populations. Finally, additional subgroup analyses were not possible due to the use of summary statistics rather than raw data in the study.

In conclusion, our bidirectional two-sample Mendelian randomization study may provide potential evidence for a causal association between COPD and an increased risk of GERD. These findings have important clinical implications, emphasizing the need for clinicians to consider both diseases simultaneously in their management strategies and the importance of early intervention and management. Further research is warranted to elucidate the underlying mechanisms linking GERD and COPD, explore potential therapeutic interventions, and investigate this relationship's bidirectional nature in different populations.

Conclusion

This study represents the first MR study to investigation of the causal relationship between GERD and COPD. Our comprehensive MR analysis revealed possible evidence for supporting a bidirectional causal relationship between GERD and COPD. By using genetic variants as IVs, we overcame confounding factors and gained a more robust understanding of the causal nature of this relationship. Further research is warranted to delve deeper into the mechanisms underlying this bidirectional causality and explore potential therapeutic interventions that could improve outcomes for individuals affected by GERD and COPD.

Abbreviations

GERD, Gastroesophageal reflux disease; COPD, Chronic Obstructive Pulmonary Disease; GWAS, genome-wide association study; MR, Mendelian randomization; IVW, Inverse-variance weighted; MR-PRESSO, MR pleiotropy residual sum and outlier; OR, odds ratio; CI, confidence interval; CRD, chronic respiratory disease; IVs, instrumental variables; SNPs, single nuclear polymorphisms; LD, linkage disequilibrium; SE, standard error; LES, lower esophageal sphincter; ECM, extracellular matrix; RBPs, RNA-binding proteins; EMT, epithelial-mesenchymal transition; TJ, tight junction.

Data Sharing Statement

All data generated or analysed during this study are included in this published article and its [Supplementary Information Files](#).

Ethics Approval and Consent to Participate

This study was approved by the Ethics Committee of the Affiliated Hospital of Southwest Medical University. Because of this study relied on the publicly available summary data from the GWAS study, the requirement for informed consent was waived.

Acknowledgments

We want to acknowledge the participants and investigators of the FinnGen study and the participants of all GWAS cohorts included in the present work.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis, and interpretation, or all these areas; took part in drafting, revising, or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

There is no funding to report.

Disclosure

The authors declare that they have no competing interests.

References

1. Divo MJ, Martinez CH, Mannino DM. Ageing and the epidemiology of multimorbidity. *Eur Respir J*. 2014;44(4):1055–1068. doi:10.1183/09031936.00059814
2. Lareau SC, Fahy B, Meek P, et al. Chronic Obstructive Pulmonary Disease (COPD). *Am J Respir Crit Care Med*. 2019;199(1):1–P2. doi:10.1164/rccm.1991P1
3. Ko FW, Chan KP, Hui DS, et al. Acute exacerbation of COPD. *Respirology*. 2016;21(7):1152–1165. doi:10.1111/resp.12780
4. Raheison C, Girodet PO. Epidemiology of COPD. *Eur Respir Rev*. 2009;18(114):213–221. doi:10.1183/09059180.00003609
5. Momtazmanesh S, Moghaddam SS, Ghamari SH, et al. Global burden of chronic respiratory diseases and risk factors, 1990–2019: an update from the global burden of disease study 2019. *E Clin Med*. 2023;59:101936.
6. Yuan S, Larsson SC. Adiposity, diabetes, lifestyle factors and risk of gastroesophageal reflux disease: a Mendelian randomization study. *Eur J Epidemiol*. 2022;37(7):747–754. doi:10.1007/s10654-022-00842-z
7. Richter JE, Rubenstein JH. Presentation and epidemiology of gastroesophageal reflux disease. *Gastroenterology*. 2018;154(2):267–276. doi:10.1053/j.gastro.2017.07.045
8. Maret-Ouda J, Markar SR, Lagergren J. Gastroesophageal reflux disease. *JAMA*. 2020;324(24):2565. doi:10.1001/jama.2020.21573
9. el-Serag HB, Sonnenberg A. Comorbid occurrence of laryngeal or pulmonary disease with esophagitis in United States military veterans. *Gastroenterology*. 1997;113(3):755–760. doi:10.1016/S0016-5085(97)70168-9
10. García Rodríguez LA, Ruigómez A, Martín-Merino E, et al. Relationship between gastroesophageal reflux disease and COPD in UK primary care. *Chest*. 2008;134(6):1223–1230. doi:10.1378/chest.08-0902
11. Menon S, Nightingale P, Trudgill N. Chronic obstructive pulmonary disease and the risk of esophagitis, barrett's esophagus, and esophageal adenocarcinoma: a primary care case-control study. *J Clin Gastroenterol*. 2019;53(10):e451–e5. doi:10.1097/MCG.0000000000001215
12. Lee AL, Goldstein RS. Gastroesophageal reflux disease in COPD: links and risks. *Int J Chron Obstruct Pulmon Dis*. 2015;10:1935–1949. doi:10.2147/COPD.S77562
13. El-Serag HB, Sweet S, Winchester CC, et al. Update on the epidemiology of gastro-oesophageal reflux disease: a systematic review. *Gut*. 2014;63(6):871–880. doi:10.1136/gutjnl-2012-304269
14. Sakae TM, Pizzichini MM, Teixeira PJ, et al. Exacerbations of COPD and symptoms of gastroesophageal reflux: a systematic review and meta-analysis. *J Bras Pneumol*. 2013;39(3):259–271. doi:10.1590/S1806-37132013000300002
15. Emdin CA, Khera AV, Kathiresan S. Mendelian Randomization. *JAMA*. 2017;318(19):1925–1926. doi:10.1001/jama.2017.17219
16. Davies NM, Holmes MV, Davey Smith G. Reading Mendelian randomisation studies: a guide, glossary, and checklist for clinicians. *BMJ*. 2018;362:k601. doi:10.1136/bmj.k601
17. An J, Gharahkhani P, Law MH, et al. Gastroesophageal reflux GWAS identifies risk loci that also associate with subsequent severe esophageal diseases. *Nat Commun*. 2019;10(1):4219. doi:10.1038/s41467-019-11968-2

18. Kurki MI, Karjalainen J, Palta P, et al. FinnGen provides genetic insights from a well-phenotyped isolated population. *Nature*. 2023;613(7944):508–518. doi:10.1038/s41586-022-05473-8
19. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports systematic causal inference across the human phenotype. *eLife*. 2018;7. doi:10.7554/eLife.34408
20. Zhou W, Cai J, Li Z, et al. Association of atopic dermatitis with autoimmune diseases: a bidirectional and multivariable two-sample Mendelian randomization study. *Front Immunol*. 2023;14:1132719. doi:10.3389/fimmu.2023.1132719
21. Pierce BL, Ahsan H, Vanderweele TJ. Power and instrument strength requirements for Mendelian randomization studies using multiple genetic variants. *Int J Epidemiol*. 2011;40(3):740–752. doi:10.1093/ije/dyq151
22. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol*. 2015;44(2):512–525. doi:10.1093/ije/dyv080
23. Verbanck M, Chen CY, Neale B, et al. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet*. 2018;50(5):693–698. doi:10.1038/s41588-018-0099-7
24. Hemani G, Tilling K, Davey Smith G, Li J. Orienting the causal relationship between imprecisely measured traits using GWAS summary data. *PLoS Genet*. 2017;13(11):e1007081. doi:10.1371/journal.pgen.1007081
25. Zagkos L, Dib MJ, Pinto R, et al. Associations of genetically predicted fatty acid levels across the phenotype: a Mendelian randomisation study. *PLoS Med*. 2022;19(12):e1004141. doi:10.1371/journal.pmed.1004141
26. Chen X, Kong J, Diao X, et al. Depression and prostate cancer risk: a Mendelian randomization study. *Cancer Med*. 2020;9(23):9160–9167. doi:10.1002/cam4.3493
27. Wang S, Zhu H, Pan L, et al. Systemic inflammatory regulators and risk of acute-on-chronic liver failure: a bidirectional Mendelian-randomization study. *Front Cell Dev Biol*. 2023;11:1125233. doi:10.3389/fcell.2023.1125233
28. Xie C, Wang J, Li Y, et al. Esophagogastric junction contractility integral reflect the anti-reflux barrier dysfunction in patients with gastroesophageal reflux disease. *J Neurogastroenterol Motil*. 2017;23(1):27–33. doi:10.5056/jnm16008
29. Houghton LA, Lee AS, Badri H, et al. Respiratory disease and the oesophagus: reflux, reflexes and microaspiration. *Nat Rev Gastroenterol Hepatol*. 2016;13(8):445–460. doi:10.1038/nrgastro.2016.91
30. Wedzicha JA, Brill SE, Allinson JP, et al. Mechanisms and impact of the frequent exacerbator phenotype in chronic obstructive pulmonary disease. *BMC Med*. 2013;11(1):181. doi:10.1186/1741-7015-11-181
31. Lee AS, Lee JS, He Z, et al. Reflux-aspiration in chronic lung disease. *Ann Am Thorac Soc*. 2020;17(2):155–164. doi:10.1513/AnnalsATS.201906-427CME
32. Chen Z, Sun L, Chen H, et al. Dorsal vagal complex modulates neurogenic airway inflammation in a guinea pig model with esophageal perfusion of HCl. *Front Physiol*. 2018;9:536. doi:10.3389/fphys.2018.00536
33. Chiu YC, Chang WP, Tang GJ, et al. Chronic obstructive pulmonary disease is associated with a higher risk of functional gastrointestinal disorders. *Respir Med*. 2022;197:106833. doi:10.1016/j.rmed.2022.106833
34. Kim J, Lee JH, Kim Y, et al. Association between chronic obstructive pulmonary disease and gastroesophageal reflux disease: a national cross-sectional cohort study. *BMC Pulm Med*. 2013;13(1):51. doi:10.1186/1471-2466-13-51
35. Shirai T, Mikamo M, Tsuchiya T, et al. Real-world effect of gastroesophageal reflux disease on cough-related quality of life and disease status in asthma and COPD. *Allergol Int*. 2015;64(1):79–83. doi:10.1016/j.alit.2014.08.001
36. Tsai CL, Lin YH, Wang MT, et al. Gastro-oesophageal reflux disease increases the risk of intensive care unit admittance and mechanical ventilation use among patients with chronic obstructive pulmonary disease: a nationwide population-based cohort study. *Crit Care*. 2015;19(1):110. doi:10.1186/s13054-015-0849-1
37. Takada K, Matsumoto S, Hiramatsu T, et al. F スケール質問票を用いた胃食道逆流症の評価と COPD との関連性の検討 [Relationship between chronic obstructive pulmonary disease and gastroesophageal reflux disease defined by the frequency scale for the symptoms of gastroesophageal reflux disease]. *Nihon Kokyuki Gakkai Zasshi*. 2010;48(9):644–648. Japanese
38. Liang B, Wang M, Yi Q, et al. Association of gastroesophageal reflux disease risk with exacerbations of chronic obstructive pulmonary disease. *Dis Esophagus*. 2013;26(6):557–560. doi:10.1111/dote.12014
39. Huang C, Liu Y, Shi G. A systematic review with meta-analysis of gastroesophageal reflux disease and exacerbations of chronic obstructive pulmonary disease. *BMC Pulm Med*. 2020;20(1):2. doi:10.1186/s12890-019-1027-z
40. Rascon-Aguilar IE, Pamer M, Wludyka P, et al. Role of gastroesophageal reflux symptoms in exacerbations of COPD. *Chest*. 2006;130(4):1096–1101. doi:10.1378/chest.130.4.1096
41. Harding SM, Allen JE, Blumin JH, et al. Respiratory manifestations of gastroesophageal reflux disease. *Ann N Y Acad Sci*. 2013;1300(1):43–52. doi:10.1111/nyas.12231
42. Mungan Z, Pınarbaşı Şimşek B. Which drugs are risk factors for the development of gastroesophageal reflux disease? *Turk J Gastroenterol*. 2017;28:S38–s43.
43. Del Grande LM, Herbella FA, Bigatao AM, et al. Inhaled beta agonist bronchodilator does not affect trans-diaphragmatic pressure gradient but decreases lower esophageal sphincter retention pressure in patients with Chronic Obstructive Pulmonary Disease (COPD) and Gastroesophageal Reflux Disease (GERD). *J Gastrointest Surg*. 2016;20(10):1679–1682. doi:10.1007/s11605-016-3192-1
44. Griffiths TL, Nassar M, Soubani AO. Pulmonary manifestations of gastroesophageal reflux disease. *Expert Rev Respir Med*. 2020;14(8):767–775. doi:10.1080/17476348.2020.1758068
45. Liao SY, Lin X, Christiani DC. Genome-wide association and network analysis of lung function in the Framingham heart study. *Genet Epidemiol*. 2014;38(6):572–578. doi:10.1002/gepi.21841
46. Merrilees MJ, Ching PS, Beaumont B, et al. Changes in elastin, elastin binding protein and versican in alveoli in chronic obstructive pulmonary disease. *Respir Res*. 2008;9(1):41. doi:10.1186/1465-9921-9-41
47. Wu L, Zhang J, Qu JM, et al. Deposition of insoluble elastin by pulmonary fibroblasts from patients with COPD is increased by treatment with versican siRNA. *Int J Chron Obstruct Pulmon Dis*. 2017;12:267–273. doi:10.2147/COPD.S116217
48. Chang MY, Kang I, Gale M, et al. Versican is produced by Trif- and type I interferon-dependent signaling in macrophages and contributes to fine control of innate immunity in lungs. *Am J Physiol Lung Cell Mol Physiol*. 2017;313(6):L1069–L1086. doi:10.1152/ajplung.00353.2017

49. Wight TN, Kang I, Evanko SP, et al. Versican-A critical extracellular matrix regulator of immunity and inflammation. *Front Immunol.* **2020**;11:512. doi:10.3389/fimmu.2020.00512
50. Andersson-Sjöland A, Hallgren O, Rolandsson S, et al. Versican in inflammation and tissue remodeling: the impact on lung disorders. *Glycobiology.* **2015**;25(3):243–251. doi:10.1093/glycob/cwu120
51. Lu Y, Wang X, Gu Q, et al. Heterogeneous nuclear ribonucleoprotein A/B: an emerging group of cancer biomarkers and therapeutic targets. *Cell Death Discov.* **2022**;8(1):337. doi:10.1038/s41420-022-01129-8
52. Sohal SS. Epithelial and endothelial cell plasticity in chronic obstructive pulmonary disease (COPD). *Respir Investig.* **2017**;55(2):104–113. doi:10.1016/j.resinv.2016.11.006
53. Sohal SS, Mahmood MQ, Walters EH. Clinical significance of epithelial mesenchymal transition (EMT) in chronic obstructive pulmonary disease (COPD): potential target for prevention of airway fibrosis and lung cancer. *Clin Transl Med.* **2014**;3(1):33. doi:10.1186/s40169-014-0033-2
54. Nowrin K, Sohal SS, Peterson G, et al. Epithelial-mesenchymal transition as a fundamental underlying pathogenic process in COPD airways: fibrosis, remodeling and cancer. *Expert Rev Respir Med.* **2014**;8(5):547–559. doi:10.1586/17476348.2014.948853
55. Morrow JD, Cho MH, Platig J, et al. Ensemble genomic analysis in human lung tissue identifies novel genes for chronic obstructive pulmonary disease. *Hum Genomics.* **2018**;12(1):1. doi:10.1186/s40246-018-0132-z
56. Yu W, Ye T, Ding J, et al. miR-4456/CCL3/CCR5 pathway in the pathogenesis of tight junction impairment in chronic obstructive pulmonary disease. *Front Pharmacol.* **2021**;12:551839. doi:10.3389/fphar.2021.551839

International Journal of Chronic Obstructive Pulmonary Disease

Dovepress

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols. This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>