

SOURCES OF THE DATA-BASES INCLUDED IN THE MANUSCRIPT: *Alpha-1 antitrypsin Pi\*Z gene frequency and Pi\*ZZ genotype numbers worldwide: an update. International Journal of Chronic Obstructive Pulmonary Disease.* (Submission ID: 125389)

#### GENERAL REVIEW ARTICLES

**Hutchison DC.** Alpha 1-antitrypsin deficiency in Europe: geographical distribution of Pi types S and Z. *Respir Med.* 1998;92:367-77.

**Blanco I,** E Fernández, EF Bustillo. Alpha-1-antitrypsin PI phenotypes S and Z in Europe: an analysis of the published surveys. *Clin Genet* 2001;60:31-41.

**Blanco I,** EF Bustillo, MC Rodríguez. Distribution of  $\alpha$ 1-antitrypsin PI S and PI Z frequencies in countries outside Europe: a meta-analysis. *Clin Genet* 2001;60:431-44.

**Miravittles M.** Alpha1-anti-trypsin deficiency: epidemiology and incidence. *Respir Med.* 2002;96:205.

**de Serres FJ.** Worldwide racial and ethnic distribution of alpha1-antitrypsin deficiency: summary of an analysis of published genetic epidemiologic surveys. *Chest.* 2002;122:1818-29.

**Luisetti M,** Seersholm N. Alpha1-antitrypsin deficiency. 1: epidemiology of alpha1-antitrypsin deficiency. *Thorax.* 2004;59:164-9.

**de Serres FJ,** Blanco I, Fernández-Bustillo E. Health implications of alpha1-antitrypsin deficiency in Sub-Sahara African countries and their emigrants in Europe and the New World. *Genet Med.* 2005;7:175-84.

**Blanco I,** de Serres FJ, Fernandez-Bustillo E, Lara B, Miravittles M. Estimated numbers and prevalence of PI\*S and PI\*Z alleles of alpha1-antitrypsin deficiency in European countries. *Eur Respir J.* 2006;27:77-84.

**de Serres FJ,** Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI\*S and PI\*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. *Eur Respir J.* 2006;28:1091-9.

**de Serres FJ,** Blanco I. Prevalence of  $\alpha$ 1-antitrypsin deficiency alleles PI\*S and PI\*Z worldwide and effective screening for each of the five phenotypic classes PI\*MS, PI\*MZ, PI\*SS, PI\*SZ, and PI\*ZZ: a comprehensive review. *Ther Adv Respir Dis.* 2012;6:277-95.

**Blanco I,** de Serres FJ, Cárcaba V, Lara B, Fernández-Bustillo E. Alpha-1 Antitrypsin Deficiency PI\*Z and PI\*S Gene Frequency Distribution Using on Maps of the World by an Inverse Distance Weighting (IDW) Multivariate Interpolation Method. *Hepat Mon.* 2012;12(10 HCC):e7434.

## SPECIFIC ARTICLES FOR DIFFERENT COUNTRIES ORDERED ALPHABETICALLY.

(Selected studies are shown in blue. Rejected studies, in red and italics)

### **Afghanistan**

*Spitsyn VA, Novoradovskii AG, Spitsyna NKh, Parik Iula. [Polymorphism of alpha1-antitrypsin in Pamir populations. Reproductive compensation--possible mechanism for maintaining genetic diversity gor PI genes in human populations]. Genetika 1989; 25: 2218–24.*

*Serum protein polymorphisms in four populations of Afghanistan. Rahimi AG, Goedde HW, Flatz G, Kaifje S, Benkmann HG, Delbrück H. Am J Hum Genet. 1977;29:356-60*

*Goedde HW, Benkmann HG, Flatz G, Bienzle U, Kroeger A. alpha 1-antitrypsin subtypes in the populations of Germany, Ecuador, Afghanistan, Cameroon, and Saudi-Arabia. Zeitschrift fur Morphologie und Anthropologie 1980;70:341–6.*

### **Albania**

*Scheil HG, Schuhen S, Huckenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (PI) in the Balkans. Coll Antropol. 2002;26:403-10.*

### **Angola**

*Dunn DS, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. Hum Hered 1986;36:238 –42.*

### **Argentina**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

*Sorroche PB, Fernández Acquier M, López Jove O, Giugno E, Pace S, Livellara B, Legal S, Oyhamburu J, Saez MS. Alpha-1 Antitrypsin Deficiency in COPD Patients: A Cross-Sectional Study. Arch Bronconeumol. 2015;51:539-43.*

### **Australia**

**Ananthakrishnan R**, Biegler B, Dennis PM. Alpha 1-antitrypsin phenotypes in paraproteinaemias. Lancet. 1979;1:561-2.

**Mulley JC**. Distribution of the six alpha 1-antitrypsin (PI) subtypes from a sample of blood donors. Aust J Exp Biol Med Sci 1980; 58: 409–14.

**McPhee H**, Ananthakrishnan R, Taft LI. Antiproteases and Down's syndrome in an Australian population. J Med Genet 1980; 17: 170–3.

**Mulley JC.** Distribution of alpha 1-antitrypsin (Pi) subtypes from newborns. *Aust J Exp Biol Med Sci* 1982; 60: 207–9.

**Clark P.** Alpha-1-protease inhibitor phenotypes in Australia. *Hum Hered* 1982;32:225–7.

**Oakeshott JG,** Muir A, Clark P et al. Effects of the protease inhibitor (Pi) polymorphism on alpha-1-antitrypsin concentration and elastase inhibitory capacity in human serum. *Ann Human Biol* 1985; 12: 149–60.

**Muir A, Whitfield JB.** Alpha-1-antitrypsin (protease inhibitor) phenotypes and longevity. *Hum Hered* 1990; 40: 22–8.

**Huggard PR,** West MJ, Summers KM. Alpha 1-antitrypsin deficiency alleles and blood pressure in an Australian population. *Clin Exp Pharmacol Physiol* 1996; 23: 600–1.

**de Serres FJ,** Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. *Clin Genet* 2003; 64: 382-97.

*Clark P. Alpha-1-protease inhibitor phenotypes in Australia. Hum Hered. 1982;32:225-7.*

#### **Austria**

**Böhme A,** Cleve H, Schönitzer D, Reissigl H, Kazda S, Müller W. alpha 1-Antitrypsin (Pi) types and subtypes in the Tyrolean population. *Hum Genet.* 1983;63:193-4.

#### **Belgium**

**Kimpen J,** Legius E, Bosmans E, et al. Alpha-1-proteinase inhibitor gene frequencies in Belgium. *Gene Geogr* 1990; 4:159–63.

#### **Bolivia**

**de Serres FJ,** Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

#### **Bosnia-Herzegovina**

**Kozarevic D,** Laban M, Budimir M, Vojvodic N, Roberts A, Gordon T, McGee DL. Intermediate alpha1-antitrypsin deficiency and chronic obstructive pulmonary disease in Yugoslavia. *Am Rev Respir Dis.* 1978;117:1039-43.

#### **Botswana**

*Dunn DS, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. Hum Hered 1986;36:238 –42.*

## **Brazil**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

*Simoës AL, Kompf J, Ritter H, Luckenbach C, Zischler H, Salzano FM. Electrophoretic and isoelectric focusing studies in Brazilian Indians: data on four systems. Hum Biol 1989; 61: 427-38.*

## **Cameroon**

**Goedde HW**, Benkmann HG, Flatz G, Bienzle U, Kroeger A. alpha 1-antitrypsin subtypes in the populations of Germany, Ecuador, Afghanistan, Cameroon, and Saudi-Arabia. *Zeitschrift fur Morphologie und Anthropologie* 1980;70:341–46.

## **Canada**

**Ashley MJ**, Corey P, Chan-Yeung M. Smoking, dust exposure, and serum alpha 1-antitrypsin. *Am Rev Respir Dis* 1980;121: 783–88.

**Chan-Yeung M**, Ashley MJ, Corey P et al. Pi phenotypes and the prevalence of chest symptoms and lung function abnormalities in workers employed in dusty industries. *Am Rev Respir Dis* 1978: 117: 239–45.

**Horne SL**, Clarke ME, Barnett GD. Alpha-1-antitrypsin deficiency: Its effect on pulmonary function in Saskatchewan grain buyers. *Occupational Pulmonary Disease: Focus on Grain Dust and Health*. New York: Academic Press, 1980: 347–56.

**Moroz SP**, Cutz E, Cox DW et al. Liver disease associated with alpha1-antitrypsin deficiency in childhood. *J Pediatr* 1976: 88:19–25.

**Talamo RC**, Thurlbeck WM. Alpha-antitrypsin Pi types in postmortem blood. *Am Rev Respir Dis* 1975;112:201–7.

**Singh R**, Horne SL, Gerrard JW et al. Alpha 1-antitrypsin phenotypes in Canadian Metis. *Hum Hered* 1983;33:181–3.

**Cox DW**, Simpson NE, Jantti R. Group-specific component, alpha1-antitrypsin and esterase D in Canadian Eskimos. *Hum Hered* 1978;28:341–50.

**Joly J**, Richer G, Boisvert F et al. alpha-1-Antitrypsin phenotypes in French Canadian newborns. *Hum Hered* 1980;30:1–2.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. *Clin Genet* 2003;64:382-97.

## **Cape Verde**

**Spínola C**, Brehm A, Spínola H. Alpha-1-antitrypsin deficiency in the Cape Verde islands (Northwest Africa): High prevalence in a sub-Saharan population. *Respir Med*. 2010;104:1069-72.

## **Chile**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

*Goedde HW, Rothhammer F, Benkmann HG, Bogdanski P. Ecogenetic studies in Atacameno Indians. Hum Genet 1984; 67:343-6.*

## **China**

**Saha N**. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. *Ann Hum Biol* 1990; 17: 229–34.

**Ying QL**, Liang ZQ. Allelic frequencies of plasma alpha-1-antitrypsin in Chinese. *Sci Sin [B]* 1984;27: 161–8.

**Ying QL**, Zhang ML, Liang CC, et al. Alpha-1-antitrypsin types in five Chinese national minorities. *Hum Genet* 1985;71: 225–6.

**Xu JJ**, Cui MY, Chen LZ, et al. Polymorphisms of Pi, Hp, ADA and AK in Mongolian, Korean and Zhuang populations of China. *Ann Hum Biol* 1986;13:245–51.

**Lee SS**, Lawton JW, Ko KH, Lam KM, Lin CK. alpha-1 antitrypsin phenotypes by isoelectric focusing in a metropolitan southern Chinese population. *J Clin Pathol*. 2001;54:798-800.

**Zhu YJ**. Epidemiological survey of chronic obstructive pulmonary disease and alpha-1-deficiency in China. *Respirology*. 2001;6 Suppl:S13-5.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI\*S and PI\*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. *Eur Respir J*. 2006;28:1091-9.

## **Colombia**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

*Kirk RL, McDermid EM, Blake NM, Gajdusek DC, Leyshon WC, MacLennan R. Blood group, serum protein and red cell enzyme groups of Amerindian populations in Colombia. Am J Phys Anthropol. 1974;41:301-16.*

#### **Costa Rica**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

#### **Cuba**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

#### **Democratic Republic of the Congo**

**Vandeville D**, Martin JP, Ropartz C. Alpha 1-antitrypsin polymorphism of a Bantu population: description of a new allele PiL. *Humangenetik.* 1974;21:33-8.

#### **Denmark**

**Thymann M.** Distribution of alpha-1-antitrypsin (Pi) phenotypes in Denmark determined by separator isoelectric focusing in agarose gel. *Hum Hered* 1986;36:19–23.

**Dahl M**, Nordestgaard BG, Lange P, Vestbo J, Tybjaerg-Hansen A. Molecular diagnosis of intermediate and severe alpha(1)-antitrypsin deficiency: MZ individuals with chronic obstructive pulmonary disease may have lower lung function than MM individuals. *Clin Chem* 2001;47:56–62.

#### **Dominican Republic**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

#### **Ecuador**

**Goedde HW**, Benkmann HG, Flatz G, Bienzle U, Kroeger A. Alpha 1-antitrypsin subtypes in the populations of Germany, Ecuador, Afghanistan, Cameroon, and Saudi-Arabia. *Zeitschrift fur Morphologie und Anthropologie* 1980;70:341-6.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

## **Egypt**

*Settin A1, El-Bendary M, Abo-Al-Kassem R, El Baz R. Molecular analysis of A1AT (S and Z) and HFE (C282Y and H63D) gene mutations in Egyptian cases with HCV liver cirrhosis. J Gastrointest Liver Dis. 2006;15:131-5.*

*Gharib AF, Karam RA, Pasha HF, Radwan MI, Elsayy WH. Polymorphisms of hemochromatosis, and alpha-1 antitrypsin genes in Egyptian HCV patients with and without hepatocellular carcinoma. Gene. 2011;489:98-102.*

## **El Salvador**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

## **Estonia**

**Uibo R**, Parik J, Maaros H-I, Samloff IM. Alpha-1-antitrypsin allo- and phenotypes in gastric and duodenal ulcer. *Scand J Gastroenterol* 1991;26:105–8.

**Beckman L**, Sikström C, Mikelsaar AV, Krumina A, Kucinskas V. a-1-Antitrypsin (PI) alleles of West European influence in the Baltic Sea region. *Hum Hered* 1999;49:52–5.

## **Finland**

**Arnaud P**, Koistinen J, Wilson GB, Fudenberg HH. Alpha-1-antitrypsin (Pi) phenotypes in a Finnish population. *Scand J Clin Lab Invest* 1977;37:339–43.

**Hägglom J**, Kettunen K, Karjalainen J, Heliövaara M, Jousilahti P, Saarelainen S. Prevalence of PI\*Z and PI\*S alleles of alpha-1-antitrypsin deficiency in Finland. *Eur Clin Respir J.* 2015;2:28829.

*Fagerhol MK, Eriksson AW, Monn E. Serum Pi types in some Lappish and Finnish populations. Hum Hered 1969;19:360–4.*

*Fagerhol MK, Gedde-Dahl T Jr. Genetics of the Pi serum types. Family studies of the inherited variants of serum alpha-1-antitrypsin. Hum Hered 1969;19:354–9.*

*Arvilommi H. Studies on serum Pi- and C3-polymorphism in Finland. Acta Pathol Microbiol Scand 1972; 234:1–50.*

## **France**

**Arnaud P**, Chapuis-Cellier C, Vittoz P, Creyssel R. Alpha-1-antitrypsin phenotypes in Lyon, France. *Hum Genet* 1977;39:63–8.

**Sesboue R**, Charlionet R, Vercaigne D, Guimbretiere J, Martin JB. Genetic variants of serum alpha-1-antitrypsin (Pi types) in Bretons. *Hum Hered* 1978;28:280–4.

**Morcamp C**, Charlionet R, Sesboue R, Lefevre F, Martin JP. Genetic variants of the Pi system in Normans: comparison with some European populations. *Bull Eur Physiopathol Respir* 1980; 16: Suppl., 285–91.

**Sesboue R**, Martin JP. Alpha-1-antitrypsin (PI) polymorphism in France, with special regard to the PI\*Z allele. *Hum Hered* 1991;41:340–6.

**Youinou P**, Gueguen A, Martin JP, Jouquan J, Salmon D, Ruffie J, Le Menn G, Salmon C. Genetic variants of serum alpha-1-antitrypsin (Pi types) in a Breton population, the Bigoudens. *Hum Hered* 1984;34:114–9.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. *Clinical Genetics* 2003;63:490-509.

*Robinet-Levy M, Rieunier M. [Method for the identification of Pi groups: 1st statistics from Languedoc]. Rev Franc Transfus 1972;15:61–72.*

## **Gambia**

**Welch SG**, McGregor IA, Williams K. Alpha 1-antitrypsin (pi) phenotypes in a village population from The Gambia, West Africa: Evidence of a new variant occurring at a polymorphic frequency. *Hum Genet* 1980;53:233–5.

## **Germany**

**Genz T**, Martin JP, Cleve H. Classification of alpha1-antitrypsin (Pi) phenotypes by isoelectrofocusing. Distinction of six subtypes of the PiM phenotype. *Hum Genet* 1977; 38:325–32.

**Bencze K**, Sabatke L, Fruhmann G. Alpha 1-antitrypsin: the PiMM subtypes. Do they play a role in development of chronic obstructive pulmonary diseases? *Chest*. 1980;77:761-3.

**Weidinger S**, Schwarzfischer F, Cleve H. Pi subtyping by isoelectric focusing: further genetic studies and application to paternity examinations. *Z Rechtsmed* 1980;86:1–7.

**Weidinger S**, Cleve H, Patutschnick W. Alpha1-antitrypsin: evidence for a fourth PiM allele. Distribution of the PiM subtypes in Southern Germany. *Z Rechtsmed* 1982;88: 203–11.



**Patzelt D**, Schroder H. Genetic-polymorphism of alpha-1-antitrypsin (Pi system) – data on the formal and populationgenetics of the Pi subtypes. *Dtsch Gesundheitsw* 1983; 38:119–23.

*Schmechta H, Geserick G. [Population genetic studies on alpha-1 antitrypsin polymorphism (Pi-system) in the population of Berlin]. Dtsch Gesundheitsw 1972; 27: 1089–91.*

### **Greece**

*Fertakis A, Tsourapas A, Douratsos D, Angelopoulos B. Pi phenotypes in Greeks. Hum Hered. 1974;24:313-6.*

*Scheil HG1, Schuhen S, Huckenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (PI) in the Balkans. Coll Antropol. 2002;26:403-10.*

### **Guatemala**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

### **Guayana**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

### **Haiti**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

### **Honduras**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

### **Hungary**

**Goedde HW**, Benkmann HG, Kriese L, Bogdanski P, Czeizel A, Bères J. Genetic markers among three populations of Hungary. *Gene Geogr* 1987; 1: 109–20.

### **Iceland**

**Kellermann G**, Walter H. Investigations on the population genetics of the alpha-1-antitrypsin polymorphism. *Humangenetik*. 1970;10:145-50.

### **India**

**Kamboh MI**, Board PG, Kirwood C. Re-evaluation of the proposed interrelationship between thyroxine-binding globulin (TBG) and alpha1-antitrypsin (PI). *Clin Chim Acta* 1984;139:65–73.

**Walter H**, Naidu JM, Danke-Hopfe H, et al. Genetic serum protein markers in eight south Indian caste and tribal populations. *Z Morphol Anthropol* 1993; 79: 355–65.

**Reddy PH**, Mastana SS. Genetic variation of serum proteins (GC, TF and PI subtypes) in the Baigas of Madhya Pradesh, India. *Anthropol Anz* 1995;53:53–6.

### **Indonesia**

**Constans J**, Gouaillard C, Brequet G. Serum protein polymorphism in Bali (Indonesia). *Ann Hum Biol* 1986;13: 537–45.

*Kirk RL, McDermid EM, Blake NM, Wight RL, Yap EH, Simons MJ. The distribution of red cell enzyme and serum protein groups in a population of Dani (Pit River, West Irian). Humangenetik 1973; 17: 345–50.*

*Malcolm LA, Woodfield DG, Blake NM, Kirk RL, McDermid EM. The distribution of blood, serum protein and enzyme groups on Manus Island (Admiralty Islands, New Guinea). Hum Hered 1972; 22: 305–22.*

### **Iran**

**Hashemi M**, Alavian SM, Ghavami S, de Serres FJ, Salehi M, Doroudi T, Fard AH, Mehrabifar H, Milani B, Shahri SJ. High prevalence of alpha 1 antitrypsin deficient phenotypes in viral hepatitis B infected patients in Iran. *Hepato Res* 2005;33:292–7.

**Walter H**, Mohammadzadeh Z, Schuler I, Farhud DD. Serum protein polymorphisms (HP, TF, GC and PI) in four Iranian population samples. *Ann Hum Biol* 1992;19:35–9.

### **Israel**

*Nevo S, Kanaaneh H, Cleve H. A genetic study of alpha 1-antitrypsin in an Israeli Arab population, with a new allele: Piv-s. Isr J Med Sci 1982;18:891–3.*

*Nevo S. Protease inhibitor subtypes in some population groups from Israel. Hum Hered 1987;37:170–81.*

Nevo S, Cleve H, Bar-Shani, Joel A., Liron M. Bulgarian Jews in Israel: genetic blood markers. Red-cell antigens, serum proteins and red-cell isozymes. *Hum Hered* 1989; 39:333–44.

Nevo S, Cleve H. PI polymorphism in Israel: report on six Jewish population groups. *Am J Med Genet* 1991;39:399–403.

Nevo S, Cleve H, Koller A, Eigel E, Patutschnick W, Kanaaneh H, Joel A. Serum protein polymorphisms in Arab Moslems and Druze of Israel: BF, F13B, AHSB, GC, PLG, PI, and TF. *Hum Biol* 1992;64:587–603.

Ablin JN, Bar-Shira A, Yaron M, Orr-Urtreger A. Candidate-gene approach in fibromyalgia syndrome: association analysis of the genes encoding substance P receptor, dopamine transporter and alpha1-antitrypsin. *Clin Exp Rheumatol*. 2009;27(5 Suppl 56):S33-8

## Italy

**Piantelli M**, Auconi P, Musiani P. Alpha-1-antitrypsin phenotypes in new-borns from Central and Southern Italy. *Hum Hered* 1978;28:468–73.

**Massi G**, Fabiano A, Ragusa D, Auconi P, Biagini R, Farinato MA, Bandelloni R. Alpha-1-antitrypsin phenotypes and Pi M subtypes in Italy. *Bull Eur Physiopathol Respir* 1980; 16, Suppl., 301–6.

**Pascali VL**, De Mercurio D. Determination of alpha-1-antitrypsin subtypes in the population of Rome: a study in ultrathin-layer isoelectric focusing. *Hum Hered* 1981;31:296–8.

**Pittschieler K**, Massi G. Alpha-1-antitrypsin deficiency in two population groups in north Italy. *Pediatr Padol* 1988;23:307–11.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. *Clinical Genetics* 2003;63:490-509.

**de Serres FJ**, Luisetti M, Ferrarotti I, Blanco I, Fernández-Bustillo E. Alpha-1 antitrypsin deficiency in Italy: regional differences of the PIS and PIZ deficiency alleles of alpha-1 antitrypsin deficiency in Italy. *Monaldi Arch Chest Dis* 2005; 63:133-41.

Seefried-Lehmann S-J, Walter H, Danker-Hopfe H, Ricjards O, De Stefano GF. Investigations on the variability of four genetic serum protein markers (HP; TF; CG and PI subtypes) in Italy. *Z Morphol Anthropol* 1992;79:215–31.

*Massi G, Cotumaccio R, Auconi P. Alpha-1-antitrypsin (alpha-1-AT) phenotypes and PiM subtypes in Italy. Evidence of considerable geographic variability. Hum Genet 1982;61:76–7.*

*Stakisaitis D, Benetis R. Does alpha-1-proteinase inhibitor play a protective role in coronary atherosclerosis? Med Sci Monit 2001;7:701–11.*

### **Jamaica**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

### **Japan**

**Harada S**, Miyake K, Suzuki H, Oda T. New phenotypes of serum alpha1-antitrypsin in Japanese detected by gel slab isoelectric focusing. *Hum Genet* 1977;38:333–6.

**Miyake K**, Suzuki H, Oka H, Oda T, Harada S. Distribution of alpha 1-antitrypsin phenotypes in Japanese: description of Pi M subtypes by isoelectric focusing. *Jinrui Idengaku Zasshi* 1979;24:55–62.

**Ohtani H**, Saito M. Alpha-1-antitrypsin: frequencies of PiM subtypes and serum concentration in the Japanese population. *Hum Hered* 1985;35:62–4.

**Sebetan IM**, Akaishi S. Alpha-1-antitrypsin (PiM) subtypes in Japanese. *Forensic Sci Int* 1981;18:155–9.

**Yuasa I**, Suenaga K, Gotoh Y, Ito K, Yokoyama N, Okada K. PI(alpha 1-antitrypsin) polymorphism in the Japanese: confirmation of PI\*M4 and description of new PI variants. *Hum Genet* 1984;67:209–12.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI\*S and PI\*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. *Eur Respir J.* 2006;28:1091-9.

### **Jordan**

*Saleh H, Davrinche C, Charlionet R, Rivat C. Alpha-1-antitrypsin phenotypes in a population of Jordan. Hum Hered 1986;36:192–4.*

### **Kazakhstan**

**Petrishev VN**, Lebedeva IA, Shneider Iu V. [Genetic polymorphism of the alpha 1-antitrypsin system in the native population of the Kazakh SSR]. *Genetika* 1987; 23:2257–64.

## Latvia

**Beckman L**, Sikstoms C, Mikelsaar AV, Krumina A, Kucinskas V.  $\alpha$ -1-Antitrypsin (PI) alleles of West European influence in the Baltic Sea region. *Hum Hered* 1999;49:52–55.

## Lithuania

**Beckman L**, Sikstoms C, Mikelsaar AV, Krumina A, Kucinskas V.  $\alpha$ -1-Antitrypsin (PI) alleles of West European influence in the Baltic Sea region. *Hum Hered* 1999;49: 52–5.

**Stakisaitis D**, Benetis R. Does alpha-1-proteinase inhibitor play a protective role in coronary atherosclerosis? *Med Sci Monit* 2001;7:701–11.

## Macedonia

**Scheil HG**, Schuhen S, Huckenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (PI) in the Balkans. *Coll Antropol*. 2002;26:403-10.

**Fertakis A**, Tsourapas A, Douratsos D, Angelopoulos B. *Pi phenotypes in Greeks. Hum Hered*. 1974;24:313-6.

## Malaysia

**Saha N**. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. *Ann Hum Biol* 1990;17:229–34.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI\*S and PI\*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. *Eur Respir J*. 2006;28:1091-9.

## Mali

Frants RR, Eriksson AW. Reliable classification of six Pi M subtypes by separator isoelectric focusing. *Hum Hered* 1978;28:201–9.

## Mexico

**Pérez-Rubio G**, Jiménez-Valverde LO, Ramírez-Venegas A, Camarena Á, Sansores RH, Flores-Trujillo F, Reséndiz-Hernández JM, Falfán-Valencia R. Prevalence of alpha-1 antitrypsin high-risk variants in Mexican mestizo population and their association with lung function values. *Arch Bronconeumol*. 2015;51:80-5.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

## Morocco

**Ezzikouri S**, El Feydi AE, El Kihal L, Afifi R, Benazzouz M, Hassar M, Chafik A, Pineau P, Benjelloun S. Prevalence of common HFE and SERPINA1 mutations in patients with hepatocellular carcinoma in a Moroccan population. *Arch Med Res.* 2008;39:236-41.

*El Fakir S, Nejjari C, Serhier Z, Tessier JF, Barberger-Gateau P, Dartigues JF, Raheison C. Alpha 1 antitrypsin phenotypes and obstructive airway disease in subjects over 65 years of age: QUID R cohort. Age Ageing. 2011;40:627-9*

### **Mozambique**

**Kellermann G**, Walter H. Investigations on the population genetics of the alpha-1-antitrypsin polymorphism. *Humangenetik* 1970;10:145–150.

### **Namibia**

*Dunn DS, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. Hum Hered 1986;36:238 –242.*

### **Nepal**

**Yuasa I**, Saneshige Y, Okamoto N, Ikawa S, Hikita T, Ikebuchi J, Inoue T, Okada K. Distribution of Hp, Tf, Gc and Pi polymorphisms in a Nepalese population. *Hum Hered.* 1983;33:302-6.

### **Netherlands**

**Frants RR**, Eriksson AW. Reliable classification of six Pi M subtypes by separator isoelectric focusing. *Hum Hered.* 1978;28:201-9.

**Hoffman JJML**, van den Broek WGM. Distribution of alpha-1-antitrypsin phenotypes in two Dutch population groups. *Hum Genet* 1976;32:43–48.

**Klasen EC**, Biemond I, Weterman IT. Alpha 1-antitrypsin levels and phenotypes in Crohn's disease in the Netherlands. *Gut* 1980;21:840–842.

**Biedmond I**, Shelby WS, Jewell DP. Alpha-1-antitrypsin serum concentration and phenotypes in ulcerative colitis. *Digestion* 1984; 29:124–128.

### **New Zealand**

**Janus ED**, Joyce PR, Sheat JM, Carrell RW. Alpha-1-antitrypsin variants in New Zealand. *N Z Med J* 1975;82:289–91.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. *Clin Genet* 2003;64:382-97.

### **Nicaragua**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

### **Nigeria**

**Olusi SO**, Adeyemo AO. Alpha-1-antitrypsin phenotypes, serum immunoglobulins and circulating bacterial antibodies in Nigerians with chronic obstructive pulmonary disease. *East Afr Med J* 1983;60:506-11.

*DeCroo S, Kamboh MI, Ferrell RE. Population genetics of alpha-1-antitrypsin polymorphism in US whites, US blacks and African blacks. Hum Hered 1991;41:215–21.*

*Awotedu AA, Adelaja AB. Alpha-1-antitrypsin and chronic bronchitis in adult Nigerians. Afr J Med Med Sci 1990;19:291–7.*

### **Norway**

**Fagerhol MK**, Tenfjord OW. Serum Pi types in some European, American, Asian and African populations. *Acta Pathol Microbiol Scand*. 1968;72:601-8.

**Fagerhol MK**, Gedde-Dahl T Jr. Serum Pi types in Norwegians. *Acta Pathol Microbiol Scand* 1967;70: 421–8.

**Moller P**, Fagerhol MK. Pi phenotypes in patients with ankylosing spondylitis, acute anterior uveitis, psoriasis and chronic prostatitis. *Clin Genet* 1984; 26: 339–41.

**Gulsvik A**, Fagerhol MK. Alpha-1-antitrypsin phenotypes and obstructive lung disease in the city of Oslo. *Scand J Respir Dis* 1979; 60: 267–74.

### **Pakistan**

**Shahid A**, Siddiqui AA, Zuberi SJ, Waqar M. Phenotypes of alpha 1 antitrypsin in Karachi, Pakistan. *J Pak Med Assoc* 2000; 50: 374–6.

### **Panama**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

### **Papua New Guinea**

**Malcolm LA**, Woodfield DG, Blake NM, Kirk RL, McDermid EM. The distribution of blood, serum protein and enzyme groups on Manus Island (Admiralty Islands, New Guinea). *Hum Hered*. 1972;22:305-22.

**Kirk RL**, McDermid EM, Blake NM, Wight RL, Yap EH, Simons MJ. The distribution of red cell enzyme and serum protein groups in a population of Dani (Pit River, West Irian). *Humangenetik*. 1973;17:345-50.

### **Paraguay**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

### **Peru**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

### **Philippines**

**Saha N**. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. *Ann Hum Biol* 1990;17:229–34.

### **Poland**

**Titenko-Holland NV**, Kowalska A. Alpha-1-antitrypsin (PI) subtypes in Russians and Poles. *HumHered* 1992; 42:384–6.

**Walter H**, Danker-Hopfe H, Lemmermann M, Lorenz M. Investigation on the variability of four genetic protein markers in Poland. *Z Morphol Anthropol* 1992;79: 203–14.

**Kolwaska A**, Rujner J. Polymorphism locus Pi (alpha-1-antitrypsin) of residents in the Poznan Province. *Polski Tygodnik Lekarski (Warszawa)* 1994;43:195–7.

**Kowalska A**, Rujner J, Titenko-Holland NV, Pitacik B. Alpha-1-antitrypsin subtypes in Polish newborns. *Hum Hered* 1995;45:351–4.

**Chorostowska-Wynimko J**, Struniawski R, Popławska B, Borszewska-Kornacka M. The incidence of alpha-1-antitrypsin (A1AT) deficiency alleles in population of Central Poland--preliminary results from newborn screening. *Pneumonol Alergol Pol*. 2012;80:450-3.

*Szczeklik AB, Turowska B, Gzerniawska-Mysik B, Opolska B, Nizankowska E. Serum alpha-1 antitrypsin in bronchial asthma. Am Rev Respir Dis 1974;109:487–90.*

### **Portugal**

**Geda HM**, Pereira-Albino JA, Manso C. Polimorphism of alpha-1-antitrypsin in a Portuguese population. *Hum Genet* 1976;32:109–13.



**Martin JP**, Seboue R, Charlionet R, Ropartz C, Pereira MT. Genetic variants of serum alpha-1-antitrypsin (Pi types) in Portuguese. *Hum Hered* 1976;26:310–4.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. *Clinical Genetics* 2003;63:490-509.

**Spínola C**, Bruges-Armas J, Pereira C, Brehm A, Spínola H. Alpha-1-antitrypsin deficiency in Madeira (Portugal): the highest prevalence in the world. *Respir Med*. 2009;103:1498-502.

*Santos-Rosa MA, Robalo-Cordeiro AJA. Alpha-1-proteinase inhibitor phenotypes in Portuguese population. Eur J Respir Dis 1986; 69: Suppl. 146,167–73.*

### **Puerto Rico**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

### **Republic of the Congo**

**Pascali VL**, Ranalletta D, Spedini G. Antitrypsin and Gc polymorphisms in some populations of Congo: an unusual, highly frequent mutant, PIS, in Bateke and Babenga. *Ann Hum Biol* 1986;13:267–71.

### **Republic of Ireland**

**Carroll TP**, O'Connor CA, Floyd O, McPartlin J, Kelleher DP, O'Brien G, Dimitrov BD, Morris VB, Taggart CC, McElvaney NG. The prevalence of alpha-1 antitrypsin deficiency in Ireland. *Respir Res*. 2011;12:91.

### **Romania**

*Scheil HG, Schuhen S, Huckenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (PI) in the Balkans. Coll Antropol. 2002;26:403-10.*

### **Russia**

**Shurkhal AV**, Podogas AV, Altukhov YP. Genetic polymorphism and rare variants of alpha-1-antitrypsin in the population of Moscow. *Genetika* 1984;20: 2066–9.

**Shurkhal AV**, Podogas AV, Dementeva GM, et al. Rare variants of alpha 1 antitrypsin in a group of families with developmental disorders in newborns. *Genetika* 1987;23:1692–8.

**Anatonov MP**, Petrukhin IS, Voitova TV, Portenko GM. Distribution of Pi phenotypes in normal subjects and subjects with diseases of the respiratory tract. *Lab Delo* 1988;3:23–7.

**Kucher AN**, Puzyrev VP, Ivanova OF, Xu JJ, Oz RF. Subtypes of serum proteins among Russian inhabitants of the Tomsk region. *Genetika* 1993;29:845–52.

**Kravchuk OI**, Spitsyn VA, Ginter EK, Makarov SV. Population genetic characteristics of Highland and Meadow Mari: genetic markers. *Hum Genet* 1996;32:570–5.

### **Saudi Arabia**

*Goedde HW, Benkmann HG, Agarwal DP, Hirth L, Bienzle U, Dietrich M, Hoppe HH, Orlowski J, Kohne E, Kleihauer E. Genetic studies in Saudi Arabia: red cell enzyme, haemoglobin and serum protein polymorphisms. Am J Phys Anthropol 1979;50:271–7.*

*Al-Balla SRS, El-Hazmi MAF, Al-Dalaan AN, Warsy AS. Alpha-1-antitrypsin phenotypes in rheumatoid-arthritis. Saudi Med J 1992;13:555–9.*

*Warsy AS, Elhazmi MAF, Sedrani SH. Alpha-1-antitrypsin phenotypes in Saudi-Arabia - a study in the Central Province. Ann Saudi Med 1991;11:159–62.*

*Warsy AS, el-Hazim MAF, Hamooda H, Kilic N. Alpha-1-antitrypsin: frequencies of Pim subtypes in a Saudi Population. Saudi Med J 1991;12:376–9.*

### **Serbia**

**Jelic-Ivanovic Z**, Spasojevic-Kalimanovska V, Topic A, Spasic S, Petrovic V. Alpha-1-antitrypsin (Pi) polymorphism in Serbia: deviation of Pi M subtype distribution from the Hardy-Weinberg equilibrium. *Gene Geogr* 1994;8:129–35.

### **Singapore**

**Saha N**. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. *Ann Hum Biol* 1990;17:229–34.

### **Somalia**

**Massi G**, Vecchio FM. Alpha-1-antitrypsin phenotypes in a group of newborn infants in Somalia. *Hum Genet* 1977;38:265–9.

### **South Africa**

**Dunn DS**, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. *Hum Hered* 1986;36:238–42.

**Hitzeroth HW**, Bissbort S, van den Berg CT, van Niekerk MM, Op't Hof J. Alpha-1-antitrypsin genetic polymorphism in South Africa. *S Afr Med J* 1987;71:755–9.

**Walter H**, Eberhardt D, Tegeler M, Wiechmann I, Dankerhopfe H, Bhattachaochya SK, Das MK, Mukherjee BN, Saha PN, Das K. HP, GC and PI polymorphisms in sixteen central Indian tribal populations. *Gene Geogr* 1993;7:193–202.

### **South Korea**

**Lee SS**, Lawton JW, Ko KH. Alpha1 antitrypsin phenotypic variability is not associated with ANCA in southern Chinese. *Ann Rheum Dis* 2001;60:725–6.

*Saha N. Alpha 1-protease inhibitor (PI) subtypes in seven populations of East Asia. Ann Hum Biol* 1990;17:229–34.

### **Spain**

**Vidal R**, Miravittles M, Jardí R, Torrella M, Rodríguez-Frías F, Moral P, Vaqué J. [Study of the frequency of different phenotypes of alpha-1-antitrypsin in a population of Barcelona]. *Med Clin (Barcelona)* 1996;107:211–14.

**Blanco I**, Fernández E, Rodríguez MC, Fernández A. [Allelic frequency of the gene of alpha-1-antitrypsin in the general population in a county in Asturias]. *Med Clin (Barcelona)* 1999;133:366–70.

**Puertas MJA**, Arranz-Peña ML. Valoración de fenotipos de alfa-1-antitripsina en Castilla-León: estudio en una población de recién nacidos [Evaluation of phenotypes of alpha-1-antitrypsin in Castilla-Leon: study in a population of newborns]. *Rev Clin Esp* 1989;184:106–8.

**Moral P**. Estudio antropogenético de diversos polimorfismos hematológicos en la isla de Menorca [Anthropogenetic study of various haematological polymorphisms on Menorca island]. PhD thesis. University of Barcelona, Barcelona, Spain, 1986.

**Blanco I**, Fernández-Bustillo E, de Serres FJ, Alkassam D, Rodríguez Menéndez C. PI\*S and PI\*Z alpha 1-antitrypsin deficiency: estimated prevalence and number of deficient subjects in Spain. *Med Clin (Barc)* 2004; 123:761-5.

*Blanco I, Fernandez E Alpha1-antitrypsin Pi phenotypes S and Z in Spain: an analysis of the published surveys. Respir Med* 2001;95:109-14.

*de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. Clinical Genetics* 2003; 63:490-509.

### **Suriname**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

#### **Sweden**

**Hjalmarsson K.** Distribution of alpha-1-antitrypsin phenotypes in Sweden. *Hum Hered* 1988;38:27–30.

*Sveger T. Liver disease in alpha1-antitrypsin deficiency detected by screening of 200,000 infants. N Engl J Med.* 1976;294:1316-21.

#### **Switzerland**

**Bar W**, Kratzer A. Polymorphism of alpha-1-antitrypsin (Pi) in the Swiss population determined by isoelectric focusing with an immobilized pH gradient. *Hum Hered* 1988; 38:106–110.

*Scheffrahn W, Ziggotti E. [Electrophoretic alpha 1-antitrypsin variation in the Swiss population]. [Article in German] Anthropol Anz.* 1982;40:137-43.

#### **Tajikistan**

*Spitsyn VA, Novoradovskii AG, Spitsyna NK, Parik Iula. [Polymorphism of alpha1-antitrypsin in Pamir populations. Reproductive compensation--possible mechanism for maintaining genetic diversity gor PI genes in human populations]. Genetika* 1989; 25: 2218–2224.

#### **Thailand**

**Pongaew P**, Schelp FP. Alpha-1-protease inhibitor phenotypes and serum concentrations in Thailand. *Hum Genet* 1980;54: 119–24.

*Chongsrisawat V, Jantaradsamee P, Vivatvakin B, Pongpaew P, Poovorawan Y. Alpha 1-antitrypsin phenotype of children with liver diseases in Thailand. Asian Pac J Allergy Immunol* 1998;16: 27–30.

*Saha N. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. Ann Hum Biol* 1990;17: 229–34.

#### **Trinidad and Tobago**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

#### **Tunisia**

Denden S, Zorzetto M, Amri F, Knani J, Ottaviani S, Scabini R, Gorrini M, Ferrarotti I, Campo I, Chibani JB, Khelil AH, Luisetti M. Screening for Alpha 1 antitrypsin deficiency in Tunisian subjects with obstructive lung disease: a feasibility report. *Orphanet J Rare Dis.* 2009;4:12.

*Chaabani H, Martin JP, Frants RR, Lefranc G. Genetic study of Tunisian Berbers. II. Alpha 1-antitrypsin (Pi) polymorphism: report of a new allele (Pi S Berber). Exp Clin Immunogenet. 1984;1:19-24.*

*Chaâbani H, Cox DW. Genetic characterization and origin of Tunisian Berbers. Hum Hered. 1988;38:308-16.*

*Denden S, Haj Khelil A, Perrin P, Daimi H, Leban N, Ouaja A, Mahdouani K, Hlioui L, Lefranc G, Ben Chibani J. Alpha 1 antitrypsin polymorphism in the Tunisian population with special reference to pulmonary disease. Pathol Biol (Paris). 2008;56:106-10.*

### **United Kingdom**

**Cook PJ.** The genetics of alpha1-antitrypsin: a family study in England and Scotland. *Ann Hum Genet.* 1975;38:275-87.

**Blundell G,** Frazer A. Alpha1 antitrypsin phenotypes in Northern Ireland. *Ann Hum Genet* 1975; 38:289–94.

**Arnaud P,** Galbraith RM, Faulk WP, Black C. Pi phenotypes of alpha1-antitrypsin in Southern England: identification of M subtypes and implications for genetic studies. *Clin Genet* 1979; 15:406–10.

**Phillips C,** Whitehouse D. A rapid high resolution separator IEF technique for phenotyping alpha-1 antrypsin. *Adv Foren Haemogen* 1990;8: 274–6.

*Papiha SS, Pal B, Walker D, Mangion P, Hossain MA. Alpha 1 antitrypsin (PI) phenotypes in two rheumatic diseases: a reappraisal of the association of PI subtypes in rheumatoid arthritis. Ann Rheum Dis. 1989;48:48-52.*

*Sanders PA, Thomson W, Browne DA, Dyer PA, Grennan DM. alpha 1 Antitrypsin (PI) allotypes in rheumatoid arthritis. Ann Rheum Dis. 1986;45:450-3.*

*Biernond I, Selby WS, Jewell DP, Klasen EC. Alpha 1-antitrypsin serum concentration and phenotypes in ulcerative colitis. Digestion. 1984;29:124-8.*

### **United States**

**Kueppers F.** Serum alpha-1-antitrypsin levels. *N Engl J Med* 1968;279:164–5.

**Kueppers F.** Alpha1-antitrypsin deficiency. *Birth Defects Orig Artic Ser* 1976;12:213–22.

**Kueppers F**, Utz G, Simon B. Alpha1-antitrypsin deficiency with M-like phenotype. *J Med Genet* 1977;14:183–6.

**Kueppers F**, Offord KP. Alpha-1 antitrypsin elevation in healthy neonates. *J Lab Clin Med* 1979;94:475–480.

**Roberts A**, Kagan A, Rhoads GG et al. Antitrypsin and chronic obstructive pulmonary disease among Japanese-American men. *Chest* 1977;72:489–91.

**Lee CC**, Kueppers F, Harpel B et al. Alpha-1-antitrypsin (Pi) types in Korean and Chinese populations. *Hum Genet* 1981;57:327–328.

**Evans HE**, Bognacki NS, Perrott LM et al. Prevalence of alpha 1-antitrypsin Pi types among newborn infants of different ethnic backgrounds. *J Pediatr* 1977;90:621–4.

**DeCroo S**, Kamboh MI, Ferrell RE. Population genetics of alpha-1-antitrypsin polymorphism in US whites, US blacks and African blacks. *Hum Hered* 1991;41:215–21.

**Pierce JA**, Eradio B, Dew TA. Antitrypsin phenotypes in St. Louis. *JAMA* 1975; 231:609–12.

**Lieberman J**, Gaidulis L, Roberts L. Racial distribution of alpha1-antitrypsin variants among junior high school students. *Am Rev Respir Dis* 1976;114:1194–8.

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. *Clin Genet* 2003;64:382-97.

*O'Brien ML, Buist NR, Murphey WH. Neonatal screening for alpha1-antitrypsin deficiency. J Pediatr. 1978;92:1006-10.*

*Silverman EK, Miletich JP, Pierce JA, Sherman LA, Endicott SK, Broze GJ Jr, Campbell EJ. Alpha-1-antitrypsin deficiency. High prevalence in the St. Louis area determined by direct population screening. Am Rev Respir Dis. 1989;140:961-6.*

## **Uruguay**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

## **Venezuela**

**de Serres FJ**, Blanco I, Fernández-Bustillo E. Estimates of PI\*S and PI\*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

*Fragoso AT, Barrios GB, Castañeda GC. [The value of the Pi system phenotype in alpha 1-antitrypsin deficiency]. [Spanish]. Acta Gastroenterologica Latinoamericana 1986;16:75-80.*

*Marini E, Moral P, Petralanda I. Serum protein markers in the Piaroa Indians of Amazonia (Venezuela). Hum Hered 1993;43:232-8.*

*Fonseca-Perez T, Gonzalez-Coira M, Arias S. PI locus (alpha-1-antitrypsin) allelic frequencies in an Andean Venezuelan population. Gene Geography 1996;10: 65-74*







