

Supplementary material

References for the trials included in the review

1. Ahren B, Gomis R, Standl E, Mills D, Schweizer A. Twelve- and 52-week efficacy of the dipeptidyl peptidase IV inhibitor LAF237 in metformin-treated patients with type 2 diabetes. *Diabetes care*. 2004;27(12):2874-80.
2. Al-Shaikh AR. Comparison if basal insulin added to oral agents versus twice-daily premixed insulin as initial insulin therapy for type 2 diabetes. *Pak J Med Sci*. 2005;22(1):14-7.
3. Aljabri K, Kozak SE, Thompson DM. Addition of pioglitazone or bedtime insulin to maximal doses of sulfonylurea and metformin in type 2 diabetes patients with poor glucose control: a prospective, randomized trial. *The American journal of medicine*. 2004;116(4):230-5.
4. Arechavaleta R, Seck T, Chen Y, Krobot KJ, O'Neill EA, Duran L, et al. Efficacy and safety of treatment with sitagliptin or glimepiride in patients with type 2 diabetes inadequately controlled on metformin monotherapy: a randomized, double-blind, non-inferiority trial. *Diabetes, obesity & metabolism*. 2011;13(2):160-8.
5. Aschner P, Chan J, Owens DR, Picard S, Wang E, Dain MP, et al. Insulin glargine versus sitagliptin in insulin-naive patients with type 2 diabetes mellitus uncontrolled on metformin (EASIE): a multicentre, randomised open-label trial. *Lancet*. 2012;379(9833):2262-9.
6. Barnett AH, Burger J, Johns D, Brodows R, Kendall DM, Roberts A, et al. Tolerability and efficacy of exenatide and titrated insulin glargine in adult patients with type 2 diabetes previously uncontrolled with metformin or a sulfonylurea: a multinational, randomized, open-label, two-period, crossover noninferiority trial. *Clinical therapeutics*. 2007;29(11):2333-48.
7. Bergenstal R, Lewin A, Bailey T, Chang D, Gylvin T, Roberts V. Efficacy and safety of biphasic insulin aspart 70/30 versus exenatide in subjects with type 2 diabetes failing to achieve glycemic control with metformin and a sulfonylurea. *Current medical research and opinion*. 2009;25(1):65-75.
8. Bergenstal RM, Wysham C, Macconell L, Malloy J, Walsh B, Yan P, et al. Efficacy and safety of exenatide once weekly versus sitagliptin or pioglitazone as an adjunct to metformin for treatment of type 2 diabetes (DURATION-2): a randomised trial. *Lancet*. 2010;376(9739):431-9.
9. Berhanu P, Perez A, Yu S. Effect of pioglitazone in combination with insulin therapy on glycaemic control, insulin dose requirement and lipid profile in patients with type 2 diabetes previously poorly controlled with combination therapy. *Diabetes, obesity & metabolism*. 2007;9(4):512-20.
10. Blonde L, Dagogo-Jack S, Banerji MA, Pratley RE, Marcellari A, Braceras R, et al. Comparison of vildagliptin and thiazolidinedione as add-on therapy in patients inadequately controlled with metformin: results of the GALIANT trial--a primary care, type 2 diabetes study. *Diabetes, obesity & metabolism*. 2009;11(10):978-86.
11. Bolli G, Dotta F, Colin L, Minic B, Goodman M. Comparison of vildagliptin and pioglitazone in patients with type 2 diabetes inadequately controlled with metformin. *Diabetes, obesity & metabolism*. 2009;11(6):589-95.

12. Bosi E, Camisasca RP, Collober C, Rochotte E, Garber AJ. Effects of vildagliptin on glucose control over 24 weeks in patients with type 2 diabetes inadequately controlled with metformin. *Diabetes care*. 2007;30(4):890-5.
13. Brazg R, Xu L, Dalla Man C, Cobelli C, Thomas K, Stein PP. Effect of adding sitagliptin, a dipeptidyl peptidase-4 inhibitor, to metformin on 24-h glycaemic control and beta-cell function in patients with type 2 diabetes. *Diabetes, obesity & metabolism*. 2007;9(2):186-93.
14. Bunck MC, Diamant M, Corner A, Eliasson B, Malloy JL, Shaginian RM, et al. One-year treatment with exenatide improves beta-cell function, compared with insulin glargine, in metformin-treated type 2 diabetic patients: a randomized, controlled trial. *Diabetes care*. 2009;32(5):762-8.
15. Charbonnel B, Karasik A, Liu J, Wu M, Meininger G, Sitagliptin Study G. Efficacy and safety of the dipeptidyl peptidase-4 inhibitor sitagliptin added to ongoing metformin therapy in patients with type 2 diabetes inadequately controlled with metformin alone. *Diabetes care*. 2006;29(12):2638-43.
16. Charbonnel B, Scherthaner G, Brunetti P, Matthews DR, Urquhart R, Tan MH, et al. Long-term efficacy and tolerability of add-on pioglitazone therapy to failing monotherapy compared with addition of gliclazide or metformin in patients with type 2 diabetes. *Diabetologia*. 2005;48(6):1093-104.
17. Charpentier G, Fleury F, Kabir M, Vaur L, Halimi S. Improved glycaemic control by addition of glimepiride to metformin monotherapy in type 2 diabetic patients. *Diabetic medicine : a journal of the British Diabetic Association*. 2001;18(10):828-34.
18. Charpentier G, Halimi S. Earlier triple therapy with pioglitazone in patients with type 2 diabetes. *Diabetes, obesity & metabolism*. 2009;11(9):844-54.
19. Cho YM, Koo BK, Son HY, Lee KW, Son HS, Choi DS, et al. Effect of the combination of mitiglinide and metformin on glycemic control in patients with type 2 diabetes mellitus. *Journal of diabetes investigation*. 2010;1(4):143-8.
20. Dailey GE, 3rd, Noor MA, Park JS, Bruce S, Fiedorek FT. Glycemic control with glyburide/metformin tablets in combination with rosiglitazone in patients with type 2 diabetes: a randomized, double-blind trial. *The American journal of medicine*. 2004;116(4):223-9.
21. Davies MJ, Thaware PK, Tringham JR, Howe J, Jarvis J, Johnston V, et al. A randomized controlled trial examining combinations of repaglinide, metformin and NPH insulin. *Diabetic medicine : a journal of the British Diabetic Association*. 2007;24(7):714-9.
22. DeFronzo RA, Hissa MN, Garber AJ, Luiz Gross J, Yuyan Duan R, Ravichandran S, et al. The efficacy and safety of saxagliptin when added to metformin therapy in patients with inadequately controlled type 2 diabetes with metformin alone. *Diabetes care*. 2009;32(9):1649-55.
23. DeFronzo RA, Ratner RE, Han J, Kim DD, Fineman MS, Baron AD. Effects of exenatide (exendin-4) on glycemic control and weight over 30 weeks in metformin-treated patients with type 2 diabetes. *Diabetes care*. 2005;28(5):1092-100.
24. Derosa G, Franzetti IG, Querci F, Carbone A, Ciccarelli L, Piccinni MN, et al. Exenatide plus metformin compared with metformin alone on beta-cell function in patients with Type 2 diabetes. *Diabetic medicine : a journal of the British Diabetic Association*. 2012;29(12):1515-23.

25. Derosa G, Maffioli P, Salvadeo SA, Ferrari I, Ragonesi PD, Querci F, et al. Exenatide versus glibenclamide in patients with diabetes. *Diabetes technology & therapeutics*. 2010;12(3):233-40.
26. Derosa G, Mereu R, D'Angelo A, Salvadeo SA, Ferrari I, Fogari E, et al. Effect of pioglitazone and acarbose on endothelial inflammation biomarkers during oral glucose tolerance test in diabetic patients treated with sulphonylureas and metformin. *Journal of clinical pharmacy and therapeutics*. 2010;35(5):565-79.
27. Derosa G, Salvadeo SA, D'Angelo A, Ferrari I, Mereu R, Palumbo I, et al. Metabolic effect of repaglinide or acarbose when added to a double oral antidiabetic treatment with sulphonylureas and metformin: a double-blind, cross-over, clinical trial. *Current medical research and opinion*. 2009;25(3):607-15.
28. Diamant M, Van Gaal L, Stranks S, Guerci B, MacConell L, Haber H, et al. Safety and efficacy of once-weekly exenatide compared with insulin glargine titrated to target in patients with type 2 diabetes over 84 weeks. *Diabetes care*. 2012;35(4):683-9.
29. Diamant M, Van Gaal L, Stranks S, Northrup J, Cao D, Taylor K, et al. Once weekly exenatide compared with insulin glargine titrated to target in patients with type 2 diabetes (DURATION-3): an open-label randomised trial. *Lancet*. 2010;375(9733):2234-43.
30. Dorkhan M, Dencker M, Stagmo M, Groop L. Effect of pioglitazone versus insulin glargine on cardiac size, function, and measures of fluid retention in patients with type 2 diabetes. *Cardiovascular diabetology*. 2009;8:15.
31. Einhorn D, Rendell M, Rosenzweig J, Egan JW, Mathisen AL, Schneider RL. Pioglitazone hydrochloride in combination with metformin in the treatment of type 2 diabetes mellitus: a randomized, placebo-controlled study. The Pioglitazone 027 Study Group. *Clinical therapeutics*. 2000;22(12):1395-409.
32. Esposito K, Ciotola M, Maiorino MI, Gualdiero R, Schisano B, Ceriello A, et al. Addition of neutral protamine lispro insulin or insulin glargine to oral type 2 diabetes regimens for patients with suboptimal glycemic control: a randomized trial. *Annals of internal medicine*. 2008;149(8):531-9.
33. Fadini GP, de Kreutzenberg SV, Mariano V, Boscaro E, Bertolini F, Mancuso P, et al. Optimized glycaemic control achieved with add-on basal insulin therapy improves indexes of endothelial damage and regeneration in type 2 diabetic patients with macroangiopathy: a randomized crossover trial comparing detemir versus glargine. *Diabetes, obesity & metabolism*. 2011;13(8):718-25.
34. Feinglos M, Dailey G, Cefalu W, Osei K, Tayek J, Canovatchel W, et al. Effect on glycemic control of the addition of 2.5 mg glipizide GITS to metformin in patients with T2DM. *Diabetes research and clinical practice*. 2005;68(2):167-75.
35. Ferrannini E, Fonseca V, Zinman B, Matthews D, Ahren B, Byiers S, et al. Fifty-two-week efficacy and safety of vildagliptin vs. glimepiride in patients with type 2 diabetes mellitus inadequately controlled on metformin monotherapy. *Diabetes, obesity & metabolism*. 2009;11(2):157-66.
36. Filozof C, Gautier JF. A comparison of efficacy and safety of vildagliptin and gliclazide in combination with metformin in patients with Type 2 diabetes inadequately controlled with metformin alone: a 52-week, randomized study. *Diabetic medicine : a journal of the British Diabetic Association*. 2010;27(3):318-26.

37. Fonseca V, Rosenstock J, Patwardhan R, Salzman A. Effect of metformin and rosiglitazone combination therapy in patients with type 2 diabetes mellitus: a randomized controlled trial. *Jama*. 2000;283(13):1695-702.
38. Forst T, Uhlig-Laske B, Ring A, Graefe-Mody U, Friedrich C, Herbach K, et al. Linagliptin (BI 1356), a potent and selective DPP-4 inhibitor, is safe and efficacious in combination with metformin in patients with inadequately controlled Type 2 diabetes. *Diabetic medicine : a journal of the British Diabetic Association*. 2010;27(12):1409-19.
39. Gallwitz B, Bohmer M, Segiet T, Molle A, Milek K, Becker B, et al. Exenatide twice daily versus premixed insulin aspart 70/30 in metformin-treated patients with type 2 diabetes: a randomized 26-week study on glycemic control and hypoglycemia. *Diabetes care*. 2011;34(3):604-6.
40. Gallwitz B, Guzman J, Dotta F, Guerci B, Simo R, Basson BR, et al. Exenatide twice daily versus glimepiride for prevention of glycaemic deterioration in patients with type 2 diabetes with metformin failure (EUREXA): an open-label, randomised controlled trial. *Lancet*. 2012;379(9833):2270-8.
41. Gallwitz B, Rosenstock J, Rauch T, Bhattacharya S, Patel S, von Eynatten M, et al. 2-year efficacy and safety of linagliptin compared with glimepiride in patients with type 2 diabetes inadequately controlled on metformin: a randomised, double-blind, non-inferiority trial. *Lancet*. 2012;380(9840):475-83.
42. Gao Y, Yoon KH, Chuang LM, Mohan V, Ning G, Shah S, et al. Efficacy and safety of exenatide in patients of Asian descent with type 2 diabetes inadequately controlled with metformin or metformin and a sulphonylurea. *Diabetes research and clinical practice*. 2009;83(1):69-76.
43. Garber A, Klein E, Bruce S, Sankoh S, Mohideen P. Metformin-glibenclamide versus metformin plus rosiglitazone in patients with type 2 diabetes inadequately controlled on metformin monotherapy. *Diabetes, obesity & metabolism*. 2006;8(2):156-63.
44. Goke B, Gallwitz B, Eriksson J, Hellqvist A, Gause-Nilsson I, Investigators DC. Saxagliptin is non-inferior to glipizide in patients with type 2 diabetes mellitus inadequately controlled on metformin alone: a 52-week randomised controlled trial. *Int J Clin Pract*. 2010;64(12):1619-31.
45. Gomez-Perez FJ, Fanghanel-Salmon G, Antonio Barbosa J, Montes-Villarreal J, Berry RA, Warsi G, et al. Efficacy and safety of rosiglitazone plus metformin in Mexicans with type 2 diabetes. *Diabetes/metabolism research and reviews*. 2002;18(2):127-34.
46. Goodman M, Thurston H, Penman J. Efficacy and tolerability of vildagliptin in patients with type 2 diabetes inadequately controlled with metformin monotherapy. *Hormone and metabolic research = Hormon- und Stoffwechselforschung = Hormones et metabolisme*. 2009;41(5):368-73.
47. Goudswaard AN, Stolk RP, Zuithoff P, de Valk HW, Rutten GE. Starting insulin in type 2 diabetes: continue oral hypoglycemic agents? A randomized trial in primary care. *The Journal of family practice*. 2004;53(5):393-9.
48. Halimi S, Le Berre MA, Grange V. Efficacy and safety of acarbose add-on therapy in the treatment of overweight patients with Type 2 diabetes inadequately controlled with metformin: a double-blind, placebo-controlled study. *Diabetes research and clinical practice*. 2000;50(1):49-56.
49. Hamann A, Garcia-Puig J, Paul G, Donaldson J, Stewart M. Comparison of fixed-dose rosiglitazone/metformin combination therapy with sulphonylurea plus metformin in

- overweight individuals with Type 2 diabetes inadequately controlled on metformin alone. *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association*. 2008;116(1):6-13.
50. Hartemann-Heurtier A, Halbron M, Golmard JL, Jacqueminet S, Bastard JP, Rouault C, et al. Effects of bed-time insulin versus pioglitazone on abdominal fat accumulation, inflammation and gene expression in adipose tissue in patients with type 2 diabetes. *Diabetes research and clinical practice*. 2009;86(1):37-43.
 51. Heine RJ, Van Gaal LF, Johns D, Mihm MJ, Widel MH, Brodows RG. Exenatide versus insulin glargine in patients with suboptimally controlled type 2 diabetes: a randomized trial. *Annals of internal medicine*. 2005;143(8):559-69.
 52. Herman WH, Buse JB, Arakaki RF, Dungan KM, Jiang HH, Jacobson JG, et al. Concomitant oral antihyperglycemic agent use and associated treatment outcomes after initiation of insulin therapy. *Endocrine practice : official journal of the American College of Endocrinology and the American Association of Clinical Endocrinologists*. 2011;17(4):563-7.
 53. Hermansen K, Kipnes M, Luo E, Fanurik D, Khatami H, Stein P. Efficacy and safety of the dipeptidyl peptidase-4 inhibitor, sitagliptin, in patients with type 2 diabetes mellitus inadequately controlled on glimepiride alone or on glimepiride and metformin. *Diabetes, obesity & metabolism*. 2007;9(5):733-45.
 54. Holman RR, Thorne KI, Farmer AJ, Davies MJ, Keenan JF, Paul S, et al. Addition of biphasic, prandial, or basal insulin to oral therapy in type 2 diabetes. *The New England journal of medicine*. 2007;357(17):1716-30.
 55. Home PD, Pocock SJ, Beck-Nielsen H, Curtis PS, Gomis R, Hanefeld M, et al. Rosiglitazone evaluated for cardiovascular outcomes in oral agent combination therapy for type 2 diabetes (RECORD): a multicentre, randomised, open-label trial. *Lancet*. 2009;373(9681):2125-35.
 56. Janka HU, Plewe G, Busch K. Combination of oral antidiabetic agents with basal insulin versus premixed insulin alone in randomized elderly patients with type 2 diabetes mellitus. *Journal of the American Geriatrics Society*. 2007;55(2):182-8.
 57. Janka HU, Plewe G, Riddle MC, Kliebe-Frisch C, Schweitzer MA, Yki-Jarvinen H. Comparison of basal insulin added to oral agents versus twice-daily premixed insulin as initial insulin therapy for type 2 diabetes. *Diabetes care*. 2005;28(2):254-9.
 58. Kaku K. Efficacy and safety of therapy with metformin plus pioglitazone in the treatment of patients with type 2 diabetes: a double-blind, placebo-controlled, clinical trial. *Current medical research and opinion*. 2009;25(5):1111-9.
 59. Kendall DM, Riddle MC, Rosenstock J, Zhuang D, Kim DD, Fineman MS, et al. Effects of exenatide (exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulfonylurea. *Diabetes care*. 2005;28(5):1083-91.
 60. Khanolkar MP, Morris RH, Thomas AW, Bolusani H, Roberts AW, Geen J, et al. Rosiglitazone produces a greater reduction in circulating platelet activity compared with gliclazide in patients with type 2 diabetes mellitus--an effect probably mediated by direct platelet PPARgamma activation. *Atherosclerosis*. 2008;197(2):718-24.
 61. Kilo C, Mezitis N, Jain R, Mersey J, McGill J, Raskin P. Starting patients with type 2 diabetes on insulin therapy using once-daily injections of biphasic insulin aspart 70/30, biphasic human insulin 70/30, or NPH insulin in combination with metformin. *Journal of diabetes and its complications*. 2003;17(6):307-13.

62. Ko GT, Tsang PC, Wai HP, Kan EC, Chan HC. Rosiglitazone versus bedtime insulin in the treatment of patients with conventional oral antidiabetic drug failure: a 1-year randomized clinical trial. *Advances in therapy*. 2006;23(5):799-808.
63. Kvapil M, Swatko A, Hilberg C, Shestakova M. Biphasic insulin aspart 30 plus metformin: an effective combination in type 2 diabetes. *Diabetes, obesity & metabolism*. 2006;8(1):39-48.
64. Lam KS, Tiu SC, Tsang MW, Ip TP, Tam SC. Acarbose in NIDDM patients with poor control on conventional oral agents. A 24-week placebo-controlled study. *Diabetes care*. 1998;21(7):1154-8.
65. Lopez-Alvarenga JC, Aguilar-Salinas CA, Velasco-Perez ML, Arita-Melzer O, Guillen LE, Wong B, et al. Acarbose vs. bedtime NPH insulin in the treatment of secondary failures to sulphonylurea-metformin therapy in type 2 diabetes mellitus. *Diabetes, obesity & metabolism*. 1999;1(1):29-35.
66. Marre M, Howlett H, Lehert P, Allavoine T. Improved glycaemic control with metformin-glibenclamide combined tablet therapy (Glucovance) in Type 2 diabetic patients inadequately controlled on metformin. *Diabetic Medicine*. 2002;19(8):673-80.
67. Marre M, Van Gaal L, Usadel KH, Ball M, Whatmough I, Guitard C. Nateglinide improves glycaemic control when added to metformin monotherapy: results of a randomized trial with type 2 diabetes patients. *Diabetes, obesity & metabolism*. 2002;4(3):177-86.
68. Matthews DR, Charbonnel BH, Hanefeld M, Brunetti P, Schernthaner G. Long-term therapy with addition of pioglitazone to metformin compared with the addition of gliclazide to metformin in patients with type 2 diabetes: a randomized, comparative study. *Diabetes/metabolism research and reviews*. 2005;21(2):167-74.
69. Milicevic Z, Hancu N, Car N, Ivanyi T, Schwarzenhofer M, Jermendy G. Effect of two starting insulin regimens in patients with type II diabetes not controlled on a combination of oral antihyperglycemic medications. *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association*. 2009;117(5):223-9.
70. Moses R, Slobodniuk R, Boyages S, Colagiuri S, Kidson W, Carter J, et al. Effect of repaglinide addition to metformin monotherapy on glycemic control in patients with type 2 diabetes. *Diabetes care*. 1999;22(1):119-24.
71. Nauck M, Frid A, Hermansen K, Shah NS, Tankova T, Mitha IH, et al. Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes: the LEAD (liraglutide effect and action in diabetes)-2 study. *Diabetes care*. 2009;32(1):84-90.
72. Nauck MA, Duran S, Kim D, Johns D, Northrup J, Festa A, et al. A comparison of twice-daily exenatide and biphasic insulin aspart in patients with type 2 diabetes who were suboptimally controlled with sulfonylurea and metformin: a non-inferiority study. *Diabetologia*. 2007;50(2):259-67.
73. Nauck MA, Ellis GC, Fleck PR, Wilson CA, Mekki Q. Efficacy and safety of adding the dipeptidyl peptidase-4 inhibitor alogliptin to metformin therapy in patients with type 2 diabetes inadequately controlled with metformin monotherapy: a multicentre, randomised, double-blind, placebo-controlled study. *Int J Clin Pract*. 2009;63(1):46-55.
74. Nauck MA, Hompesch M, Filipczak R, Le TD, Zdravkovic M, Gumprecht J, et al. Five weeks of treatment with the GLP-1 analogue liraglutide improves glycaemic control and

- lowers body weight in subjects with type 2 diabetes. *Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association*. 2006;114(8):417-23.
75. Nauck MA, Meininger G, Sheng D, Terranella L, Stein PP, Sitagliptin Study G. Efficacy and safety of the dipeptidyl peptidase-4 inhibitor, sitagliptin, compared with the sulfonylurea, glipizide, in patients with type 2 diabetes inadequately controlled on metformin alone: a randomized, double-blind, non-inferiority trial. *Diabetes, obesity & metabolism*. 2007;9(2):194-205.
 76. Ovalle F, Bell DS. Effect of rosiglitazone versus insulin on the pancreatic beta-cell function of subjects with type 2 diabetes. *Diabetes care*. 2004;27(11):2585-9.
 77. Owens DR, Swallow R, Dugi KA, Woerle HJ. Efficacy and safety of linagliptin in persons with type 2 diabetes inadequately controlled by a combination of metformin and sulphonylurea: a 24-week randomized study. *Diabetic medicine : a journal of the British Diabetic Association*. 2011;28(11):1352-61.
 78. Pan C, Xing X, Han P, Zheng S, Ma J, Liu J, et al. Efficacy and tolerability of vildagliptin as add-on therapy to metformin in Chinese patients with type 2 diabetes mellitus. *Diabetes, obesity & metabolism*. 2012;14(8):737-44.
 79. Papanthassiou K, Naka KK, Kazakos N, Kanioglou C, Makriyiannis D, Pappas K, et al. Pioglitazone vs glimepiride: Differential effects on vascular endothelial function in patients with type 2 diabetes. *Atherosclerosis*. 2009;205(1):221-6.
 80. Pfutzner A, Schondorf T, Tschope D, Lobmann R, Merke J, Muller J, et al. PIOfix-study: effects of pioglitazone/metformin fixed combination in comparison with a combination of metformin with glimepiride on diabetic dyslipidemia. *Diabetes technology & therapeutics*. 2011;13(6):637-43.
 81. Phillips P, Karrasch J, Scott R, Wilson D, Moses R. Acarbose improves glycemic control in overweight type 2 diabetic patients insufficiently treated with metformin. *Diabetes care*. 2003;26(2):269-73.
 82. Poon T, Nelson P, Shen L, Mihm M, Taylor K, Fineman M, et al. Exenatide improves glycemic control and reduces body weight in subjects with type 2 diabetes: a dose-ranging study. *Diabetes technology & therapeutics*. 2005;7(3):467-77.
 83. Pratley RE, Nauck M, Bailey T, Montanya E, Cuddihy R, Filetti S, et al. Liraglutide versus sitagliptin for patients with type 2 diabetes who did not have adequate glycaemic control with metformin: a 26-week, randomised, parallel-group, open-label trial. *Lancet*. 2010;375(9724):1447-56.
 84. Raskin PR, Hollander PA, Lewin A, Gabbay RA, Bode B, Garber AJ. Basal insulin or premix analogue therapy in type 2 diabetes patients. *European journal of internal medicine*. 2007;18(1):56-62.
 85. Raz I, Chen Y, Wu M, Hussain S, Kaufman KD, Amatruda JM, et al. Efficacy and safety of sitagliptin added to ongoing metformin therapy in patients with type 2 diabetes. *Current medical research and opinion*. 2008;24(2):537-50.
 86. Reynolds LR, Kingsley FJ, Karounos DG, Tannock LR. Differential effects of rosiglitazone and insulin glargine on inflammatory markers, glycemic control, and lipids in type 2 diabetes. *Diabetes research and clinical practice*. 2007;77(2):180-7.
 87. Ristic S, Collober-Maugeais C, Cressier F, Tang P, Pecher E. Nateglinide or gliclazide in combination with metformin for treatment of patients with type 2 diabetes mellitus

- inadequately controlled on maximum doses of metformin alone: 1-year trial results. *Diabetes, obesity & metabolism*. 2007;9(4):506-11.
88. Ristic S, Collober-Maugeais C, Pecher E, Cressier F. Comparison of nateglinide and gliclazide in combination with metformin, for treatment of patients with Type 2 diabetes mellitus inadequately controlled on maximum doses of metformin alone. *Diabetic medicine : a journal of the British Diabetic Association*. 2006;23(7):757-62.
 89. Rodger NW, Chiasson JL, Josse RG, Hunt JA, Palmason C, Ross SA, et al. Clinical experience with acarbose: results of a Canadian multicentre study. *Clinical and investigative medicine Medecine clinique et experimentale*. 1995;18(4):318-24.
 90. Rosenstock J, Brown A, Fischer J, Jain A, Littlejohn T, Nadeau D, et al. Efficacy and safety of acarbose in metformin-treated patients with type 2 diabetes. *Diabetes care*. 1998;21(12):2050-5.
 91. Rosenstock J, Sugimoto D, Strange P, Stewart JA, Soltes-Rak E, Dailey G. Triple therapy in type 2 diabetes: insulin glargine or rosiglitazone added to combination therapy of sulfonylurea plus metformin in insulin-naive patients. *Diabetes care*. 2006;29(3):554-9.
 92. Ross SA, Zinman B, Campos RV, Strack T. A comparative study of insulin lispro and human regular insulin in patients with type 2 diabetes mellitus and secondary failure of oral hypoglycemic agents. *Clinical and investigative medicine Medecine clinique et experimentale*. 2001;24(6):292-8.
 93. Russell-Jones D, Vaag A, Schmitz O, Sethi BK, Lalic N, Antic S, et al. Liraglutide vs insulin glargine and placebo in combination with metformin and sulfonylurea therapy in type 2 diabetes mellitus (LEAD-5 met+SU): a randomised controlled trial. *Diabetologia*. 2009;52(10):2046-55.
 94. Schernthaner G, Grimaldi A, Di Mario U, Drzewoski J, Kempler P, Kvapil M, et al. Guide study: Double-blind comparison of once-daily gliclazide MR and glimepiride in type 2 diabetic patients. *European Journal of Clinical Investigation*. 2004;34(8):535-42.
 95. Scott R, Loeys T, Davies MJ, Engel SS, Sitagliptin Study G. Efficacy and safety of sitagliptin when added to ongoing metformin therapy in patients with type 2 diabetes. *Diabetes, obesity & metabolism*. 2008;10(10):959-69.
 96. Secnik Boye K, Matza LS, Oglesby A, Malley K, Kim S, Hayes RP, et al. Patient-reported outcomes in a trial of exenatide and insulin glargine for the treatment of type 2 diabetes. *Health and quality of life outcomes*. 2006;4:80.
 97. Standl E, Schernthaner G, Rybka J, Hanefeld M, Raptis SA, Naditch L. Improved glycaemic control with miglitol in inadequately-controlled type 2 diabetics. *Diabetes research and clinical practice*. 2001;51(3):205-13.
 98. Stehouwer MH, DeVries JH, Lumeij JA, Ader HJ, Engbers AM, Iperen Av A, et al. Combined bedtime insulin--daytime sulphonylurea regimen compared with two different daily insulin regimens in type 2 diabetes: effects on HbA1c and hypoglycaemia rate--a randomised trial. *Diabetes/metabolism research and reviews*. 2003;19(2):148-52.
 99. Strojek K, Bebakar WM, Khutsoane DT, Pesic M, Smahelova A, Thomsen HF, et al. Once-daily initiation with biphasic insulin aspart 30 versus insulin glargine in patients with type 2 diabetes inadequately controlled with oral drugs: an open-label, multinational RCT. *Current medical research and opinion*. 2009;25(12):2887-94.
 100. Taskinen MR, Rosenstock J, Tamminen I, Kubiak R, Patel S, Dugi KA, et al. Safety and efficacy of linagliptin as add-on therapy to metformin in patients with type 2 diabetes: a

- randomized, double-blind, placebo-controlled study. *Diabetes, obesity & metabolism*. 2011;13(1):65-74.
101. Umpierrez G, Issa M, Vlahovic A. Glimepiride versus pioglitazone combination therapy in subjects with type 2 diabetes inadequately controlled on metformin monotherapy: results of a randomized clinical trial. *Current medical research and opinion*. 2006;22(4):751-9.
 102. Van Gaal L, Maislos M, Schernthaner G, Rybka J, Segal P. Miglitol combined with metformin improves glycaemic control in type 2 diabetes. *Diabetes, obesity & metabolism*. 2001;3(5):326-31.
 103. von Bibra H, Diamant M, Scheffer PG, Siegmund T, Schumm-Draeger PM. Rosiglitazone, but not glimepiride, improves myocardial diastolic function in association with reduction in oxidative stress in type 2 diabetic patients without overt heart disease. *Diabetes & vascular disease research : official journal of the International Society of Diabetes and Vascular Disease*. 2008;5(4):310-8.
 104. Wang JS, Lin SD, Lee WJ, Su SL, Lee IT, Tu ST, et al. Effects of acarbose versus glibenclamide on glycemic excursion and oxidative stress in type 2 diabetic patients inadequately controlled by metformin: a 24-week, randomized, open-label, parallel-group comparison. *Clinical therapeutics*. 2011;33(12):1932-42.
 105. Wolever TM, Chiasson JL, Josse RG, Hunt JA, Palmason C, Rodger NW, et al. Small weight loss on long-term acarbose therapy with no change in dietary pattern or nutrient intake of individuals with non-insulin-dependent diabetes. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*. 1997;21(9):756-63.
 106. Yang W, Chen L, Ji Q, Liu X, Ma J, Tandon N, et al. Liraglutide provides similar glycaemic control as glimepiride (both in combination with metformin) and reduces body weight and systolic blood pressure in Asian population with type 2 diabetes from China, South Korea and India: a 16-week, randomized, double-blind, active control trial(*). *Diabetes, obesity & metabolism*. 2011;13(1):81-8.
 107. Yang W, Guan Y, Shentu Y, Li Z, Johnson-Levonas AO, Engel SS, et al. The addition of sitagliptin to ongoing metformin therapy significantly improves glycemic control in Chinese patients with type 2 diabetes. *Journal of diabetes*. 2012;4(3):227-37.
 108. Yang W, Pan CY, Tou C, Zhao J, Gause-Nilsson I. Efficacy and safety of saxagliptin added to metformin in Asian people with type 2 diabetes mellitus: a randomized controlled trial. *Diabetes research and clinical practice*. 2011;94(2):217-24.
 109. Yki-Jarvinen H, Kauppinen-Makelin R, Tiikkainen M, Vahatalo M, Virtamo H, Nikkila K, et al. Insulin glargine or NPH combined with metformin in type 2 diabetes: the LANMET study. *Diabetologia*. 2006;49(3):442-51.