

Supplementary file for

Enhanced cellular uptake of phenamil through inclusion complex with histidine functionalized β -Cyclodextrin as a penetrative osteo-inductive agent

Vahid Jahed^a, Ebrahim Vasheghani Farahani^{a*}, Fatemeh Bagheri^b, Ali Zarrabi^c, Trine Fink^d, Kim Lambertsen Larsen^{e*}

^a Biomedical Engineering Division, Faculty of Chemical engineering, Tarbiat Modares University, Tehran, Iran


^b Department of Biotechnology, Faculty of Chemical engineering, Tarbiat Modares University, Tehran, Iran

^c Sabanci University Nanotechnology Research and Application Center (SUNUM), Tuzla, 34956, Istanbul, Turkey


^d Department of Health Science and Technology, Faculty of Health, Aalborg University, Aalborg, Denmark

^e Department of Chemistry and Bioscience, Faculty of Chemistry, Aalborg University, Aalborg, Denmark


* Corresponding authors emails: Ebrahim Vasheghani Farahani: evf@modares.ac.ir, Kim Lambertsen Larsen kll@bio.aau.dk




S 1. ^1H NMR spectra of A) βCD , and B) Amine- βCD in $\text{dms}\text{-d}_6$ to confirm selective reaction on primary hydroxyl (integral calibration was done on H1 peak)



S 2. TLC plates of Amine- β CD conversion to His- β CD after 3, 5 and 7 hours. Arrow shows Amine- β CD before of reaction, R and R' assigned to low and high concentration of reaction respectively.



S 3. FTIR spectra of His-βCD deprotection.



S 4. ¹H NMR spectra of A) His-βCD-PH, B) HP-βCD-PH. Due to the small amount of PH, weak peaks in the range of 7 to 8 ppm related to benzene ring of PH were appeared