

Supplementary data

Table S1 Hospital wise details of patients with multi drug resistant organisms (MDRO) infections and outcomes

Hospitals	No. of patients with COVID-19 (n)	No. of patients with secondary infections (n/%)	No. of patients with MDROs (n/%)	Outcomes in MDRO infected (n/%)
H1	1089	19 (1.7%)	3 (15.7%)	Death:3 (100%)
H2	1552	145 (9.3%)	103 (71.0%)	Death: 73 (70.8%) Recovered: 30
H3	4590	104 (2.2%)	21 (20.1%)	Death:17 (80.9%) Recovered: 4
H4	612	58 (9.4%)	25 (45.4%)	Death:14 (56.0%) Recovered: 11
H5	1082	38 (3.5%)	-	--
H6	5132	63 (1.2%)	41 (65.0%)	Death:30 (73.1%) Recovered: 11
H7	2598	11 (0.4%)	4 (36.3%)	Death:1 (25.0%) Recovered: 3
H8	459	85 (18.5%)	28 (32.9%)	Death:18 (64.2%) Recovered: 10
H9	70	17 (24.2%)	-	--
H10	350	100 (28.5%)	77 (77.0%)	Death:44 (57.1%) Recovered: 33

Table S2 Hospital wise details of sample collection for culture of patients diagnosed with SIs

Hospitals	Total patients	Total samples sent for culture	Cultures sent per patient	Total no. of patients with SIs	% patients diagnosed with SIs
H1	1089	129	0.11	19	1.7
H2	1552	2060	1.32	145	10.0
H3	4590	2373	0.51	104	2.2
H4	612	390	0.63	58	9.4
H5	1082	217	0.2	38	3.5
H6	5132	98	0.01	63	1.2
H7	2598	374	0.14	11	0.4
H8	459	570	1.24	85	18
H9	70	229	3.2	17	24
H10	350	723	2.0	100	28.5




Figure S1 Correlation between culture collected and patients diagnosed with secondary infections