Supplementary material

Statistical analysis results regarding the correlation between serum GFAP levels and the degree of neurological deficit based on the NIHSS score in ischemic stroke patients in this study

Nama	Umur	L/P	Merokok	DM	Dislipidemia	НТ	ΣFR	NIHSS	GFAP (ng/mL)
01	81	L	v		v	v	3	8	0.3
02	65	L	v			v	2	8	0.5
03	59	L	v			v	2	5	0.5
04	51	Р				v	1	4	0.3
05	57	L	v		v	v	3	4	0.7
06	41	Р			v	v	2	8	1
07	31	L	v				1	8	0.3
08	37	L	v	v		v	3	4	0.2
09	54	Р			v	v	2	5	0.5
10	59	L		v		v	2	8	0.4
11	65	L	v		v	v	3	9	0.3
12	52	L	v		v	v	3	8	0.4
13	68	Р			v	v	2	8	0.7
14	65	Р			v	v	2	9	0.4
15	66	L	v				1	7	0.7
16	62	Р			v	v	2	6	0.7
17	71	Р				v	1	6	0.3
18	45	L				v	1	15	0.7
19	91	Р				v	1	15	0.5
20	60	L				v	1	17	0.8
21	64	L	v	v	v	v	4	16	1.0
22	64	L		v		v	2	15	0.7
23	57	Р			v	v	2	14	0.5
24	53	L	v	v		v	3	20	0.5
25	71	L	v		v	v	2	18	0.7
26	73	L	v		v		2	19	0.5
27	62	L	v			v	2	20	0.6
28	72	Р		v	v	v	3	21	1.9
29	66	L				v	1	16	1.8
30	75	Р			v	v	2	18	0.7
31	63	L	v		v	v	2	21	1.0
32	65	Р	v	v		v	3	15	0.8
33	59	Р		v		v	2	16	1

Keterangan: FR: Faktor Risiko; DM: Diabetes Melitus; HT: Hipertensi; LAO: Large Artery Occlusion, NIHSS: National Institutes of Health Stroke Scale; SAO: Small Artery Occlusion