



Supplementary Table 1. Primer sequences used in the assessment of gene expression by RT-qPCR (human nucleotide sequence).

Gene	Forward sequence (5'-3')	Reverse sequence (5'-3')
CD80	CTGCCTGACCTACTGCTTIG	GGCGTACACTTTCCCTTCTC
CD163	ACATAGATCATGCATCTGTCATTTG	ATTCTCGGAATCTCACTTCTA
CD200R	GAGCAATGGCACAGTGACTGTT	GTGGCAGGTCACGGTAGACA
β -actin	AGAGCTACGAGCTGCCTGAC	AGCACTGTGTTGGCGTACAG
TNF- α	AGGCGCTCCCAAGAAGACA	TCCTTGGCAAACTGCACCT
IL-1 β	GCAGTCTACACAGCTTCGGG	CCGCCTCAGCCTCCCAAAG
IL-6	GGCTGCTCCTGGTGTTC	TCTGAAGAGGTGAGTGGCTGTC
IL-8	CAGTTTGGCCAAGAAGTGCTAAAG	GGGTGGAAGGTTTGGAGTATGTC
IL-10	TTGCTGGAGGACTTTAAGGGTTAC	CCTGATGTCTGGGTCTTGGTTCTC
IL-12	CCAGAGCAGTGAGGTCTTAGGC	GTGAAGCAGCAGGAGCGAATG



Supplementary Figure 1 Polarizing agents-induced cytotoxicity of macrophages in function of PMA concentrations. Cytotoxicity of M1 (dots plots) or M2 (empty plots) polarizing agents assessed by crystal violet test after 48h or 72h respectively. * $p < 0,05$; ** $p < 0.01$ using Student's t-test analysis as compared with PMA alone.



Supplementary Figure 2. Distribution of Foslip in macrophages in spheroids by flow cytometry.