

1 SUPPORTING INFORMATION

2 **Microcurrent cloth-assisted transdermal penetration and**
3 **follicular ducts escape of curcumin-loaded micelles for**
4 **enhanced wound healing**

5 Pei-Chi Lee^{1,†}, Cun-Zhao Li^{2,†}, Chun-Te Lu^{3,†}, Ming-Han Zhao², Syu-Ming Lai², Man-Hua Liao⁴,
6 Cheng-Liang Peng⁵, Hsin-Tung Liu¹, Ping-Shan Lai^{1*}

7

8 ¹ xTrans Corporate Research and Innovation Center, Taipei City, Taiwan

9 ² Department of Chemistry, National Chung Hsing University, Taichung, Taiwan

10 ³ Plastic Surgery, Taichung Veterans General Hospital, Taichung, Taiwan

11 ⁴ Graduate Institute of Biotechnology, National Chung Hsing University, Taichung, Taiwan

12 ⁵ Isotope Application Division, Institute of Nuclear Energy Research, Taoyuan, Taiwan

13

14 Correspondence: Ping-Shan Lai

15 Department of Chemistry, National Chung Hsing University, No. 145, Xingda Road, Taichung 402,
16 Taiwan.

17 Tel: +886-4-22840411 ext. 428.

18 Fax: +886-4-22862547.

19 E-mail address: pslai@email.nchu.edu.tw

20

21 † These authors contributed equally to this work.


22

23

24

25

26



27

28 **Figure S1.** Aqueous solubility of Cur in its native form and in CM-L. Samples containing 50 mg of
 29 curcumin were added to 2 mL eppendorf tubes with 1.5 mL water and incubated in a water bath at
 30 37°C overnight. The supernatant was collected and then analyzed by using HPLC. Data are
 31 expressed as the mean \pm SD (n = 3). Statistically significant differences are indicated by *(p < 0.05).

32 **Abbreviations:** Cur: curcumin; CM-L: curcumin-loaded micelle with large particle size; HPLC: high-
 33 performance liquid chromatography. SD: standard deviation.

34