

Cost-effectiveness analysis of memantine for moderate-to-severe Alzheimer's disease in the Netherlands

Bart Hoogveldt¹
Benoît Rive²
Johan Severens³
Khaled Maman⁴
Chantal Guilhaume⁵

¹Field Product Management, Lundbeck BV, Amsterdam, The Netherlands;

²Global Outcomes Research, Lundbeck SAS, Issy-les-Moulineaux, France; ³Institute of Health Policy and Management, Erasmus University Rotterdam, The Netherlands; ⁴Creativ Research SAS, Paris, France; ⁵Global Evidence and Value Development, Sanofi-Aventis, Paris, France

Objective: The purpose of this study was to estimate the cost-effectiveness of memantine relative to standard care in patients with moderate-to-severe Alzheimer's disease in the Netherlands.

Methods: A country-adapted five-year Markov model simulated disease progression through a series of states, defined by dependency and disease severity. Transition probabilities were derived from trials, with utility and epidemiological data obtained from a longitudinal Dutch cohort. Cost-effectiveness was described in terms of quality-adjusted life years and time spent in a nondependent state or in a moderate severity state.

Results: Memantine monotherapy versus standard care led to 0.058 quality-adjusted life years gained (1.207 versus 1.265), longer time in a nondependent state (from 1.602 to 1.751 years) and in a moderate state (from 2.051 to 2.141 years), and no additional costs (€113,927 versus €110,097). Robustness of results was confirmed through sensitivity analyses.

Conclusion: Memantine is dominant compared with standard care in the Netherlands. Results are consistent with similar economic evaluations in other countries.

Keywords: memantine, Alzheimer's disease, cost-effectiveness analysis, Netherlands, cholinesterase inhibitors

Introduction

Alzheimer's disease is the most common cause of dementia, accounting for two-thirds of all dementia cases. People with Alzheimer's disease experience gradual loss of independence, and as Alzheimer's disease becomes more severe, the burden on informal caregivers increases until institutionalization is required. A study in the Netherlands showed that dementia patients spend approximately 20–25% of their time institutionalized with the condition.¹

Previous pharmacoeconomic studies conducted in the Netherlands have investigated galantamine, an acetylcholinesterase inhibitor, for mild-to-moderate Alzheimer's disease and the general costs associated with the disease.^{2,3}

Memantine is a moderate-affinity, noncompetitive N-methyl-D-aspartate antagonist, and is approved in over 70 countries worldwide for the symptomatic treatment of patients with moderate-to-severe Alzheimer's disease. Memantine may be administered as monotherapy or in combination with an acetylcholinesterase inhibitor. In this study, a Markov model was used to determine the cost-effectiveness of memantine relative to standard care in moderate-to-severe Alzheimer's disease in the Netherlands.

Correspondence: Benoît Rive
Lundbeck SAS, 43-45 Quai du Président
Roosevelt, 92445 Issy-les-
Moulineaux Cedex, France
Tel +331 7941 2912
Fax +331 7941 2908
Email berv@lundbeck.com

Methods

This cost-effectiveness study evaluated the five-year outcomes associated with memantine monotherapy compared with no pharmacological treatment (standard care) in terms of additional time living independently, additional time in a moderate state, quality-adjusted life years, and societal costs. A Markov model adapted from a previous model constructed for the Canadian setting was used, and full methodological details are provided in Gagnon et al.⁴ The methods employed here are identical to the Canadian model unless explicitly stated otherwise. Key changes are the data source for the initial patient distribution, background mortality, cost per health state, and discount rates. Each of these inputs was adjusted to make the model directly relevant to the economic assessment of memantine in the Netherlands as described by the College Voor Zorgverzekeringen (CVZ).⁵

Model structure

In this model, health states were based on dependence and Alzheimer's disease severity, ie, a patient could be in one of five possible states: moderate, independent; moderate, dependent; severe, independent; severe, dependent; death. A six-month Markov cycle was chosen and repeated 10 times.

Perspective

This study was performed from a societal perspective, with nonmedical costs included, in line with the CVZ guidelines for pharmacoeconomic research.

Data sources

The initial patient distributions between Markov health states were computed from the Rotterdam study.⁶ This was a large study testing 7528 residents (2939 men and 4589 women) aged 55 years and over (median age 69 years) of a Rotterdam suburb for disorders associated with dementia. The Rotterdam study found a subset of 339 patients diagnosed with Alzheimer's disease, from which the health state distributions were calculated.⁷ The percentages of patients by severity stage and level of dependency were: moderate severity, independent (69.02%); moderate, dependent (20.11%); severe, independent (6.52%); and severe, dependent (4.35%).

All severity and dependency transition probabilities were assumed to depend only on patient severity and dependency stages at the beginning of the cycle and on treatment, and were estimated using pooled data from four six-month, double-blind, randomized, placebo-controlled clinical trials for memantine considering a total of 1096

patients with moderate-to-severe Alzheimer's disease (589 receiving memantine and 507 receiving placebo).⁸⁻¹¹ The efficacy of memantine was assumed to last for the first 12 months of treatment, as suggested by an open-label extension study.¹²

Probability of death over each six-month cycle was assumed to be the same for all Markov health states and treatment strategies, and was computed from published Dutch data.³ This resulted in an estimated probability of death of 7.55% for each six-month period when data were adjusted using the weighted mean of probabilities. This was based on the aforementioned initial model distribution for dependency status and an assumption (based on expert opinion) that, for patients in an institution, the split between patients living in a home for the elderly and patients living in a nursing home was 2/3:1/3 in independent patients and 1/2:1/2 in dependent patients. The uncertainty surrounding these parameters was modeled using a priori triangular distributions.

The costs per unit of care were derived from the Dutch guidelines.¹³ The number of units used per health state was estimated from a Dutch model of cost of care for patients with dementia.¹⁴ The costs of care and consumption by dependency and type of accommodation were calculated in Euros (€, value 1999, Table 1). Because no distinction between genders was made in this study, it was assumed when calculating total cost per stage that approximately 66% of the patients with Alzheimer's disease were female, based on dementia prevalence estimates among men and women from the Rotterdam study.⁶

The distribution of moderate and severe Alzheimer's disease patients in the Rotterdam study and the split between patients living in a home for the elderly and patients living in a nursing home were applied to costs per dependency and accommodation (Table 1), to compute the costs per Markov health state. Costs were updated to 2008 using the consumer price data from Statistics Netherlands restricted to the health sector. Final six-month cost estimates per Markov health state were 8,092.12 (EUR, 2008) for moderate independent patients, 26,202.87 for moderate dependent, 9,475.97 for severe independent and 27,022.75 for severe dependent. The price used for memantine in the Netherlands was €2.95 per defined-daily-dose of 20 mg. An additional pharmacist fee of €6.10 per prescription was added. The cost of the memantine treatment over a cycle was €537.10.

Level of dependency and residential status appear to be the main criteria influencing quality-adjusted life years values, and thus values were taken from the LASER-AD study from the UK that considered 224 Alzheimer's disease

Table 1 Annual costs and consumption (€, value 1999) by dependency and accommodation

Stage	Accommodation	Type of care	Costs/unit*	Units [†]	Costs	
Independent	Community	General practitioner	16.59 Euro/consult	6.09 consults	101.03 Euro	
		Family care	17.70 Euro/hour	93.59 hours	1656.54 Euro	
		District-nursing	31.76 Euro/hour	17.51 hours	556.12 Euro	
		RIAGG	102.55 Euro/contact	–	–	
		Hospital admission ¹	283.84 Euro/day	–	–	
		Day-care center (including travelling costs)	52.18 Euro/day	–	–	
		Informal care	7.94 Euro/hour	260 hours	2064.40 Euro	
		Total	–	–	4378.09 Euro	
		Home for the elderly ²	Total	30.33 Euro/day	365 days	11070.45 Euro
		Nursing home ²	Total	99.43 Euro/day ³	365 days	36291.95 Euro
Dependent	Community	General practitioner	16.59 Euro/hour	6.09 consults	101.03 Euro	
		Family care	17.70 Euro/hour	183.22 hours	3242.99 Euro	
		District nursing	31.76 Euro/hour	213.53 hours	6781.71 Euro	
		RIAGG	102.55 Euro/contact	9.4 contacts	963.97 Euro	
		Hospital admission	283.84 Euro/day	41.65 days	11821.94 Euro	
		Day-care center (including travelling costs)	52.18 Euro/day	92 days	4800.56 Euro	
		Informal care	7.94 Euro/hour	322 hours	2556.68 Euro	
		Total	–	–	30268.88 Euro	
		Home for the elderly ²	Total	94.01 Euro/day	365 days	34313.65 Euro
		Nursing home ²	Total	171.02 Euro/day ³	365 days	62422.30 Euro

Notes: ¹Mean academic and general hospital; ²These do not include housing costs; ³Mean psychogeriatric and combined nursing homes; *Derived from Oostenbrink et al;¹³ [†]Derived from Van der Roer.¹⁴

Abbreviation: RIAGG, regional institutes for mental health.

patients (independent in community, 0.608 [standard error (SE) 0.028]; independent in institution, 0.543 [SE 0.064]; dependent in community, 0.340 [SE 0.058]; dependent in institution, 0.169 [SE 0.058]).¹⁵

Following Dutch guidelines, health outcomes were discounted at an annual rate of 1.5%, and costs outcomes at 4%.

Analyses

The base-case scenario reported the mean number of years of independence and moderate severity state, quality-adjusted life years and average costs over five years. Probabilistic analyses using Monte-Carlo simulations comprised of 10,000 draws were conducted, which attributed a value for each of the a priori distributions. The a priori distributions were beta distributions for severity and dependency transition probabilities, log-normal for the odds ratios measuring impact of memantine on severity and dependency transitions, normal distributions for utilities per health states, triangular for the nursing home/home for elderly proportions, and triangular ($\pm 20\%$) for costs per health states. A worst-case scenario was included whereby it was assumed that patients received treatment for two years, but the treatment effect only lasted for six months. One-way sensitivity analyses were conducted on the odds ratios of memantine applied to dependency and severity probabilities. A sensitivity analysis was also conducted on the discount rates

(applied to both costs and health benefits) by using the values 0% and 7% per annum in the model. Finally, a sensitivity analysis was conducted on the nursing home/home for elderly distributions.

Results

Compared with the standard care strategy, over five years memantine treatment produced an additional 0.149 years (54.39 days, 95% credible intervals [CrI] = 0.036; 0.252) of independence, 0.091 years (33.22 days, 95% CrI = -0.027; 0.197) in the moderate disease state, and 0.058 (95% CrI = 0.014; 0.108) additional quality-adjusted life years (Table 2). Furthermore, memantine treatment was associated with a cost reduction of €3,830 compared with standard care. Memantine was the dominant strategy being more effective and less costly than standard care. There was a 96.8% probability of memantine being dominant.

Results were robust to sensitivity analyses. For the worst-case scenario, the memantine strategy was associated with an additional 0.078 years (28.47 days; 95% CrI = 0.019; 0.132) of independence, 0.048 years (17.52 days; 95% CrI = -0.015; 0.104) in the moderate disease state, and 0.031 additional quality-adjusted life years (95% CrI = 0.007; 0.057). Memantine was associated with a cost reduction of €920 over the five-year evaluation period compared with

Table 2 The cost-effectiveness model outcomes of memantine compared with standard care

	Memantine		Standard care	
	Mean	2.5%; 97.5% CrI	Mean	2.5%; 97.5% CrI
Distribution of model outcomes and costs per strategy				
QALYs	1.265	0.999; 1.527	1.207	0.940; 1.471
Years in independent state	1.751	1.463; 2.039	1.602	1.335; 1.878
Years in moderate state	2.141	1.852; 2.415	2.051	1.780; 2.322
Costs (€, value 2006)	110,097	96,599; 124,223	113,927	100,364; 128,146
Distribution of incremental model outcomes and costs associated with memantine				
QALYs	0.058	0.014; 0.108		
Years in independent state	0.149	0.036; 0.252		
Years in moderate state	0.091	-0.027; 0.197		
Costs (€, value 2006)	-3,830	-7,793; 208		

Abbreviations: CrI, credible interval; QALYs, quality-adjusted life years.

standard care. The probability of memantine being dominant was 80.7% in this scenario.

Discussion

This study has shown that over a five-year period in a Dutch setting, compared with standard care in patients with moderate-to-severe Alzheimer's disease, memantine was associated with additional quality-adjusted life years, prolonged time of independence and prolonged time in the moderate Alzheimer's disease state.

These health benefits translated into cost savings, which fully offset the drug cost, making memantine the dominant option. The internal validity and robustness of the model were ensured through numerous sensitivity analyses conducted around key parameters. A major assumption is that the structure of the Canadian model and several inputs are considered generalizable without transferability adaptation.¹⁶

This study confirmed results from previous cost-effectiveness analyses performed in other countries such as the UK, Finland, Norway, Sweden, Spain, and Canada. In each of these evaluations, the additional cost of memantine was offset by health benefits which translated into cost savings. Results were consistent, despite differences in model design, perspective, data sources, geographical location, and time horizon. The estimated gain in terms of months of independence (increase of 9.3% for 1.8 months) in this study was similar to other evaluations of memantine. Although not directly comparable, it is interesting to note that in an analysis of galantamine for Alzheimer's disease in a number of countries (including the Netherlands), galantamine delayed time to full-time care by 6.8%.¹⁷ As with all pharmacoeconomic analyses, this cost-effectiveness evaluation is subject to a number of limitations because it extrapolates data from the clinical trial setting to model real-life practice and is associated with a number of assumptions.

Conclusion

This analysis demonstrated that memantine monotherapy produced health benefits compared with standard care which translated into cost savings. Memantine was associated with additional quality-adjusted life years and an extended period of nondependence. These results for the Netherlands are consistent with economic evaluations of memantine in Canada and a number of other countries in Europe.

Acknowledgment

The authors thank Monique Breteler of Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands, for her contribution to this research.

Disclosure

Chantal Guilhaume was an employee of H Lundbeck A/S at the time the analysis was conducted. Otherwise, the authors report no conflicts of interest in this work.

References

1. Perenboom RJ, Boshuizen HC, Breteler MM, Ott A, Van de Water HP. Dementia-free life expectancy (DemFLE) in The Netherlands. *Soc Sci Med.* 1996;43:1703-1707.
2. Caro JJ, Salas M, Ward A, Getsios D, Mehnert A. Economic analysis of galantamine, a cholinesterase inhibitor, in the treatment of patients with mild to moderate Alzheimer's disease in the Netherlands. *Dement Geriatr Cogn Disord.* 2002;14:84-89.
3. McDonnell J, Redekop WK, Van der Roer N, et al. The cost of treatment of Alzheimer's disease in The Netherlands: A regression-based simulation model. *Pharmacoeconomics.* 2001;19:379-390.
4. Gagnon M, Rive B, Hux M, Guilhaume C. Cost-effectiveness of memantine compared with standard care in moderate-to-severe Alzheimer disease in Canada. *Can J Psychiatry.* 2007;52:519-526.
5. College voor Zorgverzekeringen. *Guidelines for Pharmacoeconomic Research, Updated Version.* Diemen, The Netherlands: College voor Zorgverzekeringen; 2006.
6. Breteler MM. A community-based study of dementia: The Rotterdam Elderly Study. *Neuroepidemiology.* 1992;11 Suppl 1:S23-S28.
7. Ott A, Breteler M, Van Harskamp F, et al. Prevalence of Alzheimer's disease and vascular dementia: Association with education. The Rotterdam study. *BMJ.* 1995;310:970-973.

8. Reisberg B, Doody R, Stöffler A, Schmitt F, Ferris S, Möbius HJ. Memantine in moderate-to-severe Alzheimer's disease. *N Engl J Med*. 2003;348:1333–1341.
9. Peskind ER, Potkin SG, Pomara N, et al. Memantine treatment in mild to moderate Alzheimer disease: A 24-week randomized controlled trial. *Am J Geriatr Psychiatry*. 2006;14:704–715.
10. Bakchine S, Pascual-Gangnant L, Loft H. Results of a randomised, placebo-controlled 6-month study of memantine in the treatment of mild to moderate Alzheimer's disease in Europe. Abstract presented at the Ninth Congress of the European Federation of the Neurological Societies (EFNS), September 17–20, 2005, Athens, Greece.
11. Van Dyck CH, Tariot PN, Meyers B, Malca Resnick E. A 24-week randomized, controlled trial of memantine in patients with moderate-to-severe Alzheimer disease. *Alzheimer Dis Assoc Disord*. 2007;21:136–143.
12. Reisberg B, Doody R, Stoffler A, Schmitt F, Ferris S, Mobius HJ. A 24-week open-label extension study of memantine in moderate to severe Alzheimer disease. *Arch Neurol*. 2006;63:49–54.
13. Oostenbrink JB, Koopmanschap MA, Rutten FFH. Handleiding voor Kostenonderzoek. Amstelveen, The Netherlands: College voor Zorgverzekeringen; 2000.
14. Van der Roer N. Model of costs of care for dementia: community-dwelling vs institutionalization. *Tijdschr Gerontol Geriatr*. 2000;31:55–61. Dutch.
15. Sapin C, Livingston G, Katona C. Main factors influencing preference-based utility values in Alzheimer's disease: Results from the LASER-AD Study. Poster presented at the Seventh International Conference on Alzheimer's and Parkinson's Disease (AD/PD), March 9–13, 2005, Sorrento, Italy.
16. Drummond M, Barbieri M, Cook J, et al. Transferability of economic evaluations across jurisdictions: ISPOR Good Research Practices Task Force report. *Value Health*. 2009;12:409–418.
17. Caro J, Salas M, Ward A, Getsios D, Migliaccio-Walle K, Garfield F. Assessing the health and economic impact of galantamine treatment in patients with Alzheimer's disease in the health care systems of different countries. *Drugs Aging*. 2004;21:677–686.

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PscINFO' database and CAS, and is the official

journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>