Comorbidity and confounding factors in attention-deficit/hyperactivity disorder and sleep disorders in children

Ya-Wen Jan^{1,2}

Chien-Ming Yang^{1,3}

Yu-Shu Huang^{4,5}

¹Department of Psychology, National Cheng-Chi University, Taipei; ²Sleep Center of Taipei Medical University Hospital, Taipei; ³The Research Center for Mind Brain and Learning, National Cheng-Chi University, Taipei; ⁴Department of Child Psychiatry and Sleep Center, Chang Gung Memorial Hospital, Taoyuan; ⁵College of Medicine, Chang Gung University, Taoyuan, Taiwan

Abstract: Sleep problems are commonly reported in children with attention-deficit/hyperactivity disorder (ADHD) symptoms. Research data regarding the complex and reciprocal relationship between ADHD and sleep disturbances has now accumulated. This paper is focused on the types of sleep problems that are associated with ADHD symptomatology, and attempts to untangle confounding factors and overlapping symptoms. The goal is also to present an updated overview of the pathophysiology of and treatment strategies for sleep problems in children with ADHD. The review also points out that future research will be needed to clarify further the other psychiatric comorbidities and side effects of medication in order to improve treatment outcomes and prevent misdiagnosis in clinical practice.

Keywords: attention-deficit/hyperactivity disorder, sleep, children

Introduction

Attention-deficit/hyperactivity disorder (ADHD) is one of the most commonly studied and diagnosed psychiatric disorders in children, and is characterized by inattention, overactivity, and impulsivity.¹ ADHD is estimated to affect about 3%–5% of children globally,² although prevalence rates may vary depending on whether samples are obtained from community or clinical settings.^{3,4} Complaints of sleep problems in children with ADHD are not uncommon. In 1957, Laufer and Denhoff⁵ first reported the association between sleep disturbances and ADHD. They noted “Generally, the parents of hyperkinetic children are so desperate over the night problems that the daytime ones pale in significance”. Restless and disturbed sleep were, at one time, part of the Diagnostic and Statistical Manual of Mental Disorders (DSM) diagnostic criteria for ADHD, although the symptoms were eliminated from subsequent DSM editions.^{1,6,7} In recent years, there has been growing interest in sleep disorders associated with ADHD. Manifestations of primary sleep disorders may mimic ADHD symptomatology, so might lead to misdiagnosis in clinical practice. Moreover, the primary sleep disorders may exacerbate underlying ADHD symptoms, which might interfere with the treatment and impair daytime performance. In addition, psychotropic medications that are used to treat ADHD may also result in sleep problems, such as sleep initiation difficulty. Therefore, the management of sleep problems in children with ADHD may significantly reduce behavioral symptom severity and improve the quality of life of these children, as well as that of their families.⁸

However, variations in the inclusion criteria and measurements across the studies have impeded our understanding of the ways in which sleep and ADHD interact. This

Correspondence: Yu-Shu Huang
Department of Child Psychiatry
and Sleep Center, Chang Gung Memorial
Hospital, Linkou,
Taiwan No 5, Fu-Shing Street,
Kwei-Shan, Taoyuan 333,
Taipei, Taiwan
Tel +886 3328 1200 ext 8541
Fax +886 3318 4541
Email hu1109s@yahoo.com.tw

review attempts to clarify the potential confounding factors of previous research regarding the relationships between sleep problems and ADHD in children, and outlines some treatment considerations in the management of sleep disturbances in ADHD. We performed a PubMed search from 1994 (when the American Psychiatric Association first published the DSM-IV) to May 2011, using the following key words in various combinations: “ADHD”, “attention-deficit/hyperactivity disorder”, “sleep”, “sleep disorder”, and “children”. References from each relevant paper, including 14 existing reviews of the literature, were examined to determine if any relevant studies had been missed during the database searches. Case reports and descriptive reports without data analysis were excluded in our review.

Diagnosis and comorbidity in ADHD

Some of the published studies have reported that the prevalence of sleep disturbances and sleep disorders among children with ADHD are significantly higher than in normal controls, but others failed to replicate these findings.^{9,10} One possible reason for the inconsistent results may be that the methods used to define ADHD differed across studies, ranging from subjective rating scales for symptoms of hyperactivity/impulsivity and inattention to clinical judgments made by experienced physicians who were following DSM diagnostic criteria. The studies in which ADHD was defined by questionnaires might be confounded by factors such as psychiatric comorbidity.¹¹ Many studies that applied DSM diagnostic criteria did not exclude or control the effects of psychiatric comorbidity as well. Comorbid disorders frequently associated with ADHD, such as tic disorder,¹² oppositional defiant disorder,⁹ depressive disorders,¹³ and anxiety disorders,^{13–15} may account for some of the sleep disturbances.

Subjective and objective measures of sleep in ADHD children

Subjective reports of sleep in children with ADHD have primarily been collected through parental-report questionnaires and sleep logs. The data collected usually include: sleep quality and quantity, with variables (Table 1) such as difficulty in sleep initiation and/or sleep onset latency, total sleep time, wake after sleep onset, number of night-time awakenings, and sleep efficiency; sleep habits and bedtime behaviors, like sleep resistance; and symptoms of sleep disorders, including sleep-related breathing disorders and movement disorders. However, the reliability and validity of subjective sleep measurement may be questionable. It has been reported that the sleep data assessed by the Child Behavior Checklist, a

Table 1 Definition of the sleep parameters

Sleep parameters	Definition
Wake after sleep onset (WASO)	Amount of total wake time between sleep-onset and final wake-up
Total sleep time (TST)	Amount of actual sleep time in a sleep period
Sleep onset latency (SOL)	Time period measured from light-off to the beginning of sleep
Sleep efficiency (SE)	Ratio of total sleep time to time in bed
Apnea-hypopnea index (AHI)	An index used to assess severity of sleep apnea based on total number of complete cessations (apnea) and partial obstructions (hypopnea) of breathing occurring per hour of sleep
Periodic limb movement index (PLMI)	An index used to assess severity of periodic limb movement disorder based on total number of periodic limb movements occurring per hour of sleep

measure frequently used to evaluate sleep disturbance among children with ADHD, was incongruent with the data from sleep logs, actigraphy, and polysomnography.

Objective measurements used to study sleep and ADHD include polysomnography and actigraphy. Polysomnography allows researchers to examine basic sleep parameters, sleep architecture (eg, total sleep time, wake after sleep onset, sleep onset latency, sleep efficiency), and pathological indices (eg, apnea-hypopnea index, periodic limb movement index). However, polysomnography requires subjects to sleep within laboratories, which may produce a sleep pattern that differs from their habitual sleep patterns. Prihodova et al¹⁶ pointed out that the “first-night effect” on sleep variables was more significant in the ADHD group than normal controls. Therefore, it might be necessary to record sleep for more than one night to ensure validity of the data. On the other hand, actigraphy assesses sleep patterns through the recording and analyses of activity patterns. It can provide sleep information, such as sleep onset, sleep offset, sleep onset latency, and wake after sleep onset. The biggest advantage of actigraphy is its capacity to collect data from more than 20 consecutive nights in natural settings. Nevertheless, it does not record pathophysiological indices that are necessary for the diagnosis of some sleep disorders. In addition to the disruption of night-time sleep, excessive daytime sleepiness is one of the major clinical characteristics in patients with sleep disorders. Hence, the multiple sleep latency test is also frequently used to measure daytime sleepiness in ADHD.^{16–18} The multiple sleep latency test assesses the physiologic sleep tendency by measuring the time it takes to fall asleep when one begins a nap in the absence of alerting factors, such as noise and light.¹⁹

Table 2 Published studies utilizing subjective measures of sleep in ADHD

Study	Sample size (n)	Age (years)	Study design	Medication	Measurement	Major findings
ADHD symptoms						
O'Brien et al ³¹	44 significant ADHD symptoms 53 mild ADHD symptoms 39 controls	5–7	Cross-sectional, community-based	NA	Parent report	Subjective sleep disturbances, such as difficulty initiating sleep, restless sleep, night-time awakenings, snoring, and willingness to fall asleep were frequently reported among significant ADHD symptoms in children.
Gau ³²	2463	6–15	Cross-sectional, community-based	NA	Parent and teacher report SHQ, CPRS-R:S, and CTRS-R:S	Dysomnia, SDB, and daytime sleepiness were related to mother's and teacher's reports on child's symptoms of inattention, and hyperactivity/impulsivity. However, parasomnia was only related to mother's reports on ADHD-related symptoms.
Willoughby et al ³⁸	1073	2–5	Cross-sectional, clinic-based	NA	Parent report CBCL	Relationship between hyperactive-impulsive symptomatology and sleep problems was attenuated once ODD/CD, anxiety, and depressive symptom counts were included as additional covariates.
ADHD diagnosis						
Owens et al ³³	46 ADHD 46 normal	5–10	Cross-sectional, clinic-based	Unmedicated	Parent and self-report Sleep self-report	Sleep disturbances, particularly at bedtime, are frequently reported by both parents and children with ADHD.
Ivanenko et al ³⁴	29 ADHD alone 50 ADHD comorbid with mood and anxiety disorders 67 mood and anxiety disorders alone 28 other psychiatric disorders 174 controls	5–18	Cross-sectional, clinic-based and community-based	NA	Parent report CSQ	ADHD group was reported to have longer sleep latency, more night-time awakenings, nightmares, snoring, restless sleep, bedtime resistance, and leg jerks compared with controls. However, sleep complaints were more likely associated with symptoms of mood disturbances and anxiety rather than derived from the presence of ADHD.
ADHD comorbid with other diagnosis						
Corkum et al ³⁵	25 ADHD 25 normals	7–11	Cross-sectional, community-based	Drug-naïve	Parent report CSQ-P	ADHD group was reported to have longer sleep duration, more difficulty with sleep onset, awakening in the morning, and bedtime resistance compared with the normal group.
Sung et al ³⁶	239 ADHD	5–18	Cross-sectional, clinic-based	Medicated	Parent report	Difficulty falling asleep, resisting going to bed, and tiredness on waking were extremely common in children with ADHD.
Lim et al ³⁷	114 ADHD 60 normals	5–13	Cross-sectional, clinic-based	Medicated	Parent report CBCL	Children with ADHD reported less sleep and more sleep-related problems compared with controls.

Abbreviations: ADHD, attention deficit/hyperactivity disorder; CD, compulsive disorder; ODD, oppositional defiant disorder; SDB, sleep-disordered breathing; SHQ, Sleep Habits Questionnaire; CTRS-R:S, Conners' Teacher Rating Scale-Revised/Short Form; CPRS-R:S, Conners' Parent Rating Scale-Revised/Short Form; CBCL, Child Behavior Checklist; CSQ, Children's Sleep Habits Questionnaire; CSQ-P, Child Sleep Questionnaire; Parent Version; NA, not applicable.

Table 3 Studies utilizing objective measures of sleep in ADHD

Study	Sample size (n)	Age (years)	Study design	Medication	Measurement	Major findings
ADHD symptoms						
Paavonen et al ⁴⁵	280 children	7–8	Cross-sectional, community-based	NA	Actigraph	Children's short sleep duration measured by actigraphs increase the risk for behavioral symptoms of ADHD.
O'Brien et al ³¹	44 significant ADHD symptoms 53 mild ADHD symptoms 39 controls	5–7	Cross-sectional, Community-based	NA	PSG	REM latency and proportion of REM sleep (%TST) were more likely to be affected in the group with significant ADHD symptoms.
ADHD diagnosis						
Gruber et al ³⁹	38 ADHD boys 64 normal boys	6–14	Cross-sectional, community-based	Medication-naïve	Actigraphy and Sleep diaries	No significant differences were found in SOL, WASO, SE, and TST between the ADHD and control groups.
Gruber and Sadeh ⁴⁰	24 ADHD boys 25 normal boys	7–11	Cross-sectional, community-based	Medication-naïve	Actigraphs	No significant differences were found between the ADHD and control groups on SOL, WASO, SE, and TST.
Dagan et al ⁴¹	12 ADHD 12 controls	6–12	Cross-sectional, clinic-based	Medicated	Actigraph	SOL and sleep duration were not significantly different between the two groups. However, quiet sleep percentage and SE were found to be significantly lower in ADHD children than in controls.
Gruber et al ⁴⁶	15 ADHD 23 controls	7–11	Cross-sectional, community-based	Unmedicated	PSG	Children with ADHD had significantly shorter sleep and REM sleep durations, and a smaller percentage of REM sleep of total sleep time compared with controls.
Kirov et al ¹²	22 healthy controls 25 ADHD 32 TD 24 ADHD + TD	8–16	Cross-sectional, clinic-based	Unmedicated	PSG	Children with ADHD had significant longer TST, shorter REM sleep, increased REM sleep percentage, and higher number of sleep cycles than children without ADHD.
O'Brien et al ⁴⁷	47 ADHD clinic 53 ADHD community 49 controls	5–9	Cross-sectional, clinic-based and community-based	Medicated	PSG	ADHD clinic group showed significantly lower spontaneous arousals, longer REM sleep latency, and lower REM sleep percentage than both ADHD community group and controls.
Picchietti et al ⁴⁸	14 ADHD 10 controls	5–12	Cross-sectional, clinic-based	Unmedicated	PSG	Duration of deep sleep (Stage 3 and 4), and REM sleep was decreased in ADHD compared with control group.
Prihodova et al ¹⁶	31 ADHD 26 controls	6–12	Cross-sectional, clinic-based	Unmedicated	PSG	Basic sleep macrostructure parameters revealed no differences between control and ADHD groups either the first or the second night.
Leclercq et al ¹⁸	33 ADHD boys 23 learning disorder boys	5–10	Cross-sectional, clinic-based	Unmedicated	PSG	No significant differences in TST, SOL, number of awakenings, and percentage of different stages were found between children with ADHD and controls.

Konofal et al ⁴⁹	30 ADHD boys 19 learning disorder boys	5–10	Cross-sectional, clinic-based	Unmedicated	PSG	TST, SOL, number of awakenings, and percentage of different stages in PSG did not differ significantly between the two groups.
Cooper et al ⁵⁰	18 ADHD 20 normal controls	4–16	Cross-sectional, clinic-based	Unmedicated	PSG	PSG showed normal arousal indexes, and AHI for the ADHD group and normal control group. The sleep architecture was not significantly different between groups.
ADHD comorbid with other diagnosis						
Corkum et al ³⁵	25 ADHD 25 normal controls	7–11	Cross-sectional, clinic-based	Medication-naïve	Actigraphy and sleep diary	There were no group differences on TST, WASO, SOL, and night-time awakenings.
Hvolby et al ⁴²	45 ADHD 64 psychiatric control group 97 health reference group	5–12	Cross-sectional, clinic-based	Medicated	Actigraphy	There was a significant difference in SOL between the three groups. There were no group differences for TST, WASO, and night-time awakenings.
Wiggs et al ⁴³	71 ADHD 23 healthy controls	3–15	Cross-sectional, clinic-based	Medication-free	Actigraphy and sleep diary	The results suggested no significant differences between the groups for wake time, wake episodes, and sleep schedule.
Owens et al ⁴⁴	80 ADHD 45 healthy controls	6–14	Cross-sectional, community-based	Unmedicated	Actigraphy and sleep diary	Compared with the control group, the ADHD group experienced shorter actual sleep time of all epochs scored as sleep, significantly fewer sleep interruptions, but more total interrupted sleep time.
Golan et al ¹⁷	34 ADHD 32 healthy controls	7–17	Cross-sectional, clinic-based	Unmedicated	PSG	Sleep architecture did not differ significantly between the groups, with the exception of higher percentage of REM sleep in the ADHD group.
O'Brien et al ⁴⁷	47 ADHD clinic 53 ADHD community 49 controls	5–7	Cross sectional, clinic-based and community-based	Medicated	PSG	There were statistically significant differences between the ADHD community group and the control group in REM sleep latency and percentage of REM.
Miano et al ⁵¹	20 ADHD 20 normal controls	6–13	Cross-sectional, clinic-based	Medication-naïve	PSG	Children with ADHD showed significantly reduced sleep duration, increased rate of stage shifts, lower REM sleep percentage, and lower sleep efficiency.
Silvestri et al ⁵²	55 ADHD 20 healthy controls	6–11	Cross-sectional, clinic-based	NA	PSG	Significant difference in percentage of REM, N2%, N3%, SE, TST, and REM latency between ADHD children and controls.
Kirov et al ⁵³	17 ADHD boys (12 with comorbid disease) 17 normal boys	8–14	Cross-sectional, clinic-based	Unmedicated	PSG	PSG data showed a significant increase in the duration of the absolute REM sleep and the number of sleep cycles in ADHD group when compared with controls.

Abbreviations: ADHD, attention deficit/hyperactivity disorder; TD, tic disorder; PSG, polysomnography; REM, rapid eye movement; SOL, sleep onset latency; WASO, wake after sleep onset; SE, sleep efficiency; TST, total sleep time; N2, sleep stage 2; N3, merged the third and fourth stage of sleep; NA, not applicable.

Table 4 Published studies investigating the relationship between ADHD and RLS in children

Study	Sample size (n)	Age (years)	Study design	Medication	Measurement	Major findings
ADHD symptoms						
Picchietti and Stevens ⁵⁴	18 RLS	8–24	Cross-sectional, clinic-based	NA	NIH-specific pediatric criteria	ADHD (72%) is a common comorbidity of RLS.
Konofal et al ⁵⁵	12 ADHD + RLS 10 ADHD 10 controls	5–8	Cross-sectional, clinic-based	Unmedicated	NIH-specific pediatric criteria CPRS	Children with ADHD and a positive family of RLS are at risk of severe ADHD symptoms.
Rajaram et al ⁵⁷	11 ADHD 10 controls	5–12	Cross-sectional, clinic-based	NA	Children and parent's reports	Some ADHD children with growing pains may actually have RLS.
Kotagal and Silber ⁵⁶	32 RLS	<18	Cross-sectional, clinic-based	NA	PSG	Inattentiveness was seen in 8 of 32 subjects (25%).
Chervin et al ⁵⁸	866	2–13	Cross-sectional, community-based	NA	PSQ CPRS	Inattention and hyperactivity in general pediatric patients associated with symptoms of RLS.
Chervin et al ⁵⁹	27 ADHD 43 non-ADHD psychiatry subjects 73 general pediatric subjects	2–18	Cross-sectional, clinic-based	NA	PSQ CPRS	Complaint of restless legs and a composite score for daytime sleepiness showed some evidence of an association with inattention and hyperactivity.
ADHD diagnosis						
Picchietti et al ⁴⁸	14 ADHD 10 control	5–12	Cross-sectional, clinic-based	Unmedicated	PSG	ADHD children's parents (32%) were more likely to have RLS than the control parents.

Abbreviations: ADHD, attention deficit/hyperactivity disorder; PSQ, Pediatric Sleep Questionnaire; CPRS, Conners' Parent Rating Scale; NA, not applicable; NIH, National Institutes of Health; PSG, polysomnography; RLS, restless legs syndrome.

In the previous review articles on sleep and ADHD,^{10,20} it was pointed out that there is a significant discrepancy between subjective and objective reports of sleep in children with ADHD. Owens²¹ and Konofal et al²² suggested that parents of children with behavioral disorders may be more likely to perceive and report higher levels of daytime and sleep-related disruptive behaviors (bedtime struggles, difficulty with morning waking) in a sort of “negative halo effect”. In addition, caregivers are more likely to be sensitive to the side effects of medication, and report sleep-related behavioral problems, such as prolonged sleep-onset latency.

Impact of ADHD medication on sleep

Stimulant medication is one of the most effective treatments attempted for the management of ADHD.²³ It seems reasonable to assume that the central nervous system (CNS) stimulants prescribed to treat ADHD may adversely affect sleep due to their pharmacological character. For example, both statistically and clinically significant negative impacts have been reported in total sleep time²⁴ and sleep onset latency^{24,25} in children with ADHD under stimulant medication compared with no-medication conditions. However, recent randomized, double-blind, placebo-controlled trials have demonstrated that CNS stimulants do not significantly increase sleep

disturbances in ADHD.^{26,27} For example, one study examined sleep-related side effects of medications for children with ADHD and showed that there was a small increase in incidence of prolonged sleep onset latency and shortened total sleep time when a non-CNS stimulant (atomoxetine) was given compared with when a CNS stimulant (methylphenidate) was given.²⁸ On the contrary, some beneficial effects were also noted in various subjective and objective measures of sleep. In addition, clonidine, a medication that is often prescribed to treat ADHD-related sleep disturbance, was found to alter the duration of a nonrapid eye movement (NREM)-rapid eye movement (REM) cycle.²⁹

All in all, CNS stimulants may cause decreased total sleep time and prolonged sleep onset, and also improve sleep quality by consolidating sleep, decreasing nocturnal awakenings, and increasing the percentage of REM sleep.³⁰ Furthermore, non-CNS stimulants also caused adverse effects on sleep but the effects were relatively mild.

General sleep disturbances in ADHD

The relationship between sleep disturbances and ADHD has been examined using different measures. In studies based on parents' reports (Table 2), children with ADHD symptoms were found to have more night waking, difficulty initiating sleep,

Table 5 Published studies investigating the relationship between ADHD and PLMS in children

Study	Sample size (n)	Age (years)	Study design	Medication	Measurement	Major findings
ADHD symptoms						
Chervin et al ⁵⁸	866	2–13	Cross-sectional, community-based	NA	PSQ CPRS	Inattention and hyperactivity in general pediatric patients were associated with symptoms of PLMS.
Chervin and Archbold ⁷⁴	113	2–18	Cross-sectional, clinic-based	Medicated	PSG CPRS	Rate of PLMS showed a linear association with hyperactivity among subjects with SDB, but no association among those subjects without SDB.
ADHD diagnosis						
Picchiatti et al ⁴⁸	14 ADHD 10 controls	5–12	Cross-sectional, clinic-based	Unmedicated	PSG	Prevalence of PLMS was higher in the children with ADHD than in controls.
ADHD comorbid with other diagnosis						
Silvestri et al ⁵²	55 ADHD 20 healthy controls	6–11	Cross-sectional, clinic-based	NA	Nocturnal video-PSG International RLS Rating Scale	Significant correlations emerged for International RLS Rating Scale scoring, PLMS indices, hyperactivity, opposition scores, and ADHD subtypes.
Kirk and Bohn ⁷⁶	591	NA	Cross-sectional, clinic-based	NA	PSG	Prevalence of PLMS in 28 children with pre-existing diagnosis of ADHD in our population was higher (7.1%) than group prevalence (5.6%).
Huang et al ⁷⁷	88 ADHD 27 healthy controls	6–12	Cross-sectional, clinic-based	NA	PSG	Nine (10.2%) of the ADHD group had ≥ 5 PLMS per hour, but none in the control group.
Golan et al ¹⁷	34 ADHD 32 healthy controls	7–17	Cross-sectional, clinic-based	Unmedicated	PSG	Five of the ADHD group had PLMS during sleep (15%) versus none in the control group.
Crabtree et al ⁷⁵	40 PLMD/ADHD 50 PLMD only 52 controls	5–7	Cross-sectional, Community-based and clinic-based samples	Medicated	PSG	Children with PLMD and ADHD had a significantly greater number of arousals associated with PLMS than children with PLMD only.

Abbreviations: ADHD, attention deficit/hyperactivity disorder; PSQ, Pediatric Sleep Questionnaire; CPRS, Conners' Parent Rating Scale; PLMD, periodic limb movement disorder; RLS, restless legs syndrome; PLMS, periodic limb movements; SDB, sleep-disordered breathing; PSG, polysomnography.

bedtime resistance, and daytime sleepiness compared with normal control subjects.^{31,32} Children with ADHD were also reported to have significantly more bedtime resistance, longer sleep onset latency, shorter sleep duration, more parasomnias, more night waking, and higher level daytime sleepiness by parent's report.^{33,34} Children who have ADHD comorbid with other disorders were also reported to have longer sleep duration, more difficulty with sleep onset, more bedtime resistance, more difficulties awakening in the morning, and more nocturnal activities during sleep,^{35,36} but another study reported that they sleep less compared with normal subjects.³⁷ However, one study reported that sleep complaints were more likely associated with symptoms of mood disturbance and anxiety rather than derived from the presence of ADHD.³⁴ Symptoms of hyperactivity and impulsivity were also reported to have no significant association with parent-reported sleep problems once psychiatric comorbidity was controlled for in another study.³⁸

In studies using actigraphy (Table 3), children with ADHD showed no significant difference in sleep onset latency and sleep efficiency compared with controls.^{39,40} However, sleep efficiency and the amount of "quiet" sleep was found to be significantly lower in medicated ADHD children than in controls.⁴¹ When looking at ADHD comorbid with other psychiatric diagnoses, results from actigraphy were less consistent. Most of these studies failed to find night-time awakenings that were clearly described in subjective reports.^{35,42,43} Only one study indicated that children who have ADHD comorbid with other psychiatric diagnoses experience longer total interrupted sleep time,⁴⁴ while another study showed that they experienced significantly longer sleep onset latency.⁴² In addition, another study indicated that healthy children with short nocturnal sleep duration as recorded by actigraphy were found to exhibit more symptoms of hyperactivity/impulsivity.⁴⁵

Table 6 Published studies investigating the relationship between ADHD and SDB in children

Study	Sample size	Age	Study design	Medication	Measurement	Major findings
ADHD symptoms						
Chervin et al ⁸¹	229	2–13	Longitudinal study, clinic-based	Medicated	PSQ CPRS	Snoring and other symptoms of SDB are strong risk factors for future emergence or exacerbation of hyperactive behavior.
Gottlieb et al ⁸⁰	3019	5	Cross-sectional, community-based	NA	CPRS	Children with SDB symptoms were more likely to have hyperactivity, inattention, and aggressiveness.
O'Brien et al ³¹	44 significant ADHD symptoms 53 mild ADHD symptoms 39 controls	5–7	Cross-sectional, community-based	NA	PSG	SDB was highly prevalent in children with mild ADHD symptoms but not in those with significant ADHD symptoms.
Chervin et al ⁷⁹	866	2–13	Cross-sectional, community-based	NA	PSQ CPRS	Inattention and hyperactivity were associated with snoring and other symptoms of SDB.
Chervin and Archbold ⁷⁴	113	2–18	Cross-sectional, clinic-based	Medicated	PSG CPRS	Hyperactivity showed no significant associations with the rate of apneas and hypopneas.
ADHD comorbid with other diagnosis						
Huang et al ⁸²	66 ADHD 20 healthy controls	6–12	Cross sectional, clinic based	Medicated	PSG	Inattention and hyperactivity reduced after treating OSA by adenotonsillectomy or treating ADHD with stimulant
Huang et al ⁷⁷	88 ADHD 27 healthy controls	6–12	Cross-sectional, clinic based	NA	PSG	ADHD had a higher AHI (56.8% AHI > 1, 19.3% AHI > 5) than healthy controls
O'Brien et al ⁴⁷	47 ADHD clinic 53 ADHD community 49 controls	5–7	Cross-sectional, clinic-based and community-based	Medicated	PSG	AHI and apnea index did not differ in the 3 groups

Abbreviations: ADHD, attention deficit/hyperactivity disorder; AHI, apnea-hypopnea index; PSQ, Pediatric Sleep Questionnaire; CPRS, Conners' Parent Rating Scale; OSA, obstructive sleep apnea; PSG, polysomnography; SDB, sleep disordered breathing.

Many studies have tried to compare results of actigraphy with those from subjective reports by parents. Most of them found that subjective reports tended to overestimate sleep disturbances.⁴² It is also possible that the sleep disturbances that come to the attention of parents are not best detected by actigraphy.⁴³ Results highlight the prominence of parent-reported sleep disturbances in children with ADHD and the need for clinicians to screen routinely for the presence of sleep disorders and to assess detailed sleep physiology whenever sleep disturbances are indicated.

With regard to studies using polysomnography (Table 3), one study showed that nonmedicated children with ADHD were significantly different in their total sleep time compared with the control group.⁴⁶ Three studies have reported inconsistent findings for REM duration.^{12,46,47} One study showed that the duration of stage 3 sleep was shorter in children with ADHD than in controls,⁴⁸ while four studies found no significant differences in sleep variables.^{16,18,49,50}

Compared with normal controls, children who have ADHD comorbid with other neurological and psychiatric disorders spent decreased time in bed,⁵¹ and had a shorter total sleep time,^{51,52} increased REM latency,⁴⁷ higher arousal index,^{17,52} increased inconsistency in their REM sleep duration,^{17,47,51–53} and lower sleep efficiency⁵² on polysomnography. Thus, the hypothesis that sleep disturbances and specific changes in sleep macrostructure are connected with pure ADHD does not seem to be supported in this review of the literature.

Even though sleep difficulties might be common in young people with ADHD, results were not consistent across various measures of sleep. There might be many underlying factors which influence the relationship between ADHD and sleep problems. Therefore, correct identification of such factors may facilitate appropriate management of sleep disturbances in this population. The following section reviews the specific primary sleep disorders identified commonly in children with ADHD.

Sleep disorders in ADHD

Restless legs syndrome

Restless legs syndrome (RLS) is characterized by uncomfortable leg or body part sensations with an irresistible urge to move the legs. Recent attention focused on the association between ADHD and sleep disorders has led to a series of studies examining RLS in children with ADHD symptoms^{54–59} and with confirmed ADHD⁴⁸ (Table 4). In a review of the literature,⁶⁰ it was concluded that approximately 44% of patients with ADHD have RLS or RLS symptoms, and up to 26% of patients with RLS were found to have ADHD or ADHD symptoms. With the exclusion of “motor restlessness” as part of the diagnostic criteria for RLS in 2003, there is no longer any overlap of the diagnostic criteria for ADHD and RLS,⁶¹ and it has become easier to differentiate these distinct but symptomatically similar conditions.

Several mechanisms have been proposed to explain the association between ADHD (or ADHD symptoms) and RLS.⁶⁰ In children, sleep disturbance may precede or overshadow the complaint of leg discomfort.⁵⁶ The quality and quantity of a child’s sleep is often diminished with RLS. This sleep deprivation resulting from RLS can cause moodiness, irritability, inattentiveness, fatigue, and hyperactivity.^{35,62,63} Therefore, RLS might lead to symptoms of ADHD through sleep disruption. Secondly, diurnal manifestations of RLS, hyperactivity and inattentiveness, might mimic ADHD symptoms or be misclassified as symptoms of hyperactivity.⁶⁴ A third possibility is that RLS and ADHD might share a common pathology, such as a dopaminergic deficit.⁵⁸ Such a possibility is supported by the previous open-label studies, showing that children with symptoms of RLS and ADHD could benefit from treatment with dopaminergic agents.^{65,66} However, the first double-blind study of a dopaminergic therapy found that L-dopa significantly improved symptoms of RLS and periodic limb movement disorder (PLMD) symptoms but not ADHD.⁶⁷ Because there were methodological limitations, including small sample size and inability to control for baseline differences in severity of ADHD symptoms, these results should be interpreted carefully and future studies are needed to address this issue further. The fourth explanation is that both ADHD and RLS are associated with iron deficiency.^{56,68,69} The findings described here are consistent with pathophysiological models of these disorders because iron is needed for dopamine synthesis.⁷⁰ While dopaminergic agents are not approved for use in children with RLS, another possible implication of the association between RLS and ADHD symptoms concerns iron supplement treatment. Incidentally, iron supplementation has been reported

to improve symptoms in children with ADHD with an iron-deficient diet.⁷¹ Moreover, Konofal et al⁷² have pointed out that iron supplementation appeared to improve ADHD symptoms in children with low serum ferritin levels.

Periodic limb movements

Periodic limb movements (PLMS) is objectively defined as the presence of repetitive movements of the toes, feet, legs, thighs and arms that last 0.5–5.0 seconds and occur at 5–90 second intervals during sleep. According to the reports, 80% of patients with RLS also have characteristic leg movements during sleep that meet the diagnostic criteria for PLMS.⁷³ Several investigators have reported excessive PLMS in association with ADHD symptoms^{58,74} (Table 5). The prevalence of PLMS in children diagnosed with ADHD is estimated to be as high as 7%–44%, compared with a prevalence of only 1.2%–2.0% in pediatric populations at risk for sleep disorders.^{48,52,75,76} Moreover, Huang et al⁷⁷ and Golan et al¹⁷ found on polysomnography that 10.2%–14.7% of children with ADHD comorbid with other diagnosis had PLMD compared with 0% of controls. Children with PLMD during sleep can have sleep fragmentation that can lead to increased risk of inattention and hyperactivity, which are two major symptoms of ADHD. Chervin et al⁵⁸ further reported a dose-dependent relationship between PLMS scores on the Pediatric Sleep Questionnaire and hyperactivity index on the Conners’ Parent Rating Scale-Short Form. Crabtree et al⁷⁵ also showed that children with PLMD and ADHD were more likely to have PLMS-related arousals than children with PLMD only. Thus, Crabtree et al⁷⁵ postulated that the link between ADHD and PLMD may be mediated by sleep fragmentation associated with PLMS-induced arousals and the presence of reduced REM sleep.

In addition, ADHD, RLS, and PLMS are all associated with low iron stores.^{68,78} The finding of relative iron deficiency is consistent with pathophysiological models of these disorders because iron is needed for dopamine synthesis. Walters et al⁶⁵ indicated the treatment of PLMD with dopaminergic agents in the absence of psychostimulant use resulted in marked improvements in PLMD and ADHD. Thus, both PLMS and ADHD have been shown to be related to dopamine production and metabolism.

Sleep-disordered breathing

Sleep-disordered breathing (SDB) in children includes a spectrum of disorders that vary in severity, ranging from obstructive sleep apnea at one end to primary snoring, ie,

snoring without ventilator abnormalities. Several studies have demonstrated the association between symptoms of SDB and ADHD-related symptoms,^{31,47,77,79–81} but others have failed to show the association^{74,82} (Table 6). The inconsistent results were probably due to the difference in criteria used to diagnosis SDB (apnea-hypopnea index <1 or <5), as well as the aggravating effect of SDB on ADHD. Repeated episodes of nocturnal hypoxia in SDB comprise an important causal factor for neurobehavioral deficits. Sleep fragmentation resulting from frequent nocturnal arousals may also lead to daytime symptoms.²¹ Both neurobehavioral deficits and daytime symptoms might mimic or exacerbate the cardinal symptoms of ADHD.

Regarding treatment strategies in this population, both Huang et al⁸² and Dillon et al⁸³ reported that behavioral morbidity in children with ADHD and SDB improved significantly more after adenotonsillectomy than after stimulant treatment. These results suggest that appropriate recognition and surgical treatment of underlying SDB in children with ADHD might prevent the need for long-term stimulant treatment.⁸² A recent longitudinal study has shown that improvements were maintained 2.5 years after surgery.⁸⁴

Conclusion

ADHD-related symptoms are frequently associated with sleep problems/disorders. In our literature review, sleep disturbances are less supported by objective sleep measurements in pure unmedicated ADHD. Therefore, evaluation of other psychiatric comorbidities and the side effects of medications is necessary when assessing the causes of their sleep complaints. Moreover, sleep disturbance is very complex, especially in ADHD children. It is critical for health providers caring for this population to identify correctly the exact abnormality in sleep disturbance within this group of children. Success in treating sleep disturbance in this population depends on accurate diagnosis of the exact sleep disturbance and its pathology. Our review shows that successful management of sleep problems/disorders results in considerable improvement in the level of daytime impairment. Future studies addressing the causal relationship between sleep problems and ADHD-related symptoms should take into account potential confounding variables and include better controls. In addition, research focused on surgical management (adenotonsillectomy) and iron supplementation should not only focus on reduction of the apnea-hypopnea index and the periodic limb movement index as the primary parameters of a successful treatment outcome, but also consider other measures of sleep disturbance.

Disclosure

The authors report no conflicts of interest in this work.

References

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. Washington, DC: American Psychiatric Association; 1994.
2. Nair J, Ehimare U, Beitman BD, Nair SS, Lavin A. Clinical review: Evidence-based diagnosis and treatment of ADHD in children. *Mo Med*. 2006;103(6):617–621.
3. Barkley RA. *Attention Deficit Hyperactivity Disorder: A Handbook for Diagnosis and Treatment*. New York, NY: Guilford; 1990.
4. Brown RT, Freeman WS, Perrin JM, et al. Prevalence and assessment of attention-deficit/hyperactivity disorder in primary care settings. *Pediatrics*. 2001;107(3):E43. Available from: <http://pediatrics.aappublications.org/content/107/3/e43.full.pdf+html>. Accessed July 22, 2011.
5. Laufer MW, Denhoff E. Hyperkinetic behavior syndrome in children. *J Pediatr*. 1957;50(4):463–474.
6. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 3rd ed. Washington, DC: American Psychiatric Association; 1980.
7. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 3rd ed. Washington, DC: American Psychiatric Association; 1987.
8. Owens JA. The ADHD and sleep conundrum: A review. *J Dev Behav Pediatr*. 2005;26(4):312–322.
9. O'Brien LM, Gozal D. Sleep in children with attention deficit/hyperactivity disorder. *Minerva Pediatr*. 2004;56(6):585–601.
10. Cohen-Zion M, Ancoli-Israel S. Sleep in children with attention-deficit hyperactivity disorder (ADHD): A review of naturalistic and stimulant intervention studies. *Sleep Med Rev*. 2004;8(5):379–402.
11. Johnson JH, Alvarez HK, Loftis C. ADHD comorbidity and mimicry: Implications for assessment and treatment. In: Maria BL, editor. *Current Management in Child Neurology*. 3rd ed. Hamilton, Ontario: BC Decker; 2005.
12. Kirov R, Kinkelbur J, Banaschewski T, Rothenberger A. Sleep patterns in children with attention-deficit/hyperactivity disorder, tic disorder, and comorbidity. *J Child Psychol Psychiatry*. 2007;48(6):561–570.
13. Mayes SD, Calhoun SL, Bixler EO, et al. ADHD subtypes and comorbid anxiety, depression, and oppositional-defiant disorder: Differences in sleep problems. *J Pediatr Psychol*. 2009;34(3):328–337.
14. Alfano CA, Ginsburg GS, Kingery JN. Sleep-related problems among children and adolescents with anxiety disorders. *J Am Acad Child Adolesc Psychiatry*. 2007;46(2):224–232.
15. Mick E, Biederman J, Jetton J, Faraone SV. Sleep disturbances associated with attention deficit hyperactivity disorder: The impact of psychiatric comorbidity and pharmacotherapy. *J Child Adolesc Psychopharmacol*. 2000;10(3):223–231.
16. Prihodova I, Paclt I, Kemlink D, Skibova J, Ptacek R, Nevsimalova S. Sleep disorders and daytime sleepiness in children with attention-deficit/hyperactivity disorder: A two-night polysomnographic study with a multiple sleep latency test. *Sleep Med*. 2010;11(9):922–928.
17. Golan N, Shahar E, Ravid S, Pillar G. Sleep disorders and daytime sleepiness in children with attention-deficit/hyperactive disorder. *Sleep*. 2004;27(2):261–266.
18. Lecendreux M, Konofal E, Bouvard M, Falissard B, Mouren-Simeoni MC. Sleep and alertness in children with ADHD. *J Child Psychol Psychiatry*. 2000;41(6):803–812.
19. Carskadon MA, Dement WC, Mitler MM, Roth T, Westbrook PR, Keenan S. Guidelines for the multiple sleep latency test (MSLT): A standard measure of sleepiness. *Sleep*. 1986;9(4):519–524.
20. Corkum P, Tannock R, Moldofsky H. Sleep disturbances in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry*. 1998;37(6):637–646.

21. Owens JA. Sleep disorders and attention-deficit/hyperactivity disorder. *Curr Psychiatry Rep.* 2008;10(5):439–444.
22. Konofal E, Lecendreux M, Cortese S. Sleep and ADHD. *Sleep Med.* 2010;11(7):652–658.
23. Clinical practice guideline: Treatment of the school-aged child with attention-deficit/hyperactivity disorder. *Pediatrics.* 2001;108(4):1033–1044.
24. Corkum P, Panton R, Ironside S, Macpherson M, Williams T. Acute impact of immediate release methylphenidate administered three times a day on sleep in children with attention-deficit/hyperactivity disorder. *J Pediatr Psychol.* 2008;33(4):368–379.
25. Stein MA. Unravelling sleep problems in treated and untreated children with ADHD. *J Child Adolesc Psychopharmacol.* 1999;9(3):157–168.
26. Galland BC, Tripp EG, Taylor BJ. The sleep of children with attention deficit hyperactivity disorder on and off methylphenidate: A matched case-control study. *J Sleep Res.* 2010;19(2):366–373.
27. Faraone SV, Glatt SJ, Bukstein OG, Lopez FA, Arnold LE, Findling RL. Effects of once-daily oral and transdermal methylphenidate on sleep behavior of children with ADHD. *J Atten Disord.* 2009;12(4):308–315.
28. Sangal RB, Owens J, Allen AJ, Sutton V, Schuh K, Kelsey D. Effects of atomoxetine and methylphenidate on sleep in children with ADHD. *Sleep.* 2006;29(12):1573–1585.
29. Miyazaki S, Uchida S, Mukai J, Nishihara K. Clonidine effects on all-night human sleep: Opposite action of low- and medium-dose clonidine on human NREM-REM sleep proportion. *Psychiatry Clin Neurosci.* 2004;58(2):138–144.
30. Huang YS, Tsai MH, Guillemainault C. Pharmacological treatment of ADHD and the short and long term effects on sleep. *Curr Pharm Des.* April 8, 2011. [Epub ahead of print].
31. O'Brien LM, Holbrook CR, Mervis CB, et al. Sleep and neurobehavioral characteristics of 5- to 7-year-old children with parentally reported symptoms of attention-deficit/hyperactivity disorder. *Pediatrics.* 2003;111(3):554–563.
32. Gau SS. Prevalence of sleep problems and their association with inattention/hyperactivity among children aged 6–15 in Taiwan. *J Sleep Res.* 2006;15(4):403–414.
33. Owens JA, Maxim R, Nobile C, McGuinn M, Msall M. Parental and self-report of sleep in children with attention-deficit/hyperactivity disorder. *Arch Pediatr Adolesc Med.* 2000;154(6):549–555.
34. Ivanenko A, Crabtree VM, O'Brien LM, Gozal D. Sleep complaints and psychiatric symptoms in children evaluated at a pediatric mental health clinic. *J Clin Sleep Med.* 2006;2(1):42–48.
35. Corkum P, Tannock R, Moldofsky H, Hogg-Johnson S, Humphries T. Actigraphy and parental ratings of sleep in children with attention-deficit/hyperactivity disorder (ADHD). *Sleep.* 2001;24(3):303–312.
36. Sung V, Hiscock H, Sciberras E, Efron D. Sleep problems in children with attention-deficit/hyperactivity disorder: Prevalence and the effect on the child and family. *Arch Pediatr Adolesc Med.* 2008;162(4):336–342.
37. Lim CG, Ooi YP, Fung DS, Mahendran R, Kaur A. Sleep disturbances in Singaporean children with attention deficit hyperactivity disorder. *Ann Acad Med Singapore.* 2008;37(8):655–661.
38. Willoughby MT, Angold A, Egger HL. Parent-reported attention-deficit/hyperactivity disorder symptomatology and sleep problems in a preschool-age pediatric clinic sample. *J Am Acad Child Adolesc Psychiatry.* 2008;47(9):1086–1094.
39. Gruber R, Sadeh A, Raviv A. Instability of sleep patterns in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry.* 2000;39(4):495–501.
40. Gruber R, Sadeh A. Sleep and neurobehavioral functioning in boys with attention-deficit/hyperactivity disorder and no reported breathing problems. *Sleep.* 2004;27(2):267–273.
41. Dagan Y, Zeevi-Luria S, Sever Y, et al. Sleep quality in children with attention deficit hyperactivity disorder: An actigraphic study. *Psychiatry Clin Neurosci.* 1997;51(6):383–386.
42. Hvolby A, Jorgensen J, Bilenberg N. Actigraphic and parental reports of sleep difficulties in children with attention-deficit/hyperactivity disorder. *Arch Pediatr Adolesc Med.* 2008;162(4):323–329.
43. Wiggs L, Montgomery P, Stores G. Actigraphic and parent reports of sleep patterns and sleep disorders in children with subtypes of attention-deficit hyperactivity disorder. *Sleep.* 2005;28(11):1437–1445.
44. Owens J, Sangal RB, Sutton VK, Bakken R, Allen AJ, Kelsey D. Subjective and objective measures of sleep in children with attention-deficit/hyperactivity disorder. *Sleep Med.* 2009;10(4):446–456.
45. Paavonen EJ, Raikonen K, Lahti J, et al. Short sleep duration and behavioral symptoms of attention-deficit/hyperactivity disorder in healthy 7- to 8-year-old children. *Pediatrics.* 2009;123(5):e857–e864. Available from: <http://pediatrics.aappublications.org/content/123/5/e857.full.pdf+html>. Accessed July 22, 2011.
46. Gruber R, Xi T, Frenette S, Robert M, Vannasinh P, Carrier J. Sleep disturbances in prepubertal children with attention deficit hyperactivity disorder: A home polysomnography study. *Sleep.* 2009;32(3):343–350.
47. O'Brien LM, Ivanenko A, Crabtree VM, et al. Sleep disturbances in children with attention deficit hyperactivity disorder. *Pediatr Res.* 2003;54(2):237–243.
48. Picchiatti DL, Underwood DJ, Farris WA, et al. Further studies on periodic limb movement disorder and restless legs syndrome in children with attention-deficit hyperactivity disorder. *Mov Disord.* 1999;14(6):1000–1007.
49. Konofal E, Lecendreux M, Bouvard MP, Mouren-Simeoni MC. High levels of nocturnal activity in children with attention-deficit hyperactivity disorder: A video analysis. *Psychiatry Clin Neurosci.* 2001;55(2):97–103.
50. Cooper J, Tyler L, Wallace I, Burgess KR. No evidence of sleep apnea in children with attention deficit hyperactivity disorder. *Clin Pediatr (Phila).* 2004;43(7):609–614.
51. Miano S, Donfrancesco R, Bruni O, et al. NREM sleep instability is reduced in children with attention-deficit/hyperactivity disorder. *Sleep.* 2006;29(6):797–803.
52. Silvestri R, Gagliano A, Arico I, et al. Sleep disorders in children with Attention-Deficit/Hyperactivity Disorder (ADHD) recorded overnight by video-polysomnography. *Sleep Med.* 2009;10(10):1132–1138.
53. Kirov R, Kinkelbur J, Heipke S, et al. Is there a specific polysomnographic sleep pattern in children with attention deficit/hyperactivity disorder? *J Sleep Res.* 2004;13(1):87–93.
54. Picchiatti DL, Stevens HE. Early manifestations of restless legs syndrome in childhood and adolescence. *Sleep Med.* 2008;9(7):770–781.
55. Konofal E, Cortese S, Marchand M, Mouren MC, Arnulf I, Lecendreux M. Impact of restless legs syndrome and iron deficiency on attention-deficit/hyperactivity disorder in children. *Sleep Med.* 2007;8(7–8):711–715.
56. Kotagal S, Silber MH. Childhood-onset restless legs syndrome. *Ann Neurol.* 2004;56(6):803–807.
57. Rajaram SS, Walters AS, England SJ, Mehta D, Nizam F. Some children with growing pains may actually have restless legs syndrome. *Sleep.* 2004;27(4):767–773.
58. Chervin RD, Archbold KH, Dillon JE, et al. Associations between symptoms of inattention, hyperactivity, restless legs, and periodic leg movements. *Sleep.* 2002;25(2):213–218.
59. Chervin RD, Dillon JE, Bassetti C, Ganoczy DA, Pituch KJ. Symptoms of sleep disorders, inattention, and hyperactivity in children. *Sleep.* 1997;20(12):1185–1192.
60. Cortese S, Konofal E, Lecendreux M, et al. Restless legs syndrome and attention-deficit/hyperactivity disorder: A review of the literature. *Sleep.* 2005;28(8):1007–1013.
61. Allen RP, Picchiatti D, Hening WA, Trenkwalder C, Walters AS, Montplaisi J. Restless legs syndrome: Diagnostic criteria, special considerations, and epidemiology. A report from the restless legs syndrome diagnosis and epidemiology workshop at the National Institutes of Health. *Sleep Med.* 2003;4(2):101–109.

62. Kaplan BJ, McNicol J, Conte RA, Moghadam HK. Sleep disturbance in preschool-aged hyperactive and nonhyperactive children. *Pediatrics*. 1987;80(6):839–844.
63. Ring A, Stein D, Barak Y, et al. Sleep disturbances in children with attention-deficit/hyperactivity disorder: A comparative study with healthy siblings. *J Learn Disabil*. 1998;31(6):572–578.
64. Picchietti DL, England SJ, Walters AS, Willis K, Verrico T. Periodic limb movement disorder and restless legs syndrome in children with attention-deficit hyperactivity disorder. *J Child Neurol*. 1998;13(12):588–594.
65. Walters AS, Mandelbaum DE, Lewin DS, Kugler S, England SJ, Miller M. Dopaminergic therapy in children with restless legs/periodic limb movements in sleep and ADHD. Dopaminergic Therapy Study Group. *Pediatr Neurol*. 2000;22(3):182–186.
66. Konofal E, Arnulf I, Lecendreux M, Mouren MC. Ropinirole in a child with attention-deficit hyperactivity disorder and restless legs syndrome. *Pediatr Neurol*. 2005;32(5):350–351.
67. England SJ, Picchietti DL, Couvadelli BV, et al. L-dopa improves restless legs syndrome and periodic limb movements in sleep but not attention-deficit-hyperactivity disorder in a double-blind trial in children. *Sleep Med*. 2011;12(5):471–477.
68. Konofal E, Lecendreux M, Arnulf I, Mouren MC. Iron deficiency in children with attention-deficit/hyperactivity disorder. *Arch Pediatr Adolesc Med*. 2004;158(12):1113–1115.
69. Kryger MH, Otake K, Foerster J. Low body stores of iron and restless legs syndrome: A correctable cause of insomnia in adolescents and teenagers. *Sleep Med*. 2002;3(2):127–132.
70. Earley CJ, Allen RP, Beard JL, Connor JR. Insight into the pathophysiology of restless legs syndrome. *J Neurosci Res*. 2000;62(5):623–628.
71. Sever Y, Ashkenazi A, Tyano S, Weizman A. Iron treatment in children with attention deficit hyperactivity disorder. A preliminary report. *Neuropsychobiology*. 1997;35(4):178–180.
72. Konofal E, Lecendreux M, Deron J, et al. Effects of iron supplementation on attention deficit hyperactivity disorder in children. *Pediatr Neurol*. 2008;38(1):20–26.
73. Chervin RD, Hedger KM. Clinical prediction of periodic leg movements during sleep in children. *Sleep Med*. 2001;2(6):501–510.
74. Chervin RD, Archbold KH. Hyperactivity and polysomnographic findings in children evaluated for sleep-disordered breathing. *Sleep*. 2001;24(3):313–320.
75. Crabtree VM, Ivanenko A, O'Brien LM, Gozal D. Periodic limb movement disorder of sleep in children. *J Sleep Res*. 2003;12(1):73–81.
76. Kirk VG, Bohn S. Periodic limb movements in children: Prevalence in a referred population. *Sleep*. 2004;27(2):313–315.
77. Huang YS, Chen NH, Li HY, Wu YY, Chao CC, Guilleminault C. Sleep disorders in Taiwanese children with attention deficit/hyperactivity disorder. *J Sleep Res*. 2004;13(3):269–277.
78. Earley CJ, Connor JR, Beard JL, Malecki EA, Epstein DK, Allen RP. Abnormalities in CSF concentrations of ferritin and transferrin in restless legs syndrome. *Neurology*. 2000;54(8):1698–1700.
79. Chervin RD, Archbold KH, Dillon JE, et al. Inattention, hyperactivity, and symptoms of sleep-disordered breathing. *Pediatrics*. 2002;109(3):449–456.
80. Gottlieb DJ, Vezina RM, Chase C, et al. Symptoms of sleep-disordered breathing in 5-year-old children are associated with sleepiness and problem behaviors. *Pediatrics*. 2003;112(4):870–877.
81. Chervin RD, Ruzicka DL, Archbold KH, Dillon JE. Snoring predicts hyperactivity four years later. *Sleep*. 2005;28(7):885–890.
82. Huang YS, Guilleminault C, Li HY, Yang CM, Wu YY, Chen NH. Attention-deficit/hyperactivity disorder with obstructive sleep apnea: A treatment outcome study. *Sleep Med*. 2007;8(1):18–30.
83. Dillon JE, Blunden S, Ruzicka DL, et al. DSM-IV diagnoses and obstructive sleep apnea in children before and 1 year after adenotonsillectomy. *J Am Acad Child Adolesc Psychiatry*. 2007;46(11):1425–1436.
84. Wei JL, Bond J, Mayo MS, Smith HJ, Reese M, Weatherly RA. Improved behavior and sleep after adenotonsillectomy in children with sleep-disordered breathing: Long-term follow-up. *Arch Otolaryngol Head Neck Surg*. 2009;135(7):642–646.

Psychology Research and Behavior Management

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered include: Neuroscience, memory & decision making; Behavior

modification & management; Clinical applications; Business & sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/psychology-research-and-behavior-management-journal>

Dovepress