Poor reproductive health among a group of socially damaged Middle Eastern women: a cross-sectional study

Gohar Mohammadi¹
Sedigheh Amiraliakbari²
Ali Ramezankhani³
Hamid Alavi Majd⁴

¹The International Branch of Shahid Beheshti University of Medical Sciences, ²School of Nursing and Midwifery, ³School of Health, ⁴Department of Biostatistics, Faculty of Paramedical Science, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Background: Despite the scope of violence against women and its importance for reproductive health, few data are available on the reproductive health issues among women having experienced violence.

Methods: This study described the reproductive disorders complicating social harm among 98 socially damaged women seeking care from drop-in centers who were of Persian ancestry, able to communicate and comprehend the contents of the questionnaire, and had history of domestic violence. The questionnaire had five dimensions: demographics, reproductive health, sexual performance, sexual behavior, and violence. Reproductive health included data on gestation, unplanned pregnancy, abortion, contraception, and cervical cancer screening. Data on sexual performance was acquired via the Persian version of sexual function scale, which has been demonstrated to have acceptable external validity in Iranian population. For sexual function, data was gathered on age at first intercourse and whether a participant had ever engaged in an oral or anal sexual activity.

Results: Mean age of participants was 33.4 years. Forty-seven percent of participants were married, 34.8% were divorced, 9.8% were widowed, and 8.7% were single. Mean age at first marriage was 16.4 (4.3) years and mean age at first sexual relationship was 16 (3.9) years. Illiteracy was observed among 18.5% of participants. Elementary education was reported by 22.8%, while only 3.3% of participants reported academic studies. Fifty-five percent were unemployed and 44.6% reported to be working at the time of the study. It was observed that 72.8% of participants were inflicted physically, as well as emotionally and sexually. The violence was reported to be exerted by husband (42.6%), parents (38.4%), or both (19.0%). Among 39 participants who ran away from home, 38 participants reported to be inflicted by violence. Unwanted pregnancy was reported by 64.6% of the participants. Abortion was reported in 50.0% of participants. Contraception was completely ignored in 44.6% of participants. Among eligible women, 53.3% never participated in cervical cancer screening examination. Mean sexual performance scale score was 21.9 (5.5) and 75 (83.3%) participants scored less than 28.

Conclusion: A high prevalence of poor reproductive health was documented among a group of Middle Eastern socially damaged women.

Keywords: sexual behavior, domestic violence, pregnancy, drop-in center, abortion, contraception, cervical cancer screening

Introduction

Violence is a major obstacle to development. Violence against women in particular hinders progress in achieving development targets. Despite the growing recognition of violence against women as a public health and human rights concern, and of the obstacle it poses for development, this type of violence continues to have an unjustifi-

Correspondence: Sedigheh Amiraliakbari
Niyayesh Complex, Niyayesh Cross-
Section, Vali Asr St, Tehran, Iran
Tel +98 21 8865 5366
Fax +98 21 2272 9238
Email asa_akbari@yahoo.com

ably low priority on the international development agenda and in planning, programming, and budgeting.^{1,2} Deviant women are at increased risk of violence. In Islamic state, a “deviant woman,” that is, one who engages in an illegitimate sexual relationship, holds a precarious position. On the other hand, she is pitied as a victim of social ills; on the other, whether she turns to selling sex because of dysfunctional family life, deception, or economic needs, she leads a “pathological” life and must be cured. She is at once “socially damaged” (*äseeb-dideh-ye ejtema’ee*) and “socially deviant” (*monhæref-e ejtema’ee*). In the past few years, World Health Organization, the American Medical Association, International Federation of Obstetricians and Gynecologists, Royal College of Nursing, and other professional medical organizations have made statements about the public health importance of violence against women.³ Islamic republic of Iran policies refer to two groups of “socially deviant” women: (1) acutely at social-harm risk (*dær mæ’ræz-e äseeb-e ejtema’ee-e häd*) are the highly at-risk who run away from their families, have no guardian or visible means of support, or the savvy to manage their lives on their own, and (2) socially damaged women (special) (*zænän-e äseebdide-ye ejtema’ee-vijheh*) includes those who engage in prostitution or “women” who do not adhere to moral and social values and engage in illegitimate sex, though accrue no income in this way.⁴ In many ways, violence against women is a very well-studied, and certainly much discussed, problem. Its legal implications have been extensively analyzed, in particular how to deal with, and prosecute perpetrators.⁵ However, there are certain issues which have been almost entirely overlooked but which are vital for a fuller understanding of the lives of women who have been harassed. The most obvious ones surround matters of reproductive health and pregnancy, which have been almost entirely neglected. This study examined reproductive health among socially harmed women with a history of domestic violence.

Methods and participants

Study design and sample

This was a cross-sectional study designed to examine the scale and scope of reproductive disorders complicating social harm among socially damaged women. The interviews were conducted between May 2010 and August 2010 in Tehran, Iran. All women ($n = 98$) who sought care from socially damaged women rehabilitation centers (also known as drop-in centers or harm reduction centers) were invited to participate in a face-to-face interview and were included in the current analyses ($n = 92$) if they were of Persian ancestry,

able to communicate and comprehend the contents of the questionnaire, and had history of domestic violence. The questionnaire had five dimensions: demographics, reproductive health, sexual performance, sexual behavior, and violence. Reproductive health included data on gestation, unplanned pregnancy, abortion, contraception, and cervical cancer screening. Data on sexual performance was acquired via the Persian version of sexual function scale, which has been demonstrated to have acceptable external validity in Iranian population.^{6–8} For sexual function, data was gathered on age at first intercourse and whether a participant had ever engaged in an oral or anal sexual activity.

Definitions of terms

Violence was defined based on the three main components namely physical, emotional, and sexual violence.⁹ Domestic violence was defined as a pattern of abusive behaviors by one or more family member; sexual disorder was defined in participants whose sexual function scale score was smaller than 28.⁸

Statistics

Data are reported as either mean (standard deviation) or frequency (%) for continuously and categorically distributed variables, respectively. The statistical significance level was set at a two-tailed type I error of 0.05. Findings among socially damaged women were compared to those obtained from general population.¹⁰ Confidence intervals for differences were derived by implementing 1000 bias-corrected bootstrap resampling method.¹¹ The hypothesis for binomial random variables was also tested. The null hypothesis was that the probability of a success in a trial (in this case reproductive health characteristics) is $\#p$.¹² Here, $\#p$ were corresponding frequencies in the general population as reported by Safarinejad.¹⁰ All statistical analyses were performed using Stata version 11 (StataCorp LP, College Station, TX).

Results

Mean (standard deviation) age of participants was 33.4 (9.6) years. Forty-seven percent (46.7) of participants were married, 34.8% were divorced, 9.8% were widowed, and 8.7% were single. Mean age at first marriage was 16.4 (4.3) years and mean age at first sexual relationship was 16 (3.9) years. Illiteracy was observed among 18.5% of participants. Elementary education was reported by 22.8%, while only 3.3% of participants reported academic studies. Fifty-five percent (55.4%) were unemployed and 44.6% reported to be working at the time of the study.

Among socially damaged women seeking care from drop-in centers, 72.8% reported to be inflicted physically, as well as emotionally and sexually. Experiences of two or one type of violence were reported by 20.7% and 6.5% of participants, respectively. The violence was reported to be exerted by husband (42.6%), parents (38.4%), or both (19.0%). Among 39 participants who ran away from home, 38 participants reported to be inflicted by violence.

Pregnancy and abortion were reported in 89.1% and 50.0% of participants, respectively. Unwanted pregnancy was reported by 64.6% of the participants. Contraception was completely ignored in 44.6% of participants. The most commonly used contraception method was condom (37.0%), followed by tubectomy (8.7%), injecting progesterone (2.2%), withdrawal (4.3%), intrauterine devices (2.2%), and oral contraception pills (1.1%). Among eligible women, 53.3% never participated in cervical cancer screening examination.

Ninety out of 92 participants reported to have sex during the last 30 days leading up to the interview, with 43.3% engaging in anal and 35.6% engaging in oral sexual activity. Mean sexual performance scale score was 21.9 (5.5) and 75 (83.3%) participants scored less than 28. Low birth weight was reported by 34 (45.9%), irregular menstrual bleeding by 58.6%, and menopause by 22 (23.9%) participants.

Table 1 demonstrates the different aspects of reproductive health among socially damaged women as compared to the general population. The reproductive health of socially damaged women was statistically significantly poorer than those of women from general population. The single exception observation was the prevalence of injectable contraceptive usage.

Discussion

In this study a high prevalence of violence among socially damaged women seeking care from socially damaged women rehabilitation centers was documented. A high prevalence of sexual dysfunction was observed. Socially damaged women were observed to have poor reproductive health. Reproductive health among women from general population was previously demonstrated with respect to unwanted pregnancy, abortion, contraception, giving birth to a low birth weight newborn, and sexual dysfunction.¹⁰ In almost all areas, socially damaged women were observed to have less favorable health.

The United Nations defines violence against women as any act of gender-based violence that results in, or is likely to result in, physical, sexual or mental harm, or suffering to women, including threats of such acts, coercion, or arbitrary deprivation of liberty, whether occurring in public or in private life.¹³ There are many forms of violence against women, including sexual, physical, or emotional abuse by an intimate partner, physical or sexual abuse by family members or others, sexual harassment and abuse by authority figures (such as teachers, police officers, or employers), trafficking for forced labor or sex, and such traditional practices as forced or child marriages, dowry-related violence, and honor killings where women are murdered in the name of family honor. Systematic sexual abuse in conflict situations is another form of violence against women.¹³ In the current study, socially damaged women reported to frequently suffer all three main aspects of domestic violence namely physical, emotional, and sexual.

Women's reproductive and sexual health clearly is affected by gender-based violence. It has been shown that

Table 1 Comparison of reproductive health characteristics of the socially damaged women with those of general population

Characteristics	Socially damaged women	General population	Difference (95% confidence intervals for difference)	P value
Unwanted pregnancy	65.0	18.6	46.4 (35.0 to 57.0)	<0.001
Abortion	50.0	3.5	46.5 (35.0 to 58.0)	<0.001
Contraception	55.4	78.9	23.5 (27.0 to 50.0)	<0.001
Condom	37.0	9.3	27.7 (-13.0 to -34.0)	0.001
Tubectomy	8.7	17.4	8.7 (-14.6 to -2.8)	0.011
Injectable progestin	2.2	2.6	-0.4 (-3.5 to 2.6)	<0.781
Interrupted coitus	4.3	19.2	-14.9 (-19.1 to -10.6)	0.001
Intrauterine devices	2.2	8.1	-5.9 (-7.0 to -2.7)	0.001
Oral contraception	1.1	19.3	-18.2 (-20.4 to -16.1)	<0.001
Low birth weight	45.9	7.2	38.7 (2.0 to 4.0)	0.001
Sexual dysfunction	83.3	31.5	50.0 (41.3 to 57.6)	0.001

Notes: Data are presented as percentages. Confidence intervals for differences were derived by implementing 1000 bias-corrected bootstrap resampling method. *P* values denote exact hypothesis tests for binomial random variables. The null hypothesis is that the probability of a success in a trial (in this case frequency of each reproductive health characteristics) is $\#p$.¹² Here, $\#p$ are corresponding frequencies in the general population as reported by Safarinejad.¹⁰

women who experienced intimate partner abuse were three times more likely to have a gynecological problem than non-abused women.¹⁴ Early childbearing, often a result of early and forced marriage, can result in a range of health problems, including effects of unsafe abortion. Abuse limits women's sexual and reproductive autonomy.^{15–17} Women who have been sexually abused are much more likely than nonabused women to use family planning clandestinely, to have had their partner stop them from using family planning, and to have a partner refuse to use a condom to prevent disease.¹⁸ In line with previous reports, a low rate of contraception among socially damaged women was observed. Although, some investigators have proposed awareness and availability of contraceptive methods as culprits; factors that might have contributed to this problem remained to be clarified.¹⁹ Condoms were observed to be the most frequently used contraception method. The same finding has been reported previously.²⁰ Although the current study, allowing for its cross-sectional nature, could not provide any insight into the reason why condoms are the most frequently used contraception method, it is hypothesized that having multiple partners, fear from sexually transmitted diseases, and availability of condoms might have potentially rendered it such a priority among socially damaged women's preferences.

Unwanted pregnancy has been previously reported to be commonly observed among socially damaged women. Violence has been shown to double the risk of unplanned pregnancy.^{21–23} Survivors of abuse have been reported to be more likely to practice high-risk sexual behaviors, experience unintended pregnancies, and suffer from sexual dysfunction than nonabused women.^{24–26} A high prevalence of sexual dysfunction among socially damaged women was documented.

Induced abortion has been shown to occur among socially damaged women with higher frequency than among general population.²⁷ Causal effect of violence on incidence of induced abortion has not been documented though. Bagherzadeh et al demonstrated that socially damaged women are more likely to give birth to a newborn with low birth weight.²⁸ Some investigators have shown the physical violence to increase the risk of sexual organ damage, unplanned pregnancy, unsafe abortion, dysmenorrhea, sexual function disorders, urinary tract infections, infertility, substance abuse, sexually transmitted disease, acquired immunodeficiency syndrome, and having multiple partners.^{29–35}

As compared to other populations, Persian socially damaged women were less likely to participate in cervical cancer screening programs, ie, Pap cervical smear test.^{36,37}

This could be alarming, taking into account the link proposed to exist between multiple-partner sexual activity and cervical cancer,³⁸ in the light of the high prevalence of multiple-partner sexual relationships observed in this subpopulation. Brady et al demonstrated that violence increases the risk of having multiple partners. They observed that associations between violence involvement and other forms of health risk behavior are bidirectional. That is, adolescents who engage in sexual behavior with multiple partners are also at risk for later violence involvement.^{39,40}

Conclusion

A high prevalence of poor reproductive health was documented among a group of Middle Eastern socially damaged women. The pervasiveness of violence and its association with reproductive health underscores that violence in general is an important determinant for reproductive health risks.

Acknowledgment

The authors appreciate and would like to thank the participants in this study for revisiting their difficult experiences.

Authors' contributions

GM prepared and analyzed the data, interpreted the results, and drafted the article. SA, AR, and HAM critically revised the manuscript.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Department of Gender, Women and Health Family and Community Health, World Health Organization. *Addressing Violence Against Women and Achieving the Millennium Development Goals*. Geneva: World Health Organization; 2005.
2. Garcia-Moreno C, Heise L, Jansen HA, Ellsberg M, Watts C. Violence against women. *Science*. 2005;310(5752):1282–1283.
3. Krug EG, Dahlberg LL, Mercy JA, Zwi AB, Lozano R, editors. *World Report on Violence and Health*. Geneva: World Health Organization; 2002.
4. Shahidian H. *Women in Iran: Gender Politics in the Islamic Republic*. Westport, CT: Greenwood Press; 2002.
5. Watts C, Zimmerman C. Violence against women: global scope and magnitude. *Lancet*. 2002;359(9313):1232–1237.
6. Wiegel M, Meston C, Rosen R. The female sexual function index (FSFI): cross-validation and development of clinical cutoff scores. *J Sex Marital Ther*. 2005;31(1):1–20.
7. Khademi A, Alleyassin A, Amini M, Ghaemi M. Evaluation of sexual dysfunction prevalence in infertile couples. *J Sex Med*. 2008;5(6):1402–1410.
8. Mohammadi K, Heydari M, Faghihzadeh S. The female sexual function index (FSFI): validation of the Iranian version. *Payesh*. 2008;7(3):269–278.

9. Dolatian M, Hesami K, Shams J, Alavi Majd H. Evaluation of the relationship between domestic violence in pregnancy and postnatal depression. *Scientific Journal of Kurdistan University of Medical Sciences*. 2008;13(2):57–68.
10. Safarinejad MR. Female sexual dysfunction in a population-based study in Iran: prevalence and associated risk factors. *Int J Impot Res*. 2006; 18(4):382–395.
11. Davison AC, Hinkley DV. *Bootstrap Methods and their Application*. Cambridge: Cambridge University Press; 2006.
12. Hoel PG, Port SC, Stone CJ. *Introduction to Stochastic Processes*. New York: John Wiley and Sons, Inc; 1984.
13. World Health Organization (WHO). Violence against women: intimate partner and sexual violence against women. Fact sheet no 239. Geneva: WHO; August 17, 2004 [updated Sep 2011]. Available from: <http://www.who.int/mediacentre/factsheets/fs239/en/>. Accessed June 11, 2011.
14. Campbell JC. Health consequences of intimate partner violence. *Lancet*. 2002;359(9314):1331–1336.
15. Kalichman SC, Williams EA, Cherry C, Belcher L, Nachimson D. Sexual coercion, domestic violence, and negotiating condom use among low-income African American women. *J Womens Health*. 1998; 7(3):371–378.
16. Tjaden P, Thoennes N. *Extent, Nature, and Consequences of Intimate Partner Violence: Findings from the National Violence Against Women Survey*. Washington, DC: National Institute of Justice; 2000.
17. O'Donnell L, Agronick G, Duran R, Myint-U A, Stueve A. Intimate partner violence among economically disadvantaged young adult women: associations with adolescent risk taking and pregnancy experiences. *Perspect Sex Reprod Health*. 2009;41(2):84–91.
18. Ellsberg M, Jansen HA, Heise L, et al. Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: an observational study. *Lancet*. 2008;371(9619):1165–1172.
19. Gelberg L, Lu MC, Leake BD, Andersen RM, Morgenstern H, Nyamathi AM. Homeless women: who is really at risk for unintended pregnancy? *Matern Child Health J*. 2008;12(1):52–60.
20. Patterson TL, Mausbach B, Lozada R, et al. Efficacy of a brief behavioral intervention to promote condom use among female sex workers in Tijuana and Ciudad Juarez, Mexico. *Am J Public Health*. 2008;98(11):2051–2057.
21. Pallitto CC, O'Campo P. The relationship between intimate partner violence and unintended pregnancy: analysis of a national sample from Colombia. *Int Fam Plan Perspect*. 2004;30(4):165–173.
22. Pallitto CC, O'Campo P. Community level effects of gender inequality on intimate partner violence and unintended pregnancy in Colombia: testing the feminist perspective. *Soc Sci Med*. 2005;60(10):2205–2216.
23. Pallitto CC, Campbell JC, O'Campo P. Is intimate partner violence associated with unintended pregnancy? A review of the literature. *Trauma Violence Abuse*. 2005;6(3):217–235.
24. Bensley LS, Van Eenwyk J, Simmons KW. Self-reported childhood sexual and physical abuse and adult HIV-risk behaviors and heavy drinking. *Am J Prev Med*. 2000;18(2):151–158.
25. Rotheram-Borus-MJ, Mahler KA, Koopman C, Langabeer K. Sexual abuse history and associated multiple risk behavior in adolescent runaways. *Am J Orthopsychiatry*. 1996;66(3):390–400.
26. Raj A, Silverman JG, Amaro H. The relationship between sexual abuse and sexual risk among high school students: findings from the 1997 Massachusetts Youth Risk Behavior Survey. *Matern Child Health J*. 2000;4(2):125–134.
27. Fergusson DM, Boden JM, Horwood LJ. Abortion among young women and subsequent life outcomes. *Perspect Sexual Reprod Health*. 2007; 39(1):6–12.
28. Bagherzadeh R, Keshavarz T, Sharif F, Dehbashi S, Tabatabaei HR. Relationship between domestic violence during pregnancy and complications of pregnancy, type of delivery and birth weight on delivered women in hospital affiliated to Shiraz university of Medical Sciences. *Ofogh-e-Danesh Journal*. 2008;13(4):51–58.
29. Straus MA, Gelles RJ, Smith C. *Physical Violence in American Families: Risk Factors and Adaptations to Violence in 8,145 Families*. New Brunswick, NJ: Transaction Publishers; 1995.
30. Kilpatrick DG, Ruggiero KJ, Acierio R, Saunders BE, Resnick HS, Best CL. Violence and risk of PTSD, major depression, substance abuse/dependence, and comorbidity: results from the National Survey of Adolescents. *J Consult Clin Psychol*. 2003;71(4):692–700.
31. Gazmararian JA, Adams MM, Saltzman LE, et al. The relationship between pregnancy intendedness and physical violence in mothers of newborns. The PRAMS Working Group. *Obstet Gynecol*. 1995; 85(6):1031–1038.
32. Dunkle KL, Jewkes RK, Brown HC, Gray GE, McIntyre JA, Harlow SD. Gender-based violence, relationship power, and risk of HIV infection in women attending antenatal clinics in South Africa. *Lancet*. 2004; 363(9419):1415–1421.
33. Silverman JG, Raj A, Mucci LA, Hathaway JE. Dating violence against adolescent girls and associated substance use, unhealthy weight control, sexual risk behavior, pregnancy, and suicidality. *JAMA*. 2001;286(5):572–579.
34. Golding JM. Intimate partner violence as a risk factor for mental disorders: a meta-analysis. *J Fam Violence*. 1999;14(2):99–132.
35. Hotaling GT, Sugarman DB. An analysis of risk markers in husband to wife violence: the current state of knowledge. *Violence Vict*. 1986;1(2): 101–124.
36. Cheek J, Fuller J, Gilchrist S, Maddock A, Ballantyne A. Vietnamese women and Pap smears: issues in promotion. *Aust NZ J Public Health*. 1999;23(1):72–76.
37. Yu C, Rymer J. Women's attitudes to and awareness of smear testing and cervical cancer. *Br J Fam Plann*. 1998;23(4):127–133.
38. Brinton LA. Epidemiology of cervical cancer – overview. *IARC Sci Publ*. 1992;119:3–23.
39. Brady SS, Donenberg GR. Mechanisms linking violence exposure to health risk behavior in adolescence: motivation to cope and sensation seeking. *J Am Acad Child Adolesc Psychiatry*. 2006;45(6):673–680.
40. Brady SS, Tschann JM, Pasch LA, Flores E, Ozer EJ. Violence involvement, substance use, and sexual activity among Mexican-American and European-American adolescents. *J Adolesc Health*. 2008;43(3):285–295.

International Journal of Women's Health

Publish your work in this journal

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. Subject areas include: Chronic conditions (migraine headaches, arthritis, osteoporosis);

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-womens-health-journal>

Endocrine and autoimmune syndromes; Sexual and reproductive health; Psychological and psychosocial conditions. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.