Identification of *Hepatocystis* species in a macaque monkey in northern Myanmar

Qiaocheng Chang^{1,*}Xiaodong Sun^{2,*}Jian Wang^{2,*}Jigang Yin¹Junpeng Song¹Shuai Peng¹Huijun Lu¹Hongning Zhou²Ning Jiang¹Qijun Chen^{1,3}

¹Key Laboratory of Zoonosis, Jilin University, Changchun; ²Institute for Parasitic Disease Control of Yunnan Province, Puer City, Yunnan; ³Institute of Pathogen Biology, Chinese Academy of Medical Sciences, Beijing, China

*These authors contributed equally to this work

Correspondence: Qijun Chen
Institute of Pathogen Biology,
Chinese Academy of Medical Sciences,
Dong Dan San Tiao 9,
Beijing 100730, China
Tel +86 10 6782 8512
Fax +86 10 6783 7321
Email qijun.chen@ipbcams.ac.cn

Ning Jiang
Key Laboratory of Zoonosis,
Jilin University,
Changchun 130062, China
Tel +86 431 8783 6721
Fax +86 431 8783 6701
Email: jiangning@jlu.edu.cn

Background: Long-tailed and pig-tailed macaque monkeys are natural hosts of *Plasmodium knowlesi*, which has been identified as a fifth malaria parasite infecting humans. In this study, we investigated possible infection by this *Plasmodium* parasite in macaque monkeys using a combination of polymerase chain reaction amplification and sequencing.

Methods: Forty-five blood samples were obtained in 2010 from macaques in northern Myanmar near Yunnan Province of China and investigated for possible infection with *Plasmodium* species using a nested polymerase chain reaction method for amplification of 18S SSU rRNA genes.

Results: Positive amplification was obtained from one monkey, and both sequence and phylogenetic analysis indicated that the parasite was of the *Hepatocystis* species lineage.

Conclusion: The results suggest that a combination of polymerase chain reaction amplification and sequence identification would be necessary for detection of *Plasmodium knowlesi* infection in both humans and its natural hosts.

Keywords: *Plasmodium knowlesi*, monkey, parasite, malaria

Background

In recent years, a fifth *Plasmodium* species, ie, *P. knowlesi*, has been identified as an invasive pathogen in humans,¹ and cases of *P. knowlesi* infection have been concentrated in southeast Asian countries.^{2,3} We recently found that the prevalence of *P. knowlesi* in the border area between China and Myanmar was more than 20% in malaria-infected patients, and the parasite was frequently found in coinfection with either *P. falciparum* or *P. vivax*.⁴

Currently, species identification of *P. knowlesi* and differentiation from other malarial parasites has relied mainly on polymerase chain reaction (PCR) amplification of 18S SSU rRNA genes.^{1,5} Studies have indicated that it is difficult to differentiate *P. knowlesi* from other parasites by examination of blood smears.⁶ Further, apart from *P. knowlesi*, *Hepatocystis* species have been reported to be common pathogens in macaque monkeys.⁷⁻⁹ The genus of blood-parasitizing hemosporines (family *Plasmodiidae*) features gametocytes in red cells and cyst-like exoerythrocytic schizonts in the liver parenchyma. These are parasitic in old world primates, bats, and squirrels, but not in domestic animals.⁷ *Hepatocystis* species are transmitted by biting *Culicoides* species, such as *C. adersi*.^{8,9} Escalante et al first identified a species of *Hepatocystis* in a phylogeny of malarial parasites and found it clustered within the primate *Plasmodium*.¹⁰ Perkins and Schall suggested that *Hepatocystis* shared a common ancestor closer to primate malaria parasites than

to avian parasites.¹¹ Infected animals are asymptomatic and do not experience hemolysis.¹² It was reported that *Hepatocystis* parasites do not produce the cyclical fever spikes typical of malaria in humans, but do cause anemia and visible merocyst formation, followed by scarring in the liver.^{7,13} Thus, it is clinically important to differentiate *Hepatocystis* parasites from other malaria parasites in both natural hosts and humans, for clinical diagnosis as well as for epidemiological clarification.

Systematic molecular studies provide an independent source of evidence for elucidating the origin of parasites. Studies found that partial gene conversion occurs among nonhomologous copies of the 18S SSU rRNA genes that are expressed during different parasite stages, which can generate misleading phylogenetic results.¹⁴ To overcome this limitation, we studied the cytochrome *b* gene independent of the mitochondrial genome, which is evolutionarily conserved.¹⁵ In this study, we investigated possible infection of *Plasmodium* parasite in macaque monkeys using a combination of PCR amplification and sequencing.

Materials and methods

Blood samples

Forty-five blood samples from macaque monkeys were collected in the border area between China and Myanmar. A 20–50 µL drop of finger-prick blood was spotted directly onto premarked filter paper which was allowed to dry in air and was then stored individually in plastic bags. A blood smear was made for each monkey and stained using Geimsa solution followed by microscopy. The study was performed with permission from the ethics committee of the Institute for Parasitic Disease Control of Yunnan province and the local administration authority in Myanmar.

DNA extraction

DNA templates were prepared from whole blood spots on filter paper according to a previously reported method.¹⁶ Each filter paper punch was soaked in 125 µL of methanol. After incubation at room temperature for 15 minutes, the methanol was removed and the samples were dried before adding 65 µL of distilled water. The punches were mashed using a new pipette tip for each punch and heated at 97°C for 15 minutes to elute the DNA.

PCR amplification of 18S SSU rRNA gene

The DNA samples were initially analyzed using specific nested PCR assays as described elsewhere.⁵ The first PCR

amplification (nest 1) using *Plasmodium* genus-specific primers for each sample was carried out in a 50 µL reaction mixture containing 2.5 mM of each primer (rPLU1 and rPLU5), 1 × PCR buffer, 2.5 mM of each dNTP, 1.25 U of *Taq* DNA polymerase (Promega, Madison, WI) and 0.5 µg of DNA template. Two microliters of nest 1 amplification was used as the template DNA in the second PCR amplification (nest 2) using the self-designed primers (CHN18-R 5'-TAA GGA TAA CTA CGG AAA AGC TGT-3' and CHN18-F 5'-AAG ATT ACG ACG GTA TCT GA-3'). The PCR conditions were 4 minutes at 94°C, 30 seconds at 94°C, 30 seconds at 58°C, one minute at 72°C for 36 cycles, and a final 10-minute extension at 72°C. Amplicons were examined by 1% agarose gel electrophoresis and ethidium bromide staining.

DNA bands were removed from the gel, purified using the QIAquick gel extraction kit (Qiagen, Valencia, CA), and ligated to a T-cloning vector (Invitrogen, Carlsbad, CA) according to protocols provided by the manufacturers. The DNA sequences were determined from at least five plasmid clones and sequenced by BGI (Beijing, China).

PCR amplification of cytochrome *b* gene of *Hepatocystis*

To confirm further that the sequences amplified with the 18S SSU rRNA primers were derived from *Hepatocystis*, the cytochrome *b* fragments of *Hepatocystis* were amplified using nested PCR. The cytochrome *b* nested PCR primers were CHNb 1 (5'-GAG AAT TAT GGA GTG GAT GGT G-3'), CHNb 2 (5'-GTG GTA ATT GAC ATC CAA TCC-3'), CHNb 3 (5'-GGT GTT TCA GAT ATA TGC ATG C-3'), and CHNb 4 (5'-CAT CCA ATC CAT AAT AAA GCA TAG-3'), which were designed based on the *Hepatocystis* cytochrome *b* gene (GU930067.1). The PCR reaction was carried out in a total volume of 25 µL under the following conditions: 1 µM of each primer, ie, CHNb 1 and CHNb 2, 2.5 mM of each dNTP, 0.5 U of DNA polymerase (Promega), 0.1 µg DNA template, and 1 × reaction buffer. The PCR conditions were 4 minutes at 94°C, 30 seconds at 94°C, 30 seconds at 55°C, 90 seconds at 72°C for 36 cycles, and a final 10-minute extension at 72°C. Nested PCR was carried out using 1 µL of the PCR products and performed with CHNb 3 and CHNb 4 primers. The PCR conditions were 4 minutes at 94°C, 30 seconds at 94°C, 30 seconds at 58°C, 30 seconds at 72°C for 35 cycles, and a final seven-minute extension at 72°C. The PCR products were detected by agarose electrophoresis, and cloned and sequenced as described earlier.

Figure 1 Polymerase chain reaction amplification of 18S SSU rRNA and cytochrome *b* genes. **(A)** Nested polymerase chain reaction amplification of the 18S SSU rRNA gene using the primers rPLU3 and rPLU4 matched to all pathogenic *Plasmodium* species. Lane 1 is a positive control. Lanes 2–11 are products of amplification of 10 monkey blood samples, and lane 12 is a negative control without DNA. **(B)** The 18S SSU rRNA and the cytochrome *b* genes were amplified by nested polymerase chain reaction with specific primers to the sequences of *Hepatocystis* species. Lanes 1 and 2 were the amplicons of 18S SSU rRNA using the primers CHN18-R and CHN18-F and the cytochrome *b* genes with the primers CHNb 3 and CHNb 4, respectively.

Sequence analysis

Sequences were first blasted in the Genbank for identification. Homologous sequences were aligned using the CLUSTAL W program.¹⁷ Phylogenetic trees were reconstructed by the neighbor-joining method.^{18,19} Reliability of clustering patterns

was tested by bootstrapping,²⁰ and 1000 bootstrap samples were used.

Results

PCR amplification

In this study, 45 macaque blood samples were first investigated for possible infection of *P. knowlesi* by nested PCR amplification of 18S SSU rRNA. A positive amplification was obtained from one monkey (Figure 1A, lane 8). Further sequence identification found that the positive product might be generated from a closely related organism, ie, *Hepatocystis* species. For further clarification, primer sets were designed based on both 18S SSU rRNA and mitochondrial cytochrome *b* genes of *Hepatocystis* species. The PCR product of the secondary amplification of the 18S SSU rRNA gene was 961 bp⁴ (Figure 1B, lane 1). The size of the PCR product of the mitochondrial cytochrome *b* gene was 788 bp (Figure 1B, lane 2).

18S SSU rRNA sequences and cytochrome *b* derived from *Hepatocystis*

Sequences of the PCR products were blasted against the sequences in GenBank (<http://blast.ncbi.nlm.nih.gov/>). Phylogenetic analysis of the 18S SSU rRNA gene (GenBank

Figure 2 Phylogenetic analysis based on the 18S SSU rRNA genes of malaria and *Hepatocystis* species using the neighbour-joining method. HQ605039 was the sequence newly obtained in this study.

Figure 3 Phylogenetic analysis based on the mitochondrial cytochrome *b* genes of malaria and *Hepatocystis* species using the neighbour-joining method. HQ605040 was newly obtained in this study.

HQ605039) and those of published malaria parasites showed that the sequence was more related to *Hepatocystis* species than to *Plasmodium* species (Figure 2).

The PCR product of cytochrome *b* gene was analyzed to confirm that the 18S SSU rRNA sequence was truly of *Hepatocystis* origin. The 788 bp PCR product was sequenced and only one sequence was obtained. A blast search indicated that the sequence has a high similarity with that of *Hepatocystis* species lineage. A phylogenetic tree was constructed with the novel sequence (HQ605040) and related cytochrome *b* sequences from *Hepatocystis* species isolated from Myanmar macaques and other malaria parasites. The sequence clearly clustered with that of *Hepatocystis* species (Figure 3).

Table 1 Sequence accession numbers and origin of the mitochondrial cytochrome *b* gene identified in *Hepatocystis* species, which was used in the phylogenetic analysis of Figure 4

GenBank accession	Geographic local source	Parasite host
GU929951	Southeast Asian	Primate
EU400409	Thailand	Primate
EU400408	Thailand	Primate
GU930067	Thailand	Primate
AF069626	Ethiopia	Primate
HQ605040	Myanmar	Primate
FJ168565	Asian	Pteropus
EF587238	Malaysia	Pteropus
DQ396531	Malaysia	Pteropus
EU254526	Singapore	Rousettus
EU254527	Guinea	Rousettus
EU254528	Guinea	Rousettus

Further phylogenetic analysis of the cytochrome *b* sequences (see Table 1) showed that these sequences form three distinct clusters (Figure 4). A cluster with 98% bootstrap indicated that our sequence (HQ605040) formed a clade of primate species (GU929951, EU400409, EU400408, GU930067) from Southeast Asian and East Africa (AF069626, Figure 4). The bat *Hepatocystis* species fell into two divergent clades. The first contained three parasite sequences from *Pteropus* (FJ168565, EF587238, DQ396531) from Southeast Asia. The other species of *Rousettus* (EU254526) from the same region formed a sister group to the *Hepatocystis* species in *Rousettus* (EU254527, EU254528) from West Africa (Guinea, Figure 4).

Discussion

In this study, 45 macaque monkeys were investigated for possible infection of *Plasmodium* species using the classical PCR method by amplification of the 18S SSU rRNA genes. The blood samples were first examined on-spot by microscopy and no parasite was observed. Surprisingly, a positive amplification was obtained from one sample using the primers specific for *Plasmodium* species. The molecular size of the PCR product was as expected (Figure 1A). Since *P. knowlesi* was detected earlier in human blood samples in the same region,⁴ we suspected that the amplified sequence might be derived from *P. knowlesi*. A blast search with the sequence against the sequences in the Genbank indicated that the sequence was more related to that of *Hepatocystis* species, although it had high similarity to the 18S SSU rRNA sequences of *Plasmodium* species (data not shown).

Figure 4 Phylogeny of *Hepatocystis* species using mitochondrial cytochrome *b* genes. Numbers on branches are percentages of 1000 bootstrap supporting a given branch. Sequences from *Hepatocystis* species are listed along with their respective GenBank accession numbers. HQ605040 was newly obtained in this study.

A literature search indicated that *Hepatocystis* has indeed been found in the monkey in the region before.^{21,22}

The result of the 18S SSU rRNA sequences was further confirmed by amplification of mitochondrial cytochrome *b* gene with a nested PCR. The sequence of the amplified product clustered within that of *Hepatocystis*. Thus, the parasite identified in the macaque monkey belonged to the genus of *Hepatocystis*. Importantly, only one sequence was obtained with the second set of primers, indicating only one species existed in the samples.

Our results are reliable for three reasons. First, nested PCR methods have proved to be more specific and sensitive than conventional microscopy, and nested PCR assays are extremely valuable tools for obtaining accurate epidemiologic data.¹⁷ Second, the 18S SSU rRNA and cytochrome *b* gene from the mitochondrial genome have been regarded as reference sequences for the detection of *Hepatocystis*,^{8,22} and we applied the sequence signatures of both 18S SSU rRNA and the cytochrome *b* gene to identify the parasite. Third, phylogenetic analysis based on the two genes supported a close phylogenetic relationship of both genes with that from the *Hepatocystis* species lineage. Finally, our data suggest that both PCR and deep sequence analysis are necessary for correct identification of primate malaria parasites which may have a similar genome sequence signature to that of *Hepatocystis* species.

Conclusion

In this study, the possible prevalence of *P. knowlesi* in 45 long-tailed macaque monkeys in northern Myanmar was investigated

using *Plasmodium* genus-specific primers followed by sequencing of the PCR product. Positive amplification was obtained from one sample. Sequence comparison indicated that the sequence had high similarity to that of *Hepatocystis* species. Further confirmation with primers to both 18S SSU rRNA and mitochondrial cytochrome *b* genes suggested that the monkey was infected by *Hepatocystis* species, but not *P. knowlesi*. Thus, the results suggest that a combination of PCR amplification and sequence identification is necessary for correct detection of *P. knowlesi* infection in both humans and its natural hosts.

Acknowledgments

This study was carried out with support to QC from the National Basic Research Program of China (973 Program, 2007CB513100), the National Science and Technology Project (2008zc-10004-011), the Chinese Natural Scientific Foundation (81171592, 81130033), and the institutional grants to QC from Institute of Pathogen Biology (IPB, CAMS) and Jilin University, China.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Singh B, Sung LK, Matusop A, et al. A large focus of naturally acquired *Plasmodium knowlesi* infections in human beings. *Lancet*. 2004;363:1017–1024.
2. Baird JK. Malaria zoonoses. *Travel Med Infect Dis*. 2009;7:269–277.
3. Cox-Singh J, Singh B. Knowlesi malaria: newly emergent and of public health importance? *Trends Parasitol*. 2008;24:406–410.

4. Jiang N, Chang Q, Sun X, et al. Co-infections with *Plasmodium knowlesi* and other malaria parasites, Myanmar. *Emerg Infect Dis*. 2010;16:1476–1478.
5. Singh B, Bobogare A, Cox-Singh J, Snounou G, Abdullah MS, Rahman HA. A genus-and species-specific nested polymerase chain reaction malaria detection assay for epidemiologic studies. *Am J Trop Med Hyg*. 1999;60:687–692.
6. Lee KS, Cox-Singh J, Brooke G, Matusop A, Singh B. *Plasmodium knowlesi* from archival blood films: Further evidence that human infections are widely distributed and not newly emergent in Malaysian Borneo. *Int J Parasitol*. 2009;39:1125–1128.
7. Garnham PCC. *Malaria Parasites and Other Haemosporidia*. Oxford, UK: Blackwell Scientific; 1966.
8. Seethamchai S, Putaporntip C, Malaivijitnond S, Cui L, Jongwutiwes S. Malaria and *Hepaticystis* species in wild macaques, southern Thailand. *Am J Trop Med Hyg*. 2008;78:646–653.
9. Garnham PC, Heisch RB, Minter DM. *Culicoides adersi* Ingram and Macfie, 1923, a presumed vector of hepaticystis (=Plasmodium) kochi (Laveran, 1899). *Nature*. 1961;190:737–741.
10. Escalante AA, Freeland DE, Collins WE, Lal AA. The evolution of primate malaria parasites based on the gene encoding cytochrome b from the linear mitochondrial genome. *Proc Natl Acad Sci U S A*. 1998;95:8124–8129.
11. Perkins SL, Schall JJ. A molecular phylogeny of malarial parasites recovered from cytochrome b gene sequences. *J Parasitol*. 2002;88:972–978.
12. Zeiss CJ, Shomer N. *Hepaticystosis* in a baboon (*Papio anubis*). *Contemp Top Lab Anim Sci*. 2001;40:41–42.
13. Tung J, Primus A, Bouley AJ, Severson TF, Alberts SC, Wray GA. Evolution of a malaria resistance gene in wild primates. *Nature*. 2009;460:388–391.
14. Escalante AA, Goldman IF, De Rijk P, et al. Phylogenetic study of the genus *Plasmodium* based on the secondary structure-based alignment of the small subunit ribosomal RNA. *Mol Biochem Parasitol*. 1997;90:317–321.
15. Milinkovitch MC, Meyer A, Powell JR. Phylogeny of all major groups of cetaceans based on DNA sequences from three mitochondrial genes. *Mol Biol Evol*. 1994;11:939–948.
16. Berezcky S, Martensson A, Gil JP, Farnert A. Short report: Rapid DNA extraction from archive blood spots on filter paper for genotyping of *Plasmodium falciparum*. *Am J Trop Med Hyg*. 2005;72:249–251.
17. Thompson JD, Gibson TJ, Higgins DG. Multiple sequence alignment using ClustalW and ClustalX. *Curr Protoc Bioinformatics*. 2002, Chapter 2:Unit 2.3.
18. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol*. 1987;4:406–425.
19. Nei M, Kumar S. *Molecular Evolution and Phylogenetics*. Oxford, UK: Oxford University Press; 2000.
20. Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*. 1985;39:783–791.
21. Takenaka T, Hashimoto K, Gotou H, Matsumoto S, Nishikawa T. Studies on *Hepaticystis* spp. in rhesus monkeys from Yunnan, China. *Jikken Dobutsu*. 1990;39:273–279. Japanese.
22. Putaporntip C, Jongwutiwes S, Thongaree S, Seethamchai S, Grynberg P, Hughes AL. Ecology of malaria parasites infecting Southeast Asian macaques: evidence from cytochrome b sequences. *Mol Ecol*. 2010;19:3466–3476.

Research and Reports in Tropical Medicine

Publish your work in this journal

Research and Reports in Tropical Medicine is an international, peer-reviewed, open access journal publishing original research, case reports, editorials, reviews and commentaries on all areas of tropical medicine, including: Diseases and medicine in tropical regions; Entomology; Epidemiology; Health economics issues; Infectious disease; Laboratory

Submit your manuscript here: <http://www.dovepress.com/research-and-reports-in-tropical-medicine-journal>

Dovepress

science and new technology in tropical medicine; Parasitology; Public health medicine/health care policy in tropical regions; and Microbiology. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.