Current and future options for the management of phantom-limb pain

Helena Knotkova^{1,2}
Ricardo A Cruciani¹⁻³
Volker M Tronnier⁴
Dirk Rasche⁴

¹Department of Pain Medicine and Palliative Care, Research Division, Institute for Non-invasive Brain Stimulation, Beth Israel Medical Center, New York, NY, USA;

²Department of Neurology, Albert Einstein College of Medicine, Bronx, NY, USA; ³Department of Anesthesiology, Albert Einstein College of Medicine, Bronx, NY, USA; ⁴Department of Neurosurgery, University of Lübeck, Germany

Abstract: Phantom-limb pain (PLP) belongs among difficult-to-treat chronic pain syndromes. Treatment options for PLP are to a large degree implicated by the level of understanding the mechanisms and nature of PLP. Research and clinical findings acknowledge the neuropathic nature of PLP and also suggest that both peripheral as well as central mechanisms, including neuroplastic changes in central nervous system, can contribute to PLP. Neuroimaging studies in PLP have indicated a relation between PLP and the neuroplastic changes. Further, it has been shown that the pathological neuroplastic changes could be reverted, and there is a parallel between an improvement (reversal) of the neuroplastic changes in PLP and pain relief. These findings facilitated explorations of novel neuromodulatory treatment strategies, adding to the variety of treatment approaches in PLP. Overall, available treatment options in PLP include pharmacological treatment, supportive non-pharmacological non-invasive strategies (eg, neuromodulation using transcranial magnetic stimulation, visual feedback therapy, or motor imagery; peripheral transcutaneous electrical nerve stimulation, physical therapy, reflexology, or various psychotherapeutic approaches), and invasive treatment strategies (eg, surgical destructive procedures, nerve blocks, or invasive neuromodulation using deep brain stimulation, motor cortex stimulation, or spinal cord stimulation). Venues of further development in PLP management include a technological and methodological improvement of existing treatment methods, an implementation of new techniques and products, and a development of new treatment approaches.

Keywords: phantom-limb pain (PLP), neuropathic pain, non-invasive treatment, invasive treatment, neuromodulation

Introduction

It is estimated that more than 80% of patients with partial or total loss of a limb develop chronic phantom-limb pain (PLP), ie, pain that seems to be located in the missing limb.¹ For the purposes of clinical assessment and treatment, it is important to differentiate PLP from other amputation-related phenomena, such as stump-pain, (ie, pain in the remaining part of the limb),² non-painful phantom-limb sensation, or telescoping, ie, sensation when the distal part of the phantom is gradually felt to approach the residual limb and may even be experienced within the stump.³ Risks factors for PLP⁴⁻⁸ include gender (PLP being more common in women), upper extremity amputation, presence of pre-amputation pain, residual pain in remaining limb, or time after amputation (there are reports of a two-peak period of onset, the first within a month after amputation and the second after 1 year).⁸ Further, stress, anxiety, depression, and other emotional triggers also highly likely contribute to PLP.⁷ Recommended preventive measures

Correspondence: Helena Knotkova
Institute for Non-invasive Brain Stimulation, Research Division,
Department of Pain Medicine and Palliative Care, Beth Israel Medical Center, 120 E 16th Street, 12th Floor, New York, NY, 10003, USA
Tel +1 212 844 8541
Fax +1 212 844 8541
Email hknotkov@chpnet.org

include using preemptive and postoperative perineural or epidural local anesthetics, intraoperative precautions related to protection of the peripheral nerves, and importance of early rehabilitation.^{9,10}

Treatment options for PLP are, to a large degree, implicated by the level of understanding the mechanisms and nature of PLP. In a historical perspective, PLP had been considered mostly of psychological origin, with the prevalent belief that PLP was generated “in the patient’s head.” However, the development of advanced diagnostic methods, recently including neuroimaging, has facilitated explorations of changes in peripheral and central neural networks after amputation and their putative contribution to the development of PLP.^{11–13} The findings acknowledged the neuropathic nature of PLP and also suggested that both peripheral, as well as central mechanisms, including neuroplastic changes in central nervous system, can contribute to PLP.^{14,15} In periphery, axotomized afferent neurons may develop retrograde degeneration and shrinking that primarily involves unmyelinated neurons, and local changes at the site of the remaining nerve endings may contribute to the development of enhanced sensitivity of the afferent fibers.¹⁶ Further, a sprouting of injured axon can lead to formation of neuromas in the residual limb. Neuromas show abnormal sensitivity to mechanical and chemical stimuli and can be a source of abnormal ectopic discharges.^{16,17} Besides neuromas, dorsal root ganglions have also been shown to generate the discharge of ectopic activity,¹⁸ substantially increasing the overall barrage of abnormal afferent input to the spinal cord and upper centers, such as the brainstem, thalamus, or cortex. Further, post-amputation changes at the level of the spinal cord include formation of new neural connections between the axonal sprouts from the proximal part of the amputated nerve and spinal cord neurons, which contributes to sensitization of pain-transmission neurons. The sensitization may manifest itself as mechanical hyperalgesia and an expansion of peripheral receptive fields,¹⁹ and is linked to an increase of activity on NMDA receptors mediated for example by substance P, neurokinins, and tachykinins at the dorsal horns of the spinal cord.²⁰ There also may be a reduction in the local intersegmental inhibitory mechanisms, resulting in spinal disinhibition and enhanced nociceptive input to the supraspinal centers. Interestingly, a pathological change of the afferent input (whether it is an increase or decrease) represents a significant source of neural plasticity, ie, dynamic changes in the function of neurons and neural networks in the central nervous system, including both spinal and cerebral parts of the Pain Matrix, the pain

processing network.^{18,21} Neuroplastic alterations at the level of the brain include changes in neuronal excitability, changes in the somatotopic organization in the cortical and subcortical areas (eg, in the somatosensory and motor cortices or in thalamus), and structural changes (thickening or thinning) of neuronal layers. Indeed, neuroimaging studies in PLP have shown presence of such neuroplastic changes in amputees and the relation to the presence of PLP.^{11–13,22–25} Further, it has been shown that the pathological neuroplastic changes can be reverted and that an improvement (reversal) of the neuroplastic changes in PLP patients was paralleled by pain relief.¹³ These findings facilitated explorations of novel neuromodulatory treatment strategies, such as transcranial magnetic stimulation (TMS) or motor imagery, adding to the variety of treatment approaches in PLP.

Overall, treatment options in PLP include pharmacological treatment, supportive non-pharmacological non-invasive strategies, and invasive treatments.

Pharmacological treatment

The pharmacological armamentarium is vast,^{26–49} (Table 1), but efficacy of most of the drugs utilized to treat PLP have not been determined from PLP controlled trials, but rather extrapolated from positive results obtained on a variety of other neuropathic pain syndromes (eg, painful polyneuropathy [PPN] or post-herpetic neuralgia [PHN]).

Acetaminophen and nonsteroidal anti-inflammatory drugs (NSAIDs)

A cross-sectional study found acetaminophen and NSAIDs to be the most common medications used in PLP.⁴ This widely-used over-the-counter medication has relatively little anti-inflammatory activity as compared to NSAIDs. However, acetaminophen and other NSAIDs all act by the same mechanism (inhibition of prostaglandin synthesis) and all show varying levels of analgesic, anti-inflammatory, antipyretic, and antiplatelet actions, and decrease the nociception at peripheral and central level.²⁸

Antidepressants

Antidepressants have been utilized for the management of neuropathic pain since the eighties after work by Mitchel Max showed analgesic effect by amitriptyline independent of improvement of depression. Since then many drugs have been added to the list (Table 1), with different degrees of success.^{26,29,30}

Tricyclics are on top of the list for their efficacy for the management of neuropathic pain syndromes, but the results

Table I Level of evidence^a for the drugs utilized to treat phantom-limb pain and other neuropathic-pain conditions

Drug	Neuropathic-pain conditions					
	PLP	PPN	PHN	TN	HIV	CP
Acetaminophen	CrS/P					
NSAIDs	CrS/P					
Antidepressants tricyclics						
Amitriptyline	CT/P/N	CT/P ^b	CT/P ^b		CT/N	CT ^b
Desipramine		CT/P ^b	CT/P ^b			b
Imipramine		CT/P ^b	CT/P ^b			b
Nortriptyline		CT/P ^b	CT/P ^b			b
Clomipramine		CT/P ^b	CT/P ^b			
Trazadone		CT/P				
Maprotiline		CT/P ^b	CT/P ^b			b
Tetracyclics						
Mirtazapine	CS					
SSRIs						
Escitalopram		CT/P				
Fluoxetine		CT/P				
Paroxetine		CT/P				
Sertraline		OLT/P				
Bupropion		CT/P	CT/P			
SNRIs						
Duloxetine	CR	CT/P ^b				
Venlafaxine ER	CT/N	CT/P ^b				CT/P
Milnacipran	CR					
Anticonvulsants calcium channel alpha 2 delta ligands						
Gabapentin	CT/P	CT/P/N ^b	CT/P ^b	CR	CT/N	CT/P ^b
Pregabalin	CT/P	CT/P ^b	CT/P		CT/N	CT/P ^b
Carbamazepine		CT/P		CT/P ^b		
Lamotrigine		CT/P/N				CT/P/N
Oxcarbamazepine				CT/P ^b		
Phenytoin		CT/P/N				
Topiramate		CT/P/N				
Valproate		CT/P/N	CT/P			CT/N
Sodium channel blockers						
Bupivacaine (inject)	CT/P					
Lidocaine						
Infusion	CT/N	CT/P				
Patches			CT/P			
Mexiletine		CT/P/N			CT/N	CT/N
NMDA receptor antagonists						
Memantine	CR/P/N	CT/P/N	CT/N			
Ketamine ^c	CT/N					
Dextromethorphan		CT/P	CT/N			
NMDA+Potassium channel blocker					CR	
Flupirtine (plus opioids)						
Opioids						
Morphine	CT/P	CT/P	CT/P			
Methadone	CS					
Buprenorphine^d	CR/P					
Oxycodone	CR/P	CT/P	CT/P			
Levorphanol		CT(M)				
Tramadol	CT/P	CT/P	CT/P			
Muscle relaxants						
Baclofen	CR ^d		CT/P			
Benzodiazepines						
Clonazepam	CS					
Diazepam		CT/P				

(Continued)

Table 1 (Continued)

Drug	Neuropathic-pain conditions					
	PLP	PPN	PHN	TN	HIV	CP
Alprazolam		OLT				
Corticosteroids						
Prednisone		CS				
Dexamethasone		CS				
Capsaicin		CT/P/N	CT/P	CT/N		

Notes: In bold characters, reports and studies specifically designed to test drugs on PLP. ^aOnly highest level of evidence for each drug and condition reported; ^bfirst-line drugs according to EFNS guidelines; ^cintravenous infusion; ^dintrathecal analgesia.

Abbreviations: CrS, cross-sectional study; CR, case report/case series; CT, controlled trial; OLT, open label trial; M, mixed patient population; P, positive clinical trial; N, negative clinical trial; P/N, mixed clinical trial results; PLP, phantom limb pain; PPN, painful polyneuropathy; PHN, post-herpetic neuralgia; TN, trigeminal neuralgia; CP, central pain (post-stroke and spinal cord injury); NSAIDs, nonsteroidal anti-inflammatory drugs; SSRIs, selective serotonin reuptake inhibitors; SNRIs, serotonin-norepinephrine reuptake inhibitors; NMDA, N-Methyl-D-Aspartate.

for the management of PLP are mixed. Due to the availability of newer agents with similar efficacy and better profile of side effects the tricyclics have been less utilized in the last few years. The anticholinergic effect presented by these drugs is responsible for the major side effects including sedation, constipation, urinary retention, and electrical conduction changes that made them potentially problematic in the elderly. Sometimes the side effects can help to manage comorbidities and a night dose of amitriptyline might be preferable for patients with insomnia. The serotonin-norepinephrine reuptake inhibitors (SNRIs) are antidepressants that block the reuptake of serotonin and norepinephrine (eg, duloxetine) and although they have become popular due to a better profile of side effects and similar good efficacy for neuropathic pain, migraines, and fibromyalgia, the level of evidence for the management of PLP is low (Table 1). Milnacipran, in the United States, has been approved for the management of fibromyalgia, but although the level of evidence to treat PLP is low, there are studies that show benefit in neuropathic pain.³¹ Agents with a preferentially dopaminergic mechanism of action like bupropion have also been shown to be beneficial for the management of neuropathic pain. The selective serotonin reuptake inhibitors (SSRIs) are considered to be less effective but there are positive controlled trials in PPN for escitalopram, fluoxetine, and paroxetine (Table 1).²⁶

Anticonvulsants

Some anticonvulsants, like gabapentin and pregabalin, have been shown to be efficacious for the treatment of PLP in controlled trials.^{26,32,33} The list grows when the evidence of efficacy is looked at on other types of neuropathic pain like painful peripheral neuropathy and post-herpetic neuralgia and includes gabapentin, pregabalin, carbamazepine, lamotrigine, phenytoin, topiramate, and valproate (Table 1). Pain with neuralgic characteristics that can occur at the level of the

stump, can be successfully treated with the older agents like carbamazepine or the newer oxcarbazepine.

Sodium channel blockers

Sodium channel blockers can be administered orally, intravenously, epidurally, or intrathecally. The evidence that supports its use varies with the agent and the formulation. The only controlled trial that supports its use for PLP is with injectable bupivacaine. A study with lidocaine infusion was negative for PLP and another positive for PPN, while there are no positive trials for the oral formulation mexiletine (Table 1).²⁹

NMDA receptor antagonists

Intravenous infusion of ketamine, an NMDA receptor antagonist, has been shown to be beneficial for PLP but other agents of this family have not been consistently shown to be efficacious. More recently it has been utilized to manage pain that does not respond to other agents as a single intravenous infusion or repetitive infusions as outpatient. Many clinicians treat patients that experience at least 50% pain reduction with the infusion, with oral ketamine that can be compounded.²⁶ The long term use is in many occasions not possible due to side effects. There are mixed case reports with memantine, another NMDA antagonist that was developed for the management of dementia in Alzheimer's disease, for PLP and positive and negative control trials for other forms of neuropathic pain. There are positive control trials with dextromethorphan for PPN and a negative trial for PHN.

Opioids

Morphine has been shown to be beneficial for the management of PLP in a controlled trial and there is case series showing analgesia with methadone and a case report with buprenorphine reporting analgesia.^{29,38} In addition morphine,

oxycodone, levorphanol, and methadone are efficacious for the management of other forms of neuropathic pain including PPN and PHN. The long term use of these drugs has been questioned due to the scarce evidence of long term benefits including function. Concerns about addiction, diversion, and side effects including decrease in testosterone level have also been raised. Despite this, the use of opioids for the management of neuropathic pain has been supported in several guidelines including the American Pain Society (APS) and the American Academy of Pain Medicine (AAPM). Methadone has received a differential treatment due to the additional concern about cardiac toxicity and the Canadian guidelines place it at a lower step than the other opioids.

Tramadol

There are positive clinical trials with tramadol supporting its use for the management of PLP and other forms of neuropathic pain including PPN and PHN.²⁹ Tramadol is an agent with a mixed mechanism of action that includes low affinity for the mu opioid receptors and the ability to blockade the reuptake of norepinephrine and serotonin. It can also produce physical dependency but it is Schedule 3 and this is one of the reasons why it is popular among prescribers.

Others

There are other drugs that have been utilized with variable degree of success, including muscle relaxants like baclofen, the centrally acting benzodiazepines, and the corticosteroids. There is a case report for baclofen in PLP that reports benefit, a positive controlled trial in PHN, a positive controlled trial for diazepam, and an open label trial for alprazolam for the management of PPN. There are case series that report benefit with corticosteroids like prednisone dexamethasone for PPN. Capsaicin has been utilized for neuropathic pain with variable success for many years, which has not been very popular due to the pain that occurs after its application. There is a new formulation in a patch form that seems to be promising, although there is no data available yet for PLP (Table 1).

In summary, there is a limited number of control studies done specifically to assess drug efficacy in PLP.²⁶ The ones showing significant beneficial effect include trials with amitriptyline,³⁰ gabapentin,³² tramadol,⁴⁰ and morphine.³³ There are also some case reports/series with mirtazapine,⁴¹ duloxetine,³³ milnacipran,³¹ memantine,^{34,35} baclofen,³⁶ buprenorphine,³⁷ and methadone,³⁸ that suggest beneficial effects on PLP. The selection of the agent should be based on the current understanding of PLP as a neuropathic type

of condition. Hence, in addition to drugs that show efficacy in PLP, other drugs with efficacy in other types of neuropathic pain could be prescribed.³⁸ However, the drug selection should take into consideration the clinical presentation of the PLP syndrome. PLP is continuous, intense, sometimes burning and for that reason a drug with efficacy in PPN or PHN may be chosen. Further, as a neuralgic type of pain may be present in PLP (possibly a result of a neuroma formation in the stump), drugs efficacious in neuralgic type of pain, like carbamazepine or oxcarbamazepine, can be added to the regimen. A trial in every patient should include appropriate titration of the drug up to levels where efficacy has been shown in other neuropathic conditions, for at least 4–6 weeks. Thus, completion of an appropriate trial usually takes 1–2 months.³⁹

It has been proposed that the administration of analgesic and anesthetics prior to a surgical intervention^{9,42} and intra-operative precautions with the handling of peripheral nerves and not to ligate with sutures, could prevent the development of central sensitization due to impulses generated at the level of the amputation, however the results have been mixed. Epidural analgesia, ropivacaine, and patient controlled analgesia during the perioperative period have shown to decrease PLP, but ketamine and ketamine plus bupivacaine showed conflicting results.^{4,44,45}

As PLP is often of severe intensity, causing a long-lasting and debilitating decrease of quality of life in many amputees,⁴⁸ pharmacological treatment itself in many patients does not provide sufficient pain relief. Therefore, supportive treatments may enhance the treatment regimen.

Supportive non-invasive treatment approaches

Non-invasive neuromodulation

Several approaches are available within non-invasive neuromodulation in PLP, for example repetitive transcranial magnetic stimulation (rTMS), visual feedback (mirror-box therapy), or motor imagery.

Transcranial magnetic stimulation (TMS)

TMS exerts its effects on brain structures via electrical currents induced by a powerful magnetic field delivered with a magnetic coil over the scalp.⁴⁹ It can be delivered as a single pulse or as sets of pulses (rTMS). Several studies have shown that a single session of rTMS can transiently relieve pain in some patients with chronic neuropathic pain,^{50–54} and a multiple application on several consecutive days can lead to prolongation of the effects.^{55,56} However, results from rTMS

treatment specifically in patients with PLP are anecdotal, based on case-reports but not sham-controlled trials, and therefore not conclusive.^{55–57} Although the application of rTMS in PLP patients resulted in pain relief,^{55–57} short-term durability of the effects is a significant issue limiting clinical use of this treatment.

Visual feedback therapy and motor imagery

Both the visual feedback therapy (mirror-box therapy) and motor imagery are based on behaviorally relevant sensory-motor stimulation of the stump.^{58–62} A rationale of this approach arises from the existence of maladaptive neuroplastic changes underlying PLP, specifically cortical reorganization in the somatosensory and motor cortices, a relation between the cortical reorganization and the occurrence of PLP, and beneficial effects of cortical normalization on PLP.

The treatment using visual feedback emerged from the observation that looking at the reflection of an intact limb in a mirror box can induce sensations of movement in the phantom limb.^{59–65} Motor imagery is based on percepts or sensations generated internally by the brain, a mental representation of an actual sensation or movement.⁵⁸ Physiological studies have shown that both the mirror-box therapy and the motor imagery resulted in increased excitability of the corticospinal pathways.⁶⁴ This at least partially depends on the so-called mirror-neuron system⁶¹ which includes neurons that are active not only during the execution of the task itself but also during the observation of the task. This finding provides support for the idea that imagined movements share the same cortical pathways as executed motor tasks. Further, it has been shown that visual feedback actually dominates somatosensory feedback and sensory experiences can be evoked by visual information alone, and it has been shown that the visual system enhances tactile sensitivity.^{61,65}

Both the mirror therapy and the motor imagery can be very beneficial in PLP patients.^{58,60,62,63,66–68} As an example, a controlled neuroimaging study of motor imagery in PLP⁵⁸ showed evidence of cortical reorganization of motor and somatosensory cortices and its correlation with patients' pain scores prior to the motor imagery training. The training resulted in a significant decrease of intensity and unpleasantness of pain which correlated with reduction (improvement) of cortical reorganization.

Overall, the mirror-box therapy and motor imagery are safe and non-expensive additions to the PLP treatment options.

Other supportive non-invasive therapies

There is preliminary evidence that PLP patients may benefit from other non-invasive supportive treatments, such as physical therapy including the stump and phantom limb, reflexology or hypnosis,^{69,70} or peripheral transcutaneous electrical nerve stimulation of the affected limb.⁷¹ Although there are multiple reports of transcutaneous electrical nerve stimulation use in PLP, a recent systematic review⁷² revealed lack of evidence from randomized controlled trials on which effectiveness of transcutaneous electrical nerve stimulation for PLP could be judged.

Further, a use of myoelectric prosthesis may improve PLP.^{73,74} Research of stimulation-induced neural plasticity indicate that extensive behaviorally relevant stimulation of the affected part of the body leads to an expansion of its representation zone, and an intensive use of a myoelectric prosthesis has been shown to be positively linked to decreased PLP as well as decreased degree of cortical reorganization.⁷⁵ Another venue of non-invasive support to PLP patients includes various psychotherapeutic approaches.⁷⁶

Invasive treatment strategies

Surgical destructive interventions and nerve blocks

Because of a high rate of refractory chronic pain among PLP patients with failed non-invasive treatments, all kinds of invasive procedures were performed in the history of those cases.^{77–93} Destructive procedures like thermal nerve root destruction, rhizotomy, spinal ganglionectomy, and also dorsal root entry zone lesion have been performed.^{83–93} As per their nature, most of these procedures led to unrestorable damage of nervous tissue and functions. Initial results with high rates of pain relief were accompanied with a high rate of complications and followed by a high rate of recurrent pain and therefore yielding disappointing long term results after several months. Nowadays, the indication for destructive procedures is limited only to a few diagnoses for severe, refractory pain patients with short life expectancy. The dorsal root entry zone lesion is the only destructive procedure for selected patients with deafferentation pain and brachial plexus avulsion or cervical nerve root injury. In total, numbers of destructive procedures are decreasing year by year.

Non-destructive interventions such as nerve blocks have also been used in the PLP treatment. For example, interscalene blocks or stellate ganglion blocks for upper extremity PLP, or lumbar sympathetic blocks for lower extremity PLP can lead to decrease of PLP.⁹⁵ These blocks are often combined with physical therapy.

Invasive neuromodulation

Invasive neuromodulation is considered the last-resort treatment for patients who failed various trials of non-invasive treatments. On contrary to surgical destructive procedures, neuromodulatory techniques are PLP-mechanisms-driven, specifically addressing maladaptive central neuroplastic changes in the Pain Matrix (brain pain-processing networks).

Deep brain stimulation (DBS)

DBS is an electrical stimulation performed after stereotactic implantation of thin stick leads into subcortical areas such as the thalamus or basal ganglia. In the early 1950s, the first clinicians used thalamic stimulation to treat chronic pain patients.^{78–83} Since these first clinical applications a lot of case series, retrospective patient samples, and reviews were published.^{81,82,88–91,93,96,97} A recent review⁹⁸ lists a total of 600 patients with DBS due to different chronic pain syndromes. Evidence up to date suggests that the significance of DBS for PLP is controversial. However, some patients clearly benefit from DBS, experiencing long term pain relief >25% and improved quality of life.⁹⁷

Motor cortex stimulation (MCS)

MCS (Figure 1) is an electrical stimulation of the precentral gyrus using epidural surgical leads and subthreshold stimulation.^{99–103}

Historically, after the application of MCS in patients with post-stroke-pain or trigeminal neuropathic pain, also patients

Figure 1 Lateral x-ray of the skull documenting the position of the implanted paddle leads. Matching of postoperative ct-scan and preoperative MRI-3D-neuronavigation data with inserted position of the leads over the pre- and postcentral gyrus.

with PLP became candidates for the MCS treatment.^{98,104–107} Some centers perform a direct implantation of the neurostimulator in the subcutaneous tissue (infraclavicular or abdominal), while others perform also a postoperative testing with externalized extensions of the lead and an external stimulation.^{98,106,108,109} Such a testing phase allows for a placebo double-blinded stimulation period, so that false positive responders can be identified.¹⁰⁶ It has been suggested¹⁰⁶ that the definite implantation of neurostimulator would be performed only in patients with a positive effect of the real MCS and a pain reduction of >30%, and the subjective impression of pain improvement by the patient. Information from different centers and experts^{98,105,109,110} show that PLP is an accepted indication for MCS. PLP of the upper limb seems to be favorable due to the large representation on the convex part of the precentral gyrus, but interhemispheric lead implantation for the lower limb has also been reported, and a review of evidence⁹⁸ suggests that MCS yields favorable results in about 53% of PLP patients.

Spinal cord stimulation (SCS)

SCS involves the placement of electrodes in the epidural space adjacent to the spinal area presumed to be the source of pain. An electric current is then applied to achieve sympatholytic and other neuromodulatory effects.¹¹¹ Typically, the first phase of treatment involves the temporary placement of an electrical stimulator. Patients are monitored over a period of time to determine the pain reduction. Only patients positively responding to the stimulation would be considered for a permanent implantation. Clinical results indicate beneficial effects of SCS in PLP patients on immediate as well as long term outcomes,^{112,113} although the percentage of benefiting patients declined with time. For example, good results have been observed at 2 year follow-up in 52.4% of 64 PLP patients, that decreased to 39% at 5 year follow-up.¹¹³ However, given the fact that PLP is a difficult-to-treat pain syndrome and many PLP patients do not respond to other treatment strategies, SCS represents a promising treatment option for selected subpopulation of patients. Similarly to other invasive treatments, such as MCS or DBS, SCS is also reserved for patients that had not benefited from non-invasive treatment strategies.

Conclusions and future directions

Although some of the currently available therapies yield promising results, many PLP patients still remain without satisfactory pain relief. Therefore, continual advances in the PLP treatment are of high importance. It can be anticipated

that pain management in PLP patients may in the future benefit from the following venues:

- Technological and methodological improvement of existing treatment methods.

Improved formulations of pharmacological agents, such as long-acting formulations or new routes of applications, may improve pain relief and overall quality of life in patients with PLP. Further, as discussed in other sections of this article, efficacy of various neuropathic-pain medications has not yet been established for PLP. Establishing the drug efficacy specifically for PLP can certainly facilitate a patient-tailored selection of more-benefiting pharmacological regimens for PLP patients.

Further, it is anticipated that an improvement of invasive DBS and MCS treatments will be facilitated by technological advances of imaging techniques, as higher resolution would provide more accurate information about the relevant brain structures. Furthermore, technological advances in the detection of pain-related electrical signals in the brain or at its surface would enable more specific and target-oriented DBS and MCS.

- Implementation of new methods and products within current PLP treatment-strategies.

Besides the fast progress of the pharmaceutical industry and implementation of new agents, also non-pharmacological treatment strategies in PLP may enrich their repertoire of available methods. For example, within the non-invasive neuromodulatory strategy, transcranial direct current stimulation¹¹⁴ that has recently shown promising clinical potential in some neuropathic pain syndromes, may be in the future explored in the treatment of PLP. As transcranial direct current stimulation yielded pain relief in various patient-populations (eg, patients with fibromyalgia, Parkinson's disease, pelvic pain, central pain due to spinal cord injuries),^{115–119} it can be anticipated that PLP patients as well may benefit from this method. Further, a wider implementation of the use of myoelectric prosthesis in clinical practice may enrich PLP management in the future. Behaviorally relevant stimulation of the stump delivered via myoelectric prosthesis addresses cortical reorganization⁷⁵ has been identified among pathological mechanisms underlying PLP.

- Development of new treatment approaches.

Pain research incorporating new perspectives, such as genetics and epi-genetics, provides an exciting opportunity to enhance the insight into the mechanisms and development of chronic pain syndromes, including PLP, and to provide an evidence base for the development of novel

treatment approaches and targets. For example, recent studies indicate that gene *CACNG2*, which is expressed in the peripheral and central nervous system and encodes neuroprotein stargazin, may be implicated in the development neuropathic pain. If confirmed by future studies, the gene may be of high significance for PLP. Understanding genetic and epigenetic factors (ie, functionally relevant modifications to the genome that do not involve a change in the nucleotide sequence) in the development of PLP may lead to exploration of epigenetic pharmaceuticals for this difficult-to-treat condition, as recently witnessed in other areas of medicine, for example in oncology.¹²⁰

Disclosure

The authors report no conflicts of interest in this work.

References

1. Kooijman CM, Dijkstra PU, Geertzen JH, Elzinga A, van der Schans CP. Phantom pain and phantom sensations in upper limb amputees: an epidemiological study. *Pain*. 2000;87(1):33–41.
2. Jensen TS, Krebs B, Nielsen J, Rasmussen P. Non-painful phantom limb phenomena in amputees: incidence, clinical characteristics and temporal course. *Acta Neurol Scand*. 1984;70(6):407–414.
3. Nikolajsen L, Jensen TS. Phantom limb pain. *Br J Anaesth*. 2001;87(1):107–116.
4. Subedi B, Grossberg GT. Phantom limb pain: mechanisms and treatment approaches. *Pain Res Treat*. 2011;2011:861605.
5. Hirsh AT, Dillworth TM, Ehde DM, Jensen MP. Sex differences in pain and psychological functioning in persons with limb loss. *J Pain*. 2010;11(1):79–86.
6. Bosmans JC, Geertzen JH, Post WJ, van der Schans CP, Dijkstra PU. Factors associated with phantom limb pain: a 3 1/2 year prospective study. *Clin Rehab*. 2010;24(5):444–453.
7. Ephraim PL, Wegener ST, MacKenzie EJ, Dillingham TR, Pezzin LE. Phantom pain, residual limb pain, and back pain in amputees: results of a national survey. *Arch Phys Med Rehabil*. 2005;86(10):1910–1919.
8. Schley MT, Wilms P, Toepfner S, et al. Painful and non-painful phantom and stump sensations in acute traumatic amputees. *J Trauma*. 2008;65(4):858–864.
9. Borghi B, D'Addabbo M, White PF, et al. The use of prolonged peripheral neural blockade after lower extremity amputation: the effect on symptoms associated with phantom limb syndrome. *Anesth analg*. 2010;111(5):1308–1315.
10. Karanikolas M, Aretha D, Tsolakis I, et al. Optimized perioperative analgesia reduces chronic phantom limb pain intensity, prevalence, and frequency: a prospective, randomized, clinical trial. *Anesthesiology*. 2011;114(5):1144–1154.
11. Cohen LG, Bandinelli S, Findley TW, Hallett M. Motor reorganization after upper limb amputation in man. A study with focal magnetic stimulation. *Brain*. 1991;114(Pt 1B):615–627.
12. Flor H, Elbert T, Knecht S, et al. Phantom-limb pain as a correlate of cortical reorganization following arm amputation. *Nature*. 1995;375(6531):482–484.
13. Birbaumer N, Lutzenberger W, Montoya P, et al. effects of regional anesthesia on phantom limb pain are mirrored in changes in cortical reorganization. *J Neurosci*. 1997;17(14):5503–5508.
14. Weeks SR, Anderson-Barnes VC, Tsao JW. Phantom limb pain: theories and therapies. *Neurologist*. 2010;16(5):277–286.
15. Flor H. Phantom-limb pain: characteristics, causes, and treatments. *Lancet Neurol*. 2002;1(3):182–189.

16. Nystrom B, Hagbarth KE. Microelectrode recordings from transected nerves in amputees with phantom limb pain. *Neurosci Lett*. 1981;27(2):211–216.
17. Wall PD, Gutnick M. Ongoing activity in peripheral nerves: the physiology and pharmacology of impulses originating from a neuroma. *Exp Neurol*. 1974;43(3):580–593.
18. Wolpaw JR, Tennissen A. Activity-dependent spinal cord plasticity in health and disease. *Annu Rev Neurosci*. 2001;24:807–843.
19. Doubell TP, Mannion RJ, Woolf CJ. The dorsal horn: state dependent sensory processing, plasticity and the generation of pain. In: Wall PD, Melzack R, editors. *Textbook of Pain*. 4th ed. Edinburgh, UK: Churchill Livingstone; 1999:165–181.
20. Baron R. Mechanisms of disease: neuropathic pain – a clinical perspective. *Nat Clin Pract Neurol*. 2006;2(2):95–106.
21. Buonomano DV, Merzenich M. Cortical plasticity: from synapse to maps. *Annu Rev Neurosci*. 1998;21:149–186.
22. Lotze M, Montoya P, Erb M, et al. Activation of cortical and cerebellar motor areas during executed and imagined hand movements: an fMRI study. *J Cogn Neurosci*. 1999;11(5):491–501.
23. Lotze M, Flor H, Grodd W, Larbig W, Birbaumer N. Phantom movements and pain. An fMRI study in upper limb amputees. *Brain*. 2001;124(Pt 11):2268–2277.
24. Karl A, Birbaumer N, Lutzenberger W, Cohen LG, Flor H. Reorganization of motor and somatosensory cortex in upper extremity amputees with phantom limb pain. *J Neurosci*. 2001;21(10):3609–3618.
25. Karl A, Mühlhnickel W, Kurth R, Flor H. Neuroelectric source imaging of steady-state movement-related cortical potentials in human upper extremity amputees with and without phantom limb pain. *Pain*. 2004;110(1–2):90–102.
26. Alviar MJ, Hale T, Duncga M. Pharmacologic interventions for treating phantom limb pain. *Cochrane Database Syst Rev*. 2011;12:CD006380.
27. O'Connor AB, Dworkin RH. Treatment of neuropathic pain: an overview of recent guidelines. *Am J Med*. 2009;122(10 Suppl 1):S22–S32.
28. Byrant B, Knights K, Salerno E. *Pharmacology for Health Professionals*. Amsterdam, Holland: Elsevier; 2007:270.
29. Attal N, Cruccua G, Baron R, et al. EFNS guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. *Eur J Neurol*. 2010;17(9):1113–e88.
30. Robinson LR, Czerniecki JM, Ehde DM, et al. Trial of amitriptyline for relief of pain in amputees: results of a randomized controlled study. *Arch Phys Med Rehabil*. 2004;85(1):1–6.
31. Sato K, Higuchi H, Hishikawa Y. Management of phantom limb pain and sensation with milnacipran. *J Neuropsychiatry Clin Neurosci*. 2008;20(3):368.
32. Nikolajsen L, Finnerup NB, Kramp S, Vimtrup AS, Keller J, Jensen TS. A randomized study of the effects of gabapentin on postamputation pain. *Anesthesiology*. 2006;105(5):1008–1015.
33. Spiegel DR, Lappinen E, Gottlieb M. A presumed case of phantom limb pain treated successfully with duloxetine and pregabalin. *Gen Hosp Psychiatry*. 2010;32(2):228.e5–228.e7.
34. Hackworth RJ, Tokarz KA, Fowler IM, Wallace SC, Stedje-Larsen ET. Profound pain reduction after induction of memantine treatment in two patients with severe phantom limb pain. *Anesth Analg*. 2008;107(4):1377–1379.
35. Schley M, Topfner S, Wiech K, et al. Continuous brachial plexus blockade in combination with the NMDA receptor antagonist memantine prevents phantom pain in acute traumatic upper limb amputees. *Eur J Pain*. 2007;11(3):299–308.
36. Zuniga RE, Schlicht CR, Abram SE. Intrathecal baclofen is analgesic in patients with chronic pain. *Anesthesiology*. 2000;92(3):876–880.
37. Omote K, Ohmori H, Kawamata M, Matsumoto M, Namiki A. Intrathecal buprenorphine in the treatment of phantom limb pain. *Anesth Analg*. 1995;80(5):1030–1032.
38. Bergmans L, Snijdelar DG, Katz J, Crul BJ. Methadone for phantom limb pain. *Clin J Pain*. 2002;18(3):203–205.
39. Cruciani RA, Strada EA, Knotkova H. Neuropathic pain. In: Bruera E, Portenoy RK, editors. *Cancer Pain – Assessment and Management*. New York, NY: Cambridge University Press; 2010:478–505.
40. Wilder-Smith CH, Hill LT, Laurent S. Postamputation pain and sensory changes in treatment-naïve patients: characteristics and responses to treatment with tramadol, amitriptyline, and placebo. *Anesthesiology*. 2005;103(3):619–628.
41. Kuiken TA, Schechtman L, Harden RN. Phantom limb pain treatment with mirtazapine: a case series. *Pain Pract*. 2005;5(4):356–360.
42. Karanikolas M, Aretha D, Tsolakis I, et al. Optimized perioperative analgesia reduces chronic phantom limb pain intensity, prevalence, and frequency: a prospective, randomized, clinical trial. *Anesthesiology*. 2011;114(5):1144–1154.
43. Rasmussen S, Kehlet H. Management of nerves during leg amputation – a neglected area in our understanding of the pathogenesis of phantom limb pain. *Acta Anaesthesiol Scand*. 2007;51(8):1115–1116.
44. Hayes C, Armstrong-Brown A, Burstal R. Perioperative intravenous ketamine infusion for the prevention of persistent post-amputation pain: a randomized, controlled trial. *Anaesth Intens Care*. 2004;32(3):330–338.
45. Wilson JA, Nimmo AF, Fleetwood-Walker SM, Colvin LA. A randomized double blind trial of the effect of preemptive epidural ketamine on persistent pain after lower limb amputation. *Pain*. 2008;135(1–2):108–118.
46. Fromm GH, Terrence CF, Chattha AS. Baclofen in the treatment of trigeminal neuralgia: double-blind study and long-term follow up. *Ann Neurol*. 1984;15(3):240–244.
47. Pandey CK, Singh N, Singh PK. Gabapentin for refractory idiopathic trigeminal neuralgia. *Indian Med Assoc*. 2008;106(2):124–125.
48. Ehde DM, Jensen MP, Engel J, Turner JA, Hoffman AJ, Cardenas DD. Chronic pain secondary to disability: a review. *Clin J Pain*. 2003;19(1):3–17.
49. Perez MA, Cohen LG. Principles and mechanisms of transcranial direct current stimulation. In: Knotkova H, Cruciani RA, Merrick J, editors. *Brain Stimulation in the Treatment of Pain*. New York, NY: Nova Science Publishers, Inc; 2010:113–129.
50. Canavero S, Bonicalzi V, Dotta M, Vighetti S, Asteggiano G, Cocito D. Transcranial magnetic cortical stimulation relieves central pain. *Stereotact Funct Neurosurg*. 2002;78(3–4):192–196.
51. Andre-Obadia N, Peyron R, Mertens P, Manguiere F, Laurent B, Garcia-Larea L. Transcranial magnetic stimulation for pain control. Double-blind study of different frequencies against placebo, and correlation with motor cortex stimulation efficacy. *Clin Neurophysiol*. 2006;117(7):1536–1544.
52. Lefaucheur J, Hatem S, Nineb A, et al. Somatotopic organization of the analgesic effects of motor cortex rTMS in neuropathic pain. *Neurology*. 2006;67(11):1998–2004.
53. Lefaucheur JP, Drouot X, Menard-Lefaucheur I, et al. Neurogenic pain relief by repetitive transcranial magnetic cortical stimulation depends on the origin and the site of pain. *J Neurol Neurosurg Psychiatry*. 2004;75(4):612–616.
54. Hirayama A, Saitoh Y, Kishima H, et al. Reduction of intractable deafferentation pain by navigation-guided repetitive transcranial magnetic stimulation of the primary motor cortex. *Pain*. 2006;122(1–2):22–27.
55. Topper R, Foltys H, Meister IG, Sparing R, Boroojerdi B. Repetitive transcranial magnetic stimulation of the parietal cortex transiently ameliorates phantom limb pain-like syndrome. *Clin Neurophysiol*. 2003;114(8):1521–1530.
56. Khedr EM, Kotb H, Kamel NF, Ahmed MA, Sadek R, Rothwell JC. Longlasting antalgic effects of daily sessions of repetitive transcranial magnetic stimulation in central and peripheral neuropathic pain. *J Neurol Neurosurg Psychiatry*. 2005;76(6):833–838.
57. Di Rollo A, Pallanti S. Phantom limb pain: low frequency repetitive transcranial magnetic stimulation in unaffected hemisphere. *Case Report Med*. 2011;2011:130751.
58. MacIver K, Lloyd DM, Kelly S, Roberts N, Nurmikko T. Phantom limb pain, cortical reorganization and the therapeutic effects of mental imagery. *Brain*. 2008;131(Pt 8):2181–2191.

59. MacLachlan M, McDonald D, Waloch J. Mirror treatment of lower limb phantom pain: a case study. *Disabil Rehabil.* 2004;26(14-15):901-904.
60. Brodie EE, Whyte A, Niven CA. Analgesia through the looking-glass? A randomized controlled trial investigating the effect of viewing a "virtual" limb upon phantom limb pain, sensation and movement. *Euro J Pain.* 2007;11(4):428-436.
61. Moseley GL, Gallace A, Spence C. Is mirror therapy all it is cracked to be? Current evidence and future directions. *Pain.* 2008;138(1):7-10.
62. Chan BL, Witt R, Charrow AP, et al. Mirror therapy for phantom limb pain. *N Engl J Med.* 2007;357(21):2206-2207.
63. Ramachandran VS, Rogers-Ramachandran D. Synaesthesia in phantom limbs induced with mirrors. *Proc Biol Sci.* 1996;263(1369):377-386.
64. Hashimoto R, Rothwell JC. Dynamic changes in corticospinal excitability during motor imagery. *Exp Brain Res.* 1999;125(1):75-81.
65. Diers M, Christmann C, Koeppel C, Ruf M, Flor H. Mirrored, imagined and executed movements differentially activate sensorimotor cortex in amputees with and without phantom limb pain. *Pain.* 2010;149(2):296-304.
66. Moseley GL. Graded motor imagery for pathologic pain: a randomised controlled trial. *Neurology.* 2006;67(12):2129-2134.
67. McCabe C. Mirror visual feedback therapy. A practical approach. *J Hand Ther.* 2011;24(2):170-178.
68. Beaumont G, Mercier C, Michon PE, Maloun F, Jackson PL. Decreasing phantom limb pain through observation of action and imagery. *Case Series.* 2011;12(2):289-299.
69. Ulger O. Effectiveness of phantom exercises for phantom limb pain: a pilot study. *J Rehab Med.* 2009;41(7):582-584.
70. Brown CA, Lido C. Reflexology treatment for patients with lower limb amputations and phantom limb pain: an exploratory pilot study. *Complement Ther Clin Pract.* 2008;14(2):124-131.
71. Giuffrida O, Simpson L, Halligan PW. Contralateral stimulation, using TENS, of phantom limb pain: two confirmatory cases. *Pain Med.* 2010;11(1):133-141.
72. Mulvey MR, Bagnall AM, Johnson MI, Marchant PR. Transcutaneous electrical nerve stimulation (TENS) for phantom pain and stump pain following amputation in adults. *Cochrane Database Syst Rev.* 2010;(5):CD007264.
73. Sumitani M, Yozu A, Tomioka T, Yamada Y, Miyauchi S. Using the intact hand for objective assessment of phantom hand-perception. *Eur J Pain.* 2010;14(3):261-265.
74. Dietrich C, Walter-Walsh K, Preißler S, et al. Sensory feedback prosthesis reduces phantom limb pain: proof of a principle. *Neurosci Lett.* 2012;507(2):97-100.
75. Lotze M, Grodd W, Birbaumer N, Erb M, Huse E, Flor H. Does use of a myoelectric prosthesis prevent cortical reorganization and phantom limb pain? *Nat Neurosci.* 1999;2(6):501-502.
76. De Roos C, Veenstra AC, DeJongh A, et al. Treatment of chronic phantom limb pain using a trauma focused psychological approach. *Pain Res Manag.* 2010;15(2):65-71.
77. Dieckmann G, Witzmann A. Initial and long-term results of deep brain stimulation for chronic intractable pain. *Appl Neurophysiol.* 1982;45(1-2):167-172.
78. Heath RG, Mickle WA. Evaluation of 7-years' experience with depth electrode studies in human patients. In: O'Doherty DS, editor. *Electrical Studies on the Unanesthetized Brain.* New York, NY: P Hoeber; 1960:214-247.
79. Hosobuchi Y. Intracerebral stimulation for the relief of chronic pain. In: Youmans JR, editor. *Neurological Surgery.* Philadelphia, PA: WB Saunders; 1990:4128-4143.
80. Hosobuchi Y, Adams JE, Linchitz R. Pain relief by electrical stimulation of the central gray matter in human and its reversal by naloxone. *Science.* 1977;197(4299):183-186.
81. Hosobuchi Y, Adams JE, Rutkin B. Chronic thalamic stimulation for the control of facial anaesthesia dolorosa. *Arch Neurol.* 1973;29(3):158-161.
82. Kumar K, Toth C, Nath RK. Deep brain stimulation for intractable pain: a 15-year experience. *Neurosurgery.* 1997;40(4):736-746.
83. Mazars G, Ruge R, Mazars Y. Résultats de la stimulation du faisceau spino-thalamique et leur incidence sur la pathophysiologie de la douleur. *Rev Neurol.* 1960;103:136-138.
84. Mazars G, Merienne L, Cioloca C. Treatment of certain types of pain by implantable thalamic stimulators. *Neurochirurgie.* 1974;20(2):117-124.
85. Mazars GJ. Intermittent stimulation of nucleus ventralis posterolateralis for intractable pain. *Surg Neurol.* 1975;4(1):93-95.
86. Meyerson BA, Boethius J, Carlsson AM. Alleviation of malignant pain by electrical stimulation in the periventricular-periaqueductal region. Pain relief as related to stimulation sites. In: Albe-Fessard D, editor. *Advances in Pain Research and Therapy.* New York, NY: Raven Press; 1979:525-533.
87. Meyerson BA, Lindblom U, Linderöth B, Lind G, Herregodts P. Motor cortex stimulation as treatment of trigeminal neuropathic pain. *Acta Neurochir Suppl (Wien).* 1993;58:150-153.
88. Richardson DE, Akil H. Pain reduction by electrical brain stimulation in man. Part 1: Acute administration in periaqueductal and periventricular sites. *J Neurosurg.* 1977;47(2):178-183.
89. Richardson DE, Akil H. Pain reduction by electrical stimulation in man. Part 2: Chronic self-administration in the periventricular gray matter. *J Neurosurg.* 1977;47(2):184-194.
90. Kringelbach ML, Aziz TZ. Neuroethical principles of deep-brain stimulation. *World Neurosurg.* 2011;76(6):518-529.
91. Richardson DE, Akil H. Long term results of periventricular gray self-stimulation. *Neurosurgery.* 1977;1(2):199-202.
92. Spiegel EA, Wycis HT. Mesencephalotomy in treatment of intractable facial pain. *AMA Arch Neurol Psychiat.* 1953;69(1):1-13.
93. Tsubokawa T, Katayama Y, Yamamoto T, Hirayama T. Deafferentation pain and stimulation of the thalamic sensory relay nucleus: clinical and experimental study. *Appl Neurophysiol.* 1985;48(1-6):166-171.
94. Siegfried J, Cetinalp E. Neurosurgical treatment of phantom limb pain. In: Zimmermann M, editor. *Phantom and Stump Pain.* Berlin, Germany: Springer-Verlag; 1981:148-155.
95. Walsh NE, Rogers JN. Injection procedures. In: DeLisa JA, Gans BM, Walsh NE, editors. *Physical Medicine and Rehabilitation: Principles and Practice, Volume 1.* Philadelphia, PA: Lippincott Williams & Wilkins; 2005:311-361.
96. Adams JE. Naloxone reversal of analgesia produced by brain stimulation in human. *Pain.* 1976;2(2):161-166.
97. Rasche D, Ruppolt M, Stippich C, Unterberg A, Tronnier VM. Motor cortex stimulation for long-term relief of chronic neuropathic pain: a 10 year experience. *Pain.* 2006;121(2):43-52.
98. Nguyen JP, Nizard J, Keravel Y, Lefaucheur JP. Invasive brain stimulation for the treatment of neuropathic pain. *Nat Rev Neurol.* 2011;7(12):699-709.
99. Garcia-Larrea L, Peyron R, Mertens P, et al. Positron emission tomography during motor cortex stimulation for pain control. *Stereotact Funct Neurosurg.* 1997;68(1-4 Pt 1):141-148.
100. Garcia-Larrea L, Peyron R, Mertens P, et al. Electrical stimulation of motor cortex for pain control: a combined PET-scan and electrophysiological study. *Pain.* 1999;83(2):259-273.
101. Mertens P, Nuti C, Sindou M, Guenot M, Peyron R, Garcia-Larrea LB. Precentral cortex stimulation for the treatment of central neuropathic pain: results of a prospective study in a 20-patient series. *Stereotact Funct Neurosurg.* 1999;73(1-4):122-125.
102. Roux FE, Ibarrola D, Lazorthes Y, Berry I. Chronic motor cortex stimulation for phantom limb pain: a functional magnetic resonance imaging study: technical case report. *Neurosurgery.* 2001;48(3):681-688.
103. Saitoh Y, Shibata M, Hirano SI, Kobayashi K, Kasai M, Oshima H, Fukaya C. Motor cortex stimulation for central and peripheral deafferentation pain. *J Neurosurg.* 2000;92(1):150-155.
104. Katayama Y, Yamamoto T, Kobayashi K, Kasai M, Oshima H, Fukaya C. Motor cortex stimulation for phantom limb pain: Comprehensive therapy with spinal cord and thalamic stimulation. *Stereotact Funct Neurosurg.* 2001;77(1-4):159-162.

105. Lima MC, Fregni F. Motor cortex stimulation for chronic pain. Systematic review and meta-analysis of the literature. *Neurology*. 2008;70(24):2329–2337.
106. Rasche D, Tronnier VM. Invasive treatment of chronic neuropathic pain syndromes: epidural stimulation of the motor cortex. In: Knotkova H, Cruciani R, Merrick J, editors. *Brain Stimulation in the Treatment of Pain*. New York, NY: Nova Science Publishers, Inc; 2010: 69–87.
107. Rasche D, Rinaldi PC, Young RF, Tronnier VM. Deep brain stimulation for the treatment of various chronic pain syndromes. *Neurosurg Focus*. 2006;21(6):E8.
108. Fagundes-Pereira WJ, Teixeira MJ, Reynolds N, et al. Motor cortex stimulation for the treatment of neuropathic pain. *Arq Neuropsiquiatr*. 2010;68(6):923–929.
109. Fontaine D, Hamani C, Lozano A. Efficacy and safety of motor cortex stimulation for chronic neuropathic pain: critical review of the literature. *J Neurosurg*. 2009;110(2):251–256.
110. Canavero S, Bonicalzi. Therapeutic extradural cortical stimulation for central and neuropathic pain: a review. *Clin J Pain*. 2002;18(1): 48–55.
111. Oakley J, Prager J. Spinal cord stimulation: mechanisms of action. *Spine (Phila Pa 1976)*. 2002;27(22):2574–2583.
112. Viswanathan A. Use of spinal cord stimulation in the treatment of phantom limb pain: case series and review of literature. *Pain Pract*. 2010;10(5):479–484.
113. Krainick JU, Thoder U. Spinal cord stimulation in post-amputation pain. In: Siegfried J, Zimmerman M, editors. *Phantom and Stump Pain*. Berlin, Germany: Springer-Verlag; 2002:527–535.
114. Nitsche MA, Paulus W. Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *J Physiol*. 2000;527(3):633–639.
115. Fregni F, Gimenes R, Valle AS, et al. A randomized, sham-controlled, proof of principle study of transcranial direct current stimulation for the treatment of pain in fibromyalgia. *Arthritis Rheum*. 2006;54(12):3988–3998.
116. Fregni F, Boggio PS, Lima MC, et al. A sham-controlled, phase II trial of transcranial direct current stimulation for the treatment of central pain in traumatic spinal cord injury. *Pain*. 2006;122(1–2):197–209.
117. Grüner U, Eggers C, Ameli M, Sarfeld AS, Fink GR, Nowak DA. 1 Hz rTMS preconditioned by tDCS over the primary motor cortex in Parkinson's disease: effects on bradykinesia of arm and hand. *J Neural Transm*. 2010;117(2):207–216.
118. Fenton B, Fanning J, Boggio P, Fregni F. A pilot efficacy trial of tDCS for the treatment of refractory chronic pelvic pain. *Brain Stimulation*. 2008;1(3):260.
119. Knotkova H, Feldman D, Dvorkin E, Cruciani RA. Transcranial direct current stimulation (tDCS): A novel strategy to alleviate neuropathic pain. *J Pain Manag*. 2008;1(3):271–275.
120. Wang LG, Chiao JW. Prostate cancer chemopreventive activity of phenethyl isothiocyanate through epigenetic regulation (review). *Int J Oncol*. 2010;3(37):533–539.

Journal of Pain Research

Publish your work in this journal

The Journal of Pain Research is an international, peer-reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication.

Submit your manuscript here: <http://www.dovepress.com/journal-of-pain-research-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.