Relative stabilities of two difluorodiazene isomers: density functional and molecular orbital studies

Jibanananda Jana

Department of Chemistry, Krishnagar
Government College, Krishnagar,
Nadia, West Bengal, India

Abstract: The N–N bond length in the cis isomer of difluorodiazene is shorter than that in the trans isomer, giving the cis isomer higher stability. The energy partitioning approach identifies the necessary and dictating parameters responsible for the higher N–N bond energy and higher non-bonded F–F interaction energy in the cis isomer. Using density functional theory, the cis isomer is found to have higher chemical hardness and lower softness, and hence, it has higher stability than the trans isomer on the basis of the principle of maximum hardness. Localized molecular orbital study shows that the cis isomer has a higher strength of delocalization of the lone pairs of electrons on the F atoms than does the trans isomer, leading to higher stability of the cis isomer.

Keywords: cis/trans isomers, difluorodiazene, PMH, energy partitioning, localized molecular orbital, chemical hardness, softness

Introduction

It is well known¹ that difluorodiazine (dinitrogen difluoride), N₂F₂, is a gas with two isomers – cis and trans – which are shown in Figure 1. It is also known that the cis isomer predominates (~90%) at 25°C over the trans isomer.

It seems to be very obvious that the two fluorine atoms in the cis isomer, due to their greater proximity than in the trans isomer, undergo strong repulsion involving the lone pairs of electrons on the F atoms and that cis isomer is less stable than the trans one. However, the reverse observation, suggests the need to consider this problem from another angle to find a more fruitful solution.

There have been both theoretical and experimental studies regarding the relative stability of cis/trans isomers.^{2–13} Mourão and Melo,² using Mayer's energy decomposition method at the Hartree–Fock level, demonstrated that the main stabilizing effects of the cis isomer in the dihalo ethylenes are the energy terms associated with the interaction of halogen substituents with carbon atoms. With the help of the simple reversed-phase column liquid chromatographic method, Ismail et al³ showed that the higher percentage composition of the monocrotophos cis isomer in the acetonitrile eluent indicates its higher stability. Concepción et al⁴ demonstrated the kinetic preferences of isomer formation. Hush et al⁵ reported the relative thermodynamic stabilities of cis- and trans-[PtCl₂(H₂O)₂] in an aqueous solution using various methods and basis sets, concentrating largely on the role of solvation. Through gamma radiation, Alfaia et al⁷ indicated the higher susceptibility of the cis double bond to autoxidation compared to the trans configuration. Paul et al¹⁰ indicated the adjustability of the cis percentage and the conformational stability toward the intermolecular H-bonding effects. Schieber

Correspondence: Jibanananda Jana
Department of Chemistry, Krishnagar
Government College, Krishnagar,
Nadia, West Bengal, India, PIN-741 101
Email jiban_jana@yahoo.co.in

Figure 1 Optimized geometries of the cis and trans isomers of difluorodiazene (N_2F_2).

Notes: Cis- N_2F_2 ; Blue balls: Nitrogen; White balls: Fluorine; Bond length (N–N) = 1.239; Bond lengths (N–F) = 1.239; Bond angle ($\angle FNN$) = 114.59. Trans- N_2F_2 ; Blue balls: Nitrogen; White balls: Fluorine; Bond length (N–N) = 1.241; Bond lengths (N–F) = 1.239; Bond angle ($\angle FNN$) = 109.93. Bond lengths are in Å and bond angles are in degrees.

Abbreviations: $\angle FNN$, angle between N–F and N–N bonds.

and Caele¹³ demonstrated that while trans configuration of carotenoids predominates, the cis isomer takes the prominent role for food processing under the alteration of provitamin activity, bioavailability, and antioxidant capacity.

The present study investigates whether there are any differences in the geometrical parameters between the two isomers, and if such differences do exist, the reasons behind them. Further investigation will study the role of global hardness on the relative stabilities of the two isomers as well as the reason behind the closer proximity of the two *F* atoms. The study will also deal with whether there is any delocalization of the lone pairs of electrons on the *F* atoms over the F–N–N–F skeleton, and if this does happen, how it dictates the relative stabilities of the two isomers.

Figure 1 shows that the N–N bond length in the cis isomer is shorter than that in the trans isomer, whereas the N–F bonds in both isomers have the same bond length. Thus, the stability of the isomers is related to the N–N bond strength. The related questions are as follows:

- How does the N–N bond strength render more stability to the cis isomer than the trans isomer and which parameter is responsible for the formation of the strong N–N bond?

- How do the highest occupied molecular orbital and lowest unoccupied molecular orbital (*HOMO–LUMO*) energy gaps and global hardnesses in the two isomers affect the stabilities of the molecules?
- What are the probable reasons for the closer proximity of the two *F* atoms in the cis isomer? On the other hand, what is the nature of the interaction between the non-bonding *F* atoms that makes them take the more stable vicinal position in the cis isomer?
- Does delocalization affect the electron pairs on the *F* atoms with regard to the extra stability of the cis isomer? If so, what is the difference in the strength of the delocalization of the lone pairs of electrons on the *F* atoms in the two cases?

It is necessary to invoke the density functional and MO theories to investigate the relative stabilities of the two isomers. The MO energy partitioning approach is to identify the parameters responsible for the N–N bond strength and the nature of the interaction between non-bonding *F* atoms in both isomers. Localized MO (LMO) study is used to investigate the nature and strength of the delocalization of the lone pairs of electrons on the *F* atoms over the F–N–N–F skeleton in the cis and trans isomers.

In the present study, I have investigated the reasons behind the unusual stability order of the two N_2F_2 isomers. The work is organized as follows:

- The *HOMO–LUMO* energy gap ($\Delta\epsilon$), global hardness (η), and global softness (*S*) for both the isomers are calculated to investigate the relative stabilities of the two N_2F_2 isomers.
- The diatomic energy components for the N–N bonded and F–F non-bonded interaction energies in both isomers are calculated to identify the parameters necessary for the formation of the stronger N–N bond and stronger non-bonded F–F interaction in the cis isomer.
- The LMOs of both the isomers are calculated to investigate the role of delocalizing lone pairs of electrons on the *F* atoms as well as their strength of delocalization with regard to the extra stability of the cis isomer over the trans one.

The conceptual density functional theory (DFT), a new branch of quantum mechanics, offers some new fundamental global parameters: chemical potential (μ),¹⁴ chemical hardness (η),¹⁵ and its reverse concept global softness (*S*),¹⁵ of which global hardness and softness are of prime importance in the present work; μ , which characterizes the escaping tendency of electrons, was discovered by Parr et al¹⁶ as a new, important property of chemical systems. Chemical hardness indicates

the resistance of charge transfer. The higher value of η makes a chemical species harder and indicates its lower chemical reactivity. Both η and μ are successful in the rationalization of chemical processes.^{17–20} μ , η , and S are new concepts of potential importance in chemistry and are established as fundamental quantities of molecular electronic structure and reactivity.^{21–25}

In this regard, the principle of maximum hardness (PMH), which was originally introduced by Pearson,²² is the one major contribution of DFT to structural and theoretical chemistry, leading to the characterization of reactants, products, and transition states of different kinds of chemical reactions.^{20,22,25–28} According to the PMH, molecules tend to attain the state of maximum hardness when they are at equilibrium. Thus, when a system moves toward a more stable configuration, its global hardness increases and reaches its maximum when electronic energy reaches its minimum. This relationship between stability and hardness indicates that the PMH is a good descriptor of the global reactivity of a system.²⁹

Rigorous and general proof of the PMH was given by Parr and Chattaraj et al.,^{30,31} and it appears that the PMH is a fundamental, broadly applicable electronic structure rule.²⁵ Although the PMH is demonstrated to be valid under the conditions of constant external potential ($v(r)$) and chemical potential, relaxation occurs in different systems.^{32,33} As any chemical reaction involves structural and electronic reordering, the external potential hardly remains constant. Morell et al.³⁴ defined a new index in terms of the variation of hardness with respect to the external potential. The PMH has been numerically tested in a number of calculations through the computation of global hardness for a variety of individual molecules during the process of structural inversion, deformation, and internal rotation in various types of simple reactions and in correlating the order of the stability and reactivity of structurally related molecules.^{27,28,35–46} The alteration of molecular polarizability with global hardness through the distortion of bond length was investigated by Roy et al.⁴⁷

In MO theory, a molecule is nothing but a collection of nuclei and electronic charges are distributed through MOs that are fully delocalized and called canonical MOs (CMOs) or spectroscopic MOs (SMOs).⁴⁸ Although CMOs are greatly helpful to spectroscopic studies, they remain silent about routine chemistry involving lone pairs and bond pairs, ie, they do not appear in the delocalized version of the quantum chemical method. It was discovered that a set of CMOs or SMOs is one of the many possible unitary bases in the Hartree–Fock space and the new sets of MOs,

which conform to the chemical intuition in terms of lone pairs and bond pairs, can be generated by suitable unitary transformation in the same Hartree–Fock space. Such MOs are called LMOs.

In the present study, the CMOs of each molecular species are generated at their optimized geometries by the complete neglect of differential overlap (CNDO)/2 method of Pople et al.⁴⁹ The CMOs are then transformed into LMOs using the method of Trindle and Sinanoğlu.⁵⁰ Through the generation of the bond pairs and lone pairs of the isomers under investigation, the goal is to identify the nature and strength of the delocalization of the lone pairs on the F atoms, and hence, to determine the relative stabilities of the isomers.

Using the energy partitioning method of Fischer and Kollmar,⁵¹ the total energy of a molecule can be partitioned into monoatomic and diatomic parts; thus, the reasons for the stronger N–N bond formation in the cis isomer as well as the stable vicinal position of the two F atoms in the cis isomer will be investigated by analyzing the diatomic interaction energy components.

General definitions of density functional theory

It should be noted that N (the number of electrons) and $v(r)$ (the external potential) determine the perturbations as occurring in a chemical reaction. Thus, the reactivity of the molecules in terms of the system response to the variation of N and $v(r)$ can be studied under the theoretical framework of DFT.⁵² The response to changes in the number of electrons when external potential remains constant is measured at the first order by μ and at the second order by η . The variation of the external potential without charge transfer is measured by the electron density ($\rho(r)$) at the first order and by the Fukui function ($f(r)$) at the second order.⁵³ As the energy is a function of N and $v(r)$, the first order variation in the total energy in terms of the simultaneous alteration of N and $v(r)$ leads to the following expression:

$$dE = (\partial E / \partial N)_{v(r)} dN + \int (\partial E / \partial v(r))_N dv(r) dr,$$

where

$$\mu = (\partial E / \partial N)_{v(r)} \quad \text{and} \quad \rho(r) = (\partial E / \partial v(r))_N.$$

μ is a global property that characterizes the electron transfer associated with any chemical process and appears as the

Lagrange multiplier with the condition that $p(r)$ integrates to N . Now, the total differential of the chemical potential and electronic density ultimately goes to the following expressions:

$$\eta = (\partial\mu/\partial N)_{v(r)} \quad \text{and} \quad f(r) = (\partial\rho(r)/\partial N)_{v(r)} = [\delta\mu/\delta v(r)]_N.$$

Hardness can be described as the resistance to charge transfer.

Operational definitions

From the rigorous density functional definition, an exact calculation is not feasible. Operational definitions of the parameters are needed. Global hardness is obtained as

$$\eta = (I - A)/2,$$

where I and A are the ionization potential and electron affinity, respectively, can be replaced by the energy of the *HOMO* and *LUMO*, respectively, using Koopmans' theorem,⁵⁴ and are given as follows:

$$I = -\mathcal{E}_{HOMO}; \quad A = -\mathcal{E}_{LUMO}.$$

As one can arrive at the DFT-based concepts in terms of semi-empirical methods¹⁴ (eg, NDO) and because conclusions of DFT can be effectively tested by semi-empirical theories,^{22,23,28,35–37} in this investigation, the CNDO/2 method of Pople et al⁴⁹ was used, taking STO as the basis set. S , the global softness, is the inverse concept of global hardness and is defined as follows:

$$S = 1/\eta.$$

For the present purposes, the operational definition of chemical potential is not required. Thus, its operational definition is avoided here.

Energy partitioning of Fischer and Kollmar

Pople et al⁵⁵ first noted the possibility of partitioning the total molecular energy (E) obtained with the semi-empirical CNDO theory into one and two center terms:

$$E = \sum_A E_A + \sum_{A > B} E_{AB}.$$

A more detailed analysis of the two center terms was done by Gordon⁵⁶ to explain the torsional barriers. For the present purposes, the matters related to the one center term

will be set aside and the explicit formulae of only two center terms will be used. However, E_A and E_{AB} can be written as follows:

$$E_A = E_A^U + E_A^J + E_A^K \quad \text{and} \\ E_{AB} = E_{AB}^R + E_{AB}^V + E_{AB}^J + E_{AB}^K + E_{AB}^N.$$

Superscripts characterize the physical nature of the energy terms and subscripts specify the atoms. E_{AB}^V = the potential energy of the electrons on atom A in the field of nucleus B plus that of the electrons on atom B in the field of nucleus A and is expressed as follows:

$$E_{AB}^V = -P_A V_{AB} - p_B V_{BA},$$

where $P_A = \sum_{\mu \in A} P_{\mu\mu}$ and $P_{\mu\mu}$ are the elements of the bond order matrix with atomic orbital μ on atom A . Similarly, $p_B = \sum_{v \in B} p_{vv}$ and p_{vv} are the elements of the bond order matrix with atomic orbital v on atom B . V_{AB} is the potential energy of an electron on atom A in the field of nucleus B , and V_{BA} is the potential energy of an electron on atom B in the field of nucleus A .

E_{AB}^R is the contribution of resonance integrals to the energy of the A – B bond and is the main feature of the covalent bond and is expressed as follows:

$$E_{AB}^R = 2 \sum_{\mu \in A} \sum_{v \in B} p_{\mu v} \beta_{\mu v} S_{\mu v},$$

where $\beta_{\mu v}$ is a parameter that is dependent on orbitals μ and v . $S_{\mu v}$ is the overlap of the STOs μ and v .

E_{AB}^J accounts for the repulsion of the electrons on the atoms A and B and is expressed as follows:

$$E_{AB}^J = P_A P_B \gamma_{AB},$$

where γ_{AB} is the electronic repulsion between an electron on atom A and an electron on atom B .

E_{AB}^K accounts for the electronic exchange interactions and is expressed as follows:

$$E_{AB}^K = -1/2 \gamma_{AB} \sum_{\mu \in A} \sum_{v \in B} P_{\mu v}^2.$$

E_{AB}^N is the nuclear repulsion energy of the nuclei A and B and is expressed as follows:

$$E_{AB}^N = \frac{Z_A Z_B}{R_{AB}},$$

where Z_A and Z_B are the nuclear charges on A and B respectively and R_{AB} is the distance between the nuclei.

Localized molecular orbitals from the method of Trindle and Sinanoğlu

Here, the definition of the localization measure, D , is based on the idea that an electron in a local orbital should interact maximally with the electron sharing the same orbital. For a given set of doubly occupied orbitals, $(\phi_1, \dots, \phi_n)$, D is defined below as the sum of orbital coulomb energies:

$$D_N = \sum_{n=1}^N [\phi_n^2 / \phi_n^2] \quad (1)$$

$$= \sum_{n=1}^N \iint \phi_n^2(1) r_{12}^{-1} \phi_n^2(2) d\tau_1 d\tau_2$$

D will be a maximum for maximally localized orbitals. A simple iterative technique was proposed by Edmiston and Reudenberg⁵⁷ for transforming the orbitals to the set that maximized D . They considered the exactly soluble two-orbital case, molecular orbitals eg, ϕ_i and ϕ_j are ones of many possible unitary transformations and through unitary transformation, another set of molecular orbitals is generated eg, u_i and u_j , which are more localized. Now the quest is for that unitary transformation which is more localized occupied molecular orbitals. That is, what is $\sin \alpha_{ij}$ and $\cos \alpha_{ij}$ in Equation (2)?

$$\begin{vmatrix} \cos \alpha_{ij} & -\sin \alpha_{ij} \\ \sin \alpha_{ij} & \cos \alpha_{ij} \end{vmatrix} \begin{vmatrix} \phi_i \\ \phi_j \end{vmatrix} = \begin{vmatrix} u_i \\ u_j \end{vmatrix} \quad (2)$$

Maximization of D_2 with respect to the angle α_{ij} leads to the expressions in Equation (3).

$$\begin{aligned} \cos 4\alpha_{ij} &= -A_{ij} / [A_{ij}^2 + B_{ij}^2]^{1/2} \\ \sin 4\alpha_{ij} &= B_{ij} / [A_{ij}^2 + B_{ij}^2]^{1/2} \\ A_{ij} &= [\phi_i \phi_j / \phi_i \phi_j] - \frac{1}{4} [\phi_i^2 - \phi_j^2 / \phi_i^2 - \phi_j^2] \\ B_{ij} &= [\phi_i^2 - \phi_j^2 / \phi_i \phi_j] \end{aligned} \quad (3)$$

The assumptions of the CNDO/2 method on the two electron integrals allows for considerable simplification of the Edmiston–Ruedenberg transformation, leading to Equation (4), which drastically reduces the time required per iteration.

$$A_{ij} = \sum_{rt} [rr/rt] \left\{ (a_{ir}^2 - a_{jr}^2) a_{it} a_{jt} - \frac{1}{4} (a_{ir}^2 - a_{jr}^2) (a_{jt}^2 - a_{it}^2) \right\} \quad (4)$$

$$B_{ij} = \sum_{rt} [rr/tt] \{ (a_{ir}^2 - a_{jr}^2) a_{it} a_{jt} \}$$

Results and discussion

Figure 1 shows the optimized geometrical parameters of both the cis and trans isomers of difluorodiazene. This figure shows that the N–N bond length in the cis isomer is shorter than that in the trans isomer, although the N–F bond length in both cases is identical. Thus, the difference in the N–N bond strength seems to be the reason behind the extra stability of the cis isomer. Table 1 shows the various energy components of the N–N bond energy for both isomers. Table 1 demonstrates that the numerical values of E_{N-N}^J – the energy of repulsion among the electrons on one N atom with those on another N atom – and E_{N-N}^N – the nuclear repulsion energy of the two N nuclei – of the cis isomer are greater than those of the trans isomer. Thus, the greater E_{N-N}^J and E_{N-N}^N values of the cis isomer tend to reduce the N–N bond strength, but all other diatomic parameters – E_{N-N}^V , the potential energy of the electrons on one N atom in the field of the nucleus of another N atom and vice versa; E_{N-N}^K , electronic exchange interactions; and E_{N-N}^R , the contribution of resonance integrals to the energy of the N–N bond – strengthen the aforesaid bond in the cis isomer; the most contributing parameter is E_{N-N}^V . Thus, the electron–nuclear attraction potential in the N–N bond takes the lead for the formation of the stronger N–N bond in the cis isomer. Table 4 shows that the difference in E^V of the two isomers corresponding to the N–N bonded interaction is 20.22 times more than the total energy difference, ΔE , whereas the differences in E^K and E^R of the two isomers are 0.61 and 2.52 times, more than the ΔE , respectively. From the above comparative result, E_{N-N}^V is found to be the controlling parameter of the large stability of the cis isomer.

Further investigation will determine the reason for the closer proximity of the two F atoms in the cis isomer compared those in the trans isomer. Table 2 shows the various energy compared to components for the non-bonded F–F diatomic interaction energy. Table 2 demonstrates that while E_{F-F}^J and E_{F-F}^N lower the F–F interaction strength in the cis isomer, E_{F-F}^V , E_{F-F}^K , and E_{F-F}^R

Table 1 N–N bond energy components and total N–N bond energy (E_{N-N}) of the cis and trans isomers of difluorodiazene, N₂F₂

Isomers	E_{N-N}^J (au)	E_{N-N}^N (au)	E_{N-N}^V (au)	E_{N-N}^K (au)	E_{N-N}^R (au)	E_{N-N} (au)
Cis-N ₂ F ₂	9.79794	10.67745	−20.01785	−0.39094	−1.65850	−1.59190
Trans-N ₂ F ₂	9.76283	10.65889	−19.96811	−0.38944	−1.65231	−1.58814

Note: Energy values are in au.

Table 2 F–F non-bonded energy components and total F–F non-bonded interaction energy (E_{F-F}) of the cis and trans isomers of difluorodiazene, N_2F_2

Isomers	E_{F-F}^J (au)	E_{F-F}^N (au)	E_{F-F}^V (au)	E_{F-F}^K (au)	E_{F-F}^R (au)	E_{F-F} (au)
Cis- N_2F_2	11.76866	11.42220	-23.18816	-0.00233	-0.00275	-0.00238
Trans- N_2F_2	8.55705	8.29248	-16.84745	-0.00206	-0.00051	-0.00049

Note: Energy values are in au.

are responsible for the stronger F–F interaction, in which E_{F-F}^V plays the primary role. Thus, the variation trends of the diatomic energy components for the F–F non-bonded interaction energy mimic those for the bonded N–N energy.

Table 4 reveals that the difference in E^V of the two isomers corresponding to the F–F non-bonded interaction is 2577.5 times greater than ΔE whereas the differences in E^K and E^R of the two isomers are 0.11 and 0.91 times more than the ΔE respectively. Thus, the comparative study reveals that the dictating parameter of the large stability of the cis isomer is E_{F-F}^V . Further comparison involving the contributing parameters of the N–N bonded and F–F non-bonded interaction energy shows that the most dictating parameter for the larger stability of the N_2F_2 cis isomer compared to the trans isomer is E_{F-F}^V .

Table 3 shows the energy of $HOMO(\epsilon_{HOMO})$, the energy of $LUMO(\epsilon_{LUMO})$, $\Delta\epsilon$, η , S , and the total energy of both isomers. From Table 3, it can be seen that the numerical value of η in the cis isomer is 0.33681 au and 0.33474 au in the trans isomer. The cis isomer has a total energy of -78.90530 au and the trans isomer has a total energy of -78.90284 au. An analysis of these values for both isomers shows that the more stable cis isomer has a higher η value than the less stable trans isomer. Thus, the above observation is in nice conformity with the PMH.

The relative capabilities of delocalization of lone pair of electrons on F atoms over the F–N–N–F skeleton in both the cis and trans isomers will be investigated. It should be noted that any unitary transformation of MOs leaves the total energy of the system intact. The chemists' routine chemistry usually deals with the lone pairs and

bond pairs for the at a glance view on the chemical system for the explanation of a behavior. In delocalized molecular orbitals, the concept of lone pairs and bond pairs vanishes. But localized molecular orbitals generate those and thus their routine chemistry is preserved. In the present study, based on the knowledge of the nature of lone pairs, their delocalization strength is easy to detect, which is related to the contribution to the stability through the lowering of kinetic energy pressure⁵⁸ and also from the view point of particle in a 3D box model.

Table 5 shows the LMOs of the cis isomer. Table 5 demonstrates that LMO 2l.p.(1F), the second lone pair on the first F atom, and LMO 3l.p.(2F), the third lone pair on the second F atom, become delocalized over the F–N–N–F skeleton to diminish the kinetic energy pressure from those electron pairs, leading to their strong contribution toward the overall stability of the cis isomer. Table 6 displays the LMOs of the trans isomer. From Table 6, it can be seen that LMO 1l.p.(1F), the first lone pair on the first F atom, and LMO 3l.p.(2F), the third lone pair on the second F atom, become delocalized over the F–N–N–F skeleton. Thus, in the trans isomer, like in the cis isomer, the third lone pair on the second F atom participates in the delocalization over the F–N–N–F skeleton; however, unlike the cis isomer, the first lone pair on the first F atom in the trans isomer takes the role of delocalization over the previous framework. Table 5 shows that the coefficient of $2p_z$ orbital on the first F atom for the formation of LMO 2l.p.(1F) and the coefficient of the $2p_z$ orbital on the second F atom for the formation of LMO 3l.p.(2F) have the same value (-0.9850). In LMO 2l.p.(1F), the $2p_z$ orbital on the first F atom is in π -type interaction with the

Table 3 Energy of $HOMO$, ϵ_{HOMO} (au); energy of $LUMO$, ϵ_{LUMO} (au); global hardness, η (au); global softness, S (au); and total energy, E (au), of the cis and trans isomers of difluorodiazene, N_2F_2

Isomers	Energy of $HOMO$ ϵ_{HOMO} (au)	Energy of $LUMO$ ϵ_{LUMO} (au)	$HOMO-LUMO$ energy gap $\Delta\epsilon$ (au)	Global hardness η (au)	Global softness S (au)	Total energy E (au)
Cis- N_2F_2	-0.56714	0.10648	0.67362	0.33681	2.96903	-78.90530
Trans- N_2F_2	-0.56671	0.10276	0.66947	0.33474	2.98739	-78.90284

Abbreviations: $HOMO$, highest occupied molecular orbital; $LUMO$, lowest unoccupied molecular orbital.

Table 4 Difference in the energy parameters (E^V , E^K , and E^R) for both the N–N bonded and F–F non-bonded interactions and difference in total energy (ΔE) of the cis and trans isomers of difluorodiazene, N_2F_2

Interaction types	ΔE^V (au)	ΔE^K (au)	ΔE^R (au)	ΔE (au)
N–N bonded	0.04974	0.00150	0.00619	0.00246
F–F non-bonded	6.34071	0.00027	0.00224	

Note: Energy values are in au.

$2p_z$ orbitals on the second F , first N , and second N . Similarly, in LMO 3l.p.(2F), the $2p_z$ orbital on the second F atom is in π -type interaction with the $2p_z$ orbital on the first F , first N , and second N . Further, Table 6 shows that the coefficient of the $2p_z$ orbital on the first F atom for the formation of LMO 1l.p.(1F) and the coefficient of the $2p_z$ orbital on the second F atom for the formation of LMO 3l.p.(2F) have the same value (-0.9858). In the formation of the previously mentioned LMOs of the trans isomer, like the cis isomer, the $2p_z$ orbital on one F atom is in π -type interaction with the $2p_z$ orbital on the other F atom and both the N atoms.

A comparison of Tables 5 and 6 shows that in Table 5, the second lone pair on the first F atom in LMO 2l.p.(1F) and the third lone pair on the second F atom in LMO 3l.p.(2F) of the cis isomer are delocalized to a greater extent than in the trans isomer, whose LMOs are shown in Table 6. Table 6 shows the greater contributions of the $2p_z$ atomic orbitals on 1F and 2F for the formation of LMOs 1l.p.(1F) and 3l.p.(2F), respectively, which indicates the higher charge accumulation on those F atoms compared to the population on the F atoms in the cis isomer. There is then greater reluctance of those lone pairs on

the F atoms in the trans isomer toward delocalization. Thus, the lower delocalization abilities of those lone pairs on the F atoms in the trans isomer make it less stable. Further comparison of Tables 5 and 6 shows that the coefficients of the $2p_z$ orbitals on the N atoms and F atom (other than the most contributing one) toward LMO 3l.p.(2F) and 1l.p.(1F) formation in the trans isomer (Table 6) are smaller than those toward the formation of LMOs 2l.p.(1F) and 3l.p.(2F) in the cis isomer, indicating a higher charge density on the N atoms in the cis form, and hence, a higher delocalization capacity of the lone pairs of electrons on the F atoms in the cis isomeric form.

Thus, from the above LMO study on both isomers, it is very much expected that the higher delocalization strengths of the delocalizing lone pairs on the F atoms in the cis isomer make the cis isomer relatively more stable than the trans isomer by lowering the kinetic energy pressure⁷² from those lone pairs, and also from the view point of particle in a 3D box model.

Conclusion

Figure 1 shows that the N–N bond length in the cis isomer of N_2F_2 is shorter than that in the trans isomer, and this difference in geometrical parameter leads to the higher stability of the cis isomer, which is nicely reflected in the DFT study. In conformity with the PMH, it is found that the more stable cis isomer has higher global hardness and lower global softness than the trans isomer.

Using CNDO/2 energy partitioning, it is found that the electron–nuclear attraction potential (E_{N-N}^V) involving the electron on one N with the nucleus of another N , and vice

Table 5 Localized molecular orbitals (LMOs) of cis-difluorodiazene, N_2F_2

LMOs/ Atomic orbitals	1l.p.(1N)	σ (1N-1F)	1l.p.(1F)	σ (N-N)	1l.p.(2F)	π (N-N)	σ (2N-2F)	2l.p.(1F)	1l.p.(2N)	3l.p.(1F)	2l.p.(2F)	3l.p.(2F)
1N _{2s}	-0.7715	0.2526	0	0.3551	-0.0215	0	0.0323	0	0.0285	-0.0281	0.0025	0
1N _{2px}	0.1676	-0.4076	-0.0001	0.5304	-0.0372	0	-0.0371	0	-0.0471	-0.0427	-0.0036	0
1N _{2py}	0.6082	0.4260	-0.0009	0.3041	-0.0171	0	0.0203	0	0.0488	-0.0233	0.0041	0
1N _{2pz}	0	0	0	0	0	0.7069	0	-0.1329	0	0	0	0.1086
2N _{2s}	0.0283	0.0324	0.0023	0.3551	0.0281	0	0.2526	0	-0.7715	0.0215	0	0
2N _{2px}	-0.0014	0.0120	-0.0004	-0.5901	-0.0466	0	-0.0131	0	-0.5486	-0.0384	0.0006	0
2N _{2py}	0.0678	0.0406	0.0056	-0.1601	-0.0136	0	0.5894	0	0.3115	-0.0142	-0.0006	0
2N _{2pz}	0	0	0	0	0	0.7069	0	0.1086	0	0	0	-0.1329
1F _{2s}	-0.0005	0.3303	-0.8962	0	-0.0012	0	-0.0174	0	-0.0159	0.0998	-0.0018	0
1F _{2px}	-0.0010	0.5146	0.2469	-0.0059	0	0	-0.0286	0	-0.0249	-0.6949	-0.0028	0
1F _{2py}		-0.4583	-0.3685	-0.0043	0.0034	0	0.0251	0	0.0223	-0.7084	0.0025	0
1F _{2pz}	0	0	0	0	0	-0.0171	0	-0.9850	0	0	0	-0.0146
2F _{2s}	-0.0162	-0.0174	-0.0018	0	-0.0996	0	0.3304	0	-0.0005	0.0013	-0.8962	0
2F _{2px}	0.0018	0.0024	0.0001	0.0072	-0.9923	0	-0.0396	0	0	0.0025	0.0858	0
2F _{2py}	0.0334	0.0379	0.0037	0.0012	0.0095	0	-0.6879	0	0.0014	-0.0023	-0.4352	0
2F _{2pz}	0	0	0	0	0	-0.0171	0	-0.0146	0	0	0	-0.9850

Table 6 Localized molecular orbitals (LMOs) of trans-difluorodiazene, N₂F₂

LMOs	$\sigma(\text{N-N})$	1l.p.(1N)	1l.p.(2N)	1l.p.(2F)	$\sigma(1\text{N}-1\text{F})$	2l.p.(2F)	3l.p.(2F)	1l.p.(1F)	2l.p.(1F)	3l.p.(1F)	$\sigma(2\text{N}-2\text{F})$	$\pi(\text{N-N})$
Atomic orbitals												
1N _{2s}	0.3508	-0.7752	0.0201	0.0293	-0.2495	-0.0008	0	0	-0.0004	0.0271	-0.0306	0
1N _{2px}	0.1219	0.5257	0.0428	0.0228	-0.5303	-0.0021	0	0	-0.0005	0.0074	-0.0383	0
1N _{2py}	0.6017	0.3454	-0.0204	0.0308	0.2563	0.0018	0	0	-0.0006	0.0462	0.0277	0
1N _{2pz}	0	0	0	0	0	0	0.1063	-0.1292	0	0	0	-0.7069
2N _{2s}	0.3508	-0.0200	0.7752	-0.0269	0.0306	-0.0004	0	0	-0.0008	-0.0292	0.2495	0
2N _{2px}	-0.1220	0.0430	0.5258	0.0075	-0.0386	0.0005	0	0	0.0023	0.0228	-0.5302	0
2N _{2py}	-0.6016	-0.0204	0.3454	0.0463	0.0278	0.0007	0	0	-0.0017	0.0308	0.2564	0
2N _{2pz}	0	0	0	0	0	0	-0.1292	0.1063	0	0	0	-0.7069
1F _{2s}	0.0011	-0.0004	-0.0113	-0.0049	-0.3313	0.0008	0	0	-0.8936	-0.1188	0.0193	0
1F _{2px}	-0.0038	0.0009	0.0203	0.0079	0.5940	-0.0015	0	0	-0.4366	0.4537	-0.0348	0
1F _{2py}	-0.0054	-0.0006	-0.0119	-0.0074	-0.3506	0.0009	0	0	0.1044	0.8801	0.0233	0
1F _{2pz}	0	0	0	0	0	0	-0.0139	-0.9858	0	0	0	0.0162
2F _{2s}	0.0012	0.0114	0.0004	0.1196	-0.0193	-0.8934	0	0	0.0008	0.0048	0.3313	0
2F _{2px}	0.0038	0.0202	0.0007	0.4533	-0.0346	0.4370	0	0	0.0014	0.0079	0.5940	0
2F _{2py}	0.0054	-0.0117	-0.0008	0.8802	0.0232	-0.1036	0	0	-0.0009	-0.0074	-0.3506	0
2F _{2pz}	0	0	0	0	0	0	-0.9858	-0.0139	0	0	0	0.0162

versa, is greater in the cis isomer compared to the trans isomer, leading to a higher N–N bond energy in the cis isomer. The exchange energy ($E_{\text{N-N}}^{\text{K}}$) and resonance energy ($E_{\text{N-N}}^{\text{R}}$) vary in the same pattern as that of ($E_{\text{N-N}}^{\text{V}}$) though variation of ($E_{\text{N-N}}^{\text{J}}$) and ($E_{\text{N-N}}^{\text{N}}$) run in the opposite direction and tend to decrease the N–N bond energy in the cis isomer.

Further, an energy partitioning study on the non-bonded F–F interaction energy was undertaken to identify the reason for the closer proximity of the two F atoms in the cis isomer. The study shows that like the N–N bond energy partitioning, $E_{\text{F-F}}^{\text{V}}$, $E_{\text{F-F}}^{\text{K}}$, and $E_{\text{F-F}}^{\text{R}}$ tend to increase the interaction energy when $E_{\text{F-F}}^{\text{V}}$ takes the lead and the opposing parameters, $E_{\text{F-F}}^{\text{J}}$ and $E_{\text{F-F}}^{\text{N}}$, are overcome by the results of the former parameters.

The LMO study shows that the second lone pair on the first F atom and the third lone pair on the second F atom in the cis isomer become delocalized in a larger amount than the third lone pair on the second F atom and the first lone pair on the first F atom in the trans isomer. Thus, the lower kinetic energy pressure due to the higher delocalization capabilities of the lone pairs in the cis isomer is expected to render higher stability to the cis isomer, and also the “particle in a 3D box” model sets its importance here.

Acknowledgment

The author is indebted to the computer center at the Department of Chemistry of this College for providing computer facilities, and the author extends his thanks to Dr Saikat Sarkar, Department Of Chemistry, Santipur College, West Bengal, INDIA, for all of his encouragement.

Disclosure

The author reports no conflicts of interest in this work.

References

1. Cotton FA, Wilkinson G, Murillo CA, Bochmann M. *Advanced Inorganic Chemistry*. 6th ed. USA: John Wiley & Sons Ltd; 2008:337.
2. Mourão ZS, Melo A. Energy decomposition analysis of cis and trans isomers of 1,2-dihaloethylenes and 2-butene. *J Mol Str THEOCHEM*. 2010;946(1-3):7-12.
3. Ismail N, Vairamani M, Rao R N. Determination of cis and trans isomers of monocrotophosin technical products reversed phase column liquid chromatography. *J Chromatogr A*. 2000;903(1-2):255-260.
4. Concepción J, Loeb B, Simón Manso Y, Zuloaga F. Influence of L-type ligand on the relative stability and inter conversion of cis-trans [Ru(Phen)₂L₂]²⁺ type complexes – a theoretical study. *Polyhedron*. 2000;19(22-23):2297-2302.
5. Hush NS, Schamburger J, Bacskey GB. A quantum chemical computational study of the relative stabilities of cis and trans platinum dichloride in aqueous solution. *Coord Chem Rev*. 2005;249(3-4):299-311.
6. Dolgonos G. Relative stability and thermodynamic properties of Si₂H₄ isomers. *Chem Phys Lett*. 2008;466(1-3):11-15.
7. Alfaia CMM, Riberio PJLC, Trigo MJP, et al. Irradiation effect on fatty acid composition and conjugated linoleic acid isomers in frozen lamb meat. *Meat Sci*. 2007;77(4):689-695.
8. Senthilkumar K, Kolandaivel P. Molecular structure conformational stability and cis effect of 1,4-dichlorobutadiene – a quantum chemical study. *J Mol Str THEOCHEM*. 2002;577(1):69-79.
9. Kuznetsov ML, Hankka M, Pombeiro JL, Nazarov AA, Kukushkin VY. Theoretical study of the relative stability of isomeric forms of platinum carboxamide complexes. *Inorganica Chim Acta*. 2003;350:245-251.
10. Paul A, Bittermann H, Gmeiner P. Triazolepeptides: chiro specific synthesis and cis/trans prolyl ratios of structural isomers. *Tetrahedron*. 2006;62(38):8919-8927.
11. Kang YK. Cis-trans isomerization of N-acetyl-N'-methyl amides of 5-methyl proline and 5,5-dimethylproline. *J Mol Str THEOCHEM*. 2002;585(1-3):209-221.
12. Dasgupta B, Chakraborti P, Basu G. Enhanced stability of cis pro-pro peptide bed in pro-pro-phe sequence motif. *FEBS Lett*. 2007;581(23):4529-4532.

13. Schieber A, Caele R. Occurrence of carotenoid isomers in food technological, analytical, and nutritional implications. *Trends Food Sci Technol*. 2005;16(9):416–422.
14. Geerlings P, De Proft F, Langenaeker W. Conceptual density functional theory. *Chem Rev*. 2003;103(5):1793–1873.
15. Chermette H. Chemical reactivity indexes in density functional theory. *J Comput Chem*. 1999;20(1):129–154.
16. Parr RG, Donnelly RA, Levy M, Palke WE. Electronegativity: the density functional view point. *J Chem Phys*. 1978;68(18):3801–3807.
17. Jaque P, Toro-Labbé A. Characterization of copper cluster through the use of density functional theory reactivity descriptors. *J Chem Phys*. 2002;117(7):3208–3218.
18. Gutierrez-oliva S, Jaque P, Toro-Labbé A. Using Sanderson's principle to estimate global electronic properties and bond energies of hydrogen-bonded complexes. *J Phys Chem A*. 2000;104(39):8955–8964.
19. Perez P, Toro-Labbé A. Characterization of keto enol tautomerism of acetyl derivatives from the analysis of energy: chemical potential and hardness. *J Phys Chem A*. 2000;104(7):1557–1562.
20. Toro-Labbé A. Characterization of chemical reactions from the profiles of energy, chemical potential and hardness. *J Phys Chem A*. 1999;103(22):4398–4403.
21. Zhou Z, Parr RG. Activation hardness: new index for describing the orientation of electrophilic aromatic substitution. *J Am Chem Soc*. 1990;112(15):5720–5724.
22. Pearson RG. Recent advances in the concept of hard and soft acids and bases. *J Chem Educ*. 1987;64(7):561–567.
23. Pearson RG. Maximum chemical and physical hardness. *J Chem Educ*. 1999;76(2):267–274.
24. Pearson RG. Electronic spectra and chemical reactivity. *J Am Chem Soc*. 1988;110(7):2092–2097.
25. Pearson RG. The principle of maximum hardness. *Acc Chem Res*. 1993;26(5):250–255.
26. Gazquez JL. Bond energies and hardness differences. *J Phys Chem A*. 1997;101(49):9464–9469.
27. Sola M, Toro-Labbé A. The Hammond postulate and the principle of maximum hardness in some intramolecular rearrangement reactions. *J Phys Chem A*. 1999;103(44):8847–8852.
28. Datta D. On Pearson's HSAB principle. *Inorg Chem*. 1992;31(13):2797.
29. Torrent-Sucarrat M, J. Luis JM, M. Duran, Sola M. The hardness profile as a tool to detect spurious stationary points in the potential energy surface. *J Chem Phys*. 2004;120(23):10914–10924.
30. Parr RG, Chattaraj PK. Principle of maximum hardness. *J Am Chem Soc*. 1991;113(5):1854–1855.
31. Chattaraj PK, Lee H, Parr RG. HSAB principle. *J Am Chem Soc*. 1991;113(5):1855–1856.
32. Gutierrez-oliva S, Jaque P, Toro-Labbé A. *Reviews in Modern Quantum Chemistry: A Celebration with Contributions of RG Parr and KD Sen*. Singapore: World Scientific Press; 2002.
33. Torrent-Sucarrat M, Luis J, Duran M, Sola M. On the validity of the maximum hardness and minimum polarizability principles for nontotally symmetric vibrations. *J Am Chem Soc*. 2001;123(32):7951–7952.
34. Morell C, Grand A, Toro-Labbé A. New dual descriptor for chemical reactivity. *J Phys Chem A*. 2005;109(1):205–212.
35. Ghosh DC, Jana J, Biswas R. Quantum chemical study of the umbrella inversion of the ammonia molecule. *Int J Quantum Chem*. 2000;80(1):1–26.
36. Ghosh DC, Jana J, Bhattacharyya S. Density functional and molecular orbital study of physical process of inversion of nitrogen trifluoride (NF_3) molecule. *Int J Quantum Chem*. 2002;87(3):111–134.
37. Ghosh DC. Density functional and frontier orbital study of the physical process of the conformational isomerism of ethane. *J Ind Chem Soc*. 2002;79(3):240–248.
38. Nguyen TN, De Proft F, Nguyen MT, Geerlings P. Theoretical study of [2+1] cycloaddition of CO and CS to acetylenes forming cyclopropanones and cyclopropenethiones. *J Org Chem*. 2001;66(12):4316–4326.
39. Nguyen LT, De Proft F, Nguyen MT, Geerlings P. Theoretical study of cyclopropanones and cyclopropenethiones: principles decomposition via intermediates. *J Chem Soc Perkin 1*. 2001;2(6):898–905.
40. Chandra AK, Nguyen MT. Density functional approach to regiochemistry, activation energy, and hardness profile in 1,3-dipolar cycloadditions. *J Phys Chem A*. 1998;102(30):6181–6185.
41. Cardens-Jiron GI, Gutierrez-oliva S, Melin J, Toro-Labbé A. Relations between potential energy, electronic chemical potential, and hardness profiles. *J Phys Chem A*. 1997;101(25):4621–4627.
42. Cardens-Jiron GI, Letelier JR, Toro-Labbé A. The internal rotation of hydrogen thioperoxide: energy, chemical potential, and hardness profiles. *J Phys Chem A*. 1998;102(40):7864–7871.
43. Ghanty TK, Ghosh SK. Molecular hardness, polarizability, and valency variation of formamide and thioformamide on internal rotation: a density functional study. *J Phys Chem A*. 2000;104(13):2975–2979.
44. Uchimaru T, Chandra AK, Kawahara S, Matsumura K, Tsuzuki S, Mikami M. Internal bond rotation in substituted methyl radicals, H_2B-CH_2 , H_3C-CH_2 , H_2N-CH_2 and $HO-CH_2$: hardness profiles. *J Phys Chem A*. 200;105(8):1343–1353.
45. Nguyen MT, Chandra AK, Sakai S, Morokuma K. Another look at the mechanism of the concerted 1,3-dipolar cycloaddition of fulminic acid to acetylene. *J Org Chem*. 1999;64(1):65–69.
46. Kar T, Scheiner S, Sannigrahi AB. Ab initio calculations of hardness and chemical potential of open shell systems using SCF, MP2 and MP4 methods. *J Mol Struct THEOCHEM*. 1998;427(1–3):79–85.
47. Roy RK, Chandra AK, Pal S. Correlation of polarizability, hardness and electronegativity: polyatomic molecules. *J Phys Chem*. 1994;98(41):10447–10450.
48. Martin RB. Localized and spectroscopic orbitals: Squirrel ears on water. *J Chem Educ*. 1988;65(8):668–670.
49. Pople JA, Beveridge DL. *Approximate Molecular Orbital Theory*. New York: Mc Graw-Hill; 1970.
50. Trindle C, Sinanoğlu O. Semiempirical method for the determination of localized orbitals in molecules. *J Chem Phys*. 1968;49(1):65–71.
51. Fischer H, Kollmar H. Energy partitioning with the CNDO method. *Theor Chim Acta*. 1970;6(3):163–174.
52. Geerlings P, De Proft F, Langenaeker W. *Density Functional Theory: A Bridge between Chemistry and Physics*. VUB University Press: 1998.
53. Parr RG, Mortier WJ. The use of global and local molecular parameters for the analysis of the gas phase basicity of amines. *J Am Chem Soc*. 1986;108(19):5708–5711.
54. Koopmans T. Ordering of wave functions and eigenenergies to the individual electrons of an atom. *Physica*. 1933;1(1–6):104–113.
55. Pople JA, Santry DP, Segal GA. Approximate self-consistent molecular orbital theory. I. invariant procedures. *J Chem Phys*. 1965;43(10):S129.
56. Gordon MS. Molecular orbital study of internal rotation. *J Am Chem Soc*. 1969;91(12):3122–3140.
57. Edmiston C, Reudenberg K. Localized atomic and molecular orbitals. *Rev Mod Phys*. 1963;35(3):457–464.
58. Reudenberg K. The physical nature of the chemical bond. *Rev Mod Phys*. 1962;34(2):326–376.

Reports in Theoretical Chemistry**Dovepress****Publish your work in this journal**

Reports in Theoretical Chemistry is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of theoretical chemistry. The manuscript management system is completely online and includes a very quick and fair

peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/reports-in-theoretical-chemistry-journal>