Mutational analysis of *APOL1* in patients with Fechtner and Epstein syndromes: no evidence of a digenic etiology in *MYH9*-related disorders with renal disease

Maxine McGredy¹
John A Martignetti¹⁻³
Melanie Babcock¹

¹Departments of Genetics and Genomic Sciences, ²Pediatrics, ³Oncological Sciences, Mount Sinai School of Medicine, New York, NY, USA

Abstract: Mutations in *MYH9* result in a group of clinically overlapping autosomal dominant macrothrombocytopenia syndromes, collectively termed the *MYH9*-related disorders (MYH9RD). Intriguingly, three of these disorders, Fechtner, Alport-like, and Epstein syndromes, are associated with the additional clinical phenotype of glomerulosclerosis. Recently, an abundance of studies have demonstrated that the *APOL1* gene, contiguous to *MYH9*, is associated with a form of kidney disease in individuals of African ancestry. Given these findings related to kidney disease arising in mutations in two contiguous genes, this study aimed to determine whether *APOL1* mutations could also be present in patients with Fechtner (FTNS) and Epstein (EPS) syndromes. This study used sequence analysis to investigate a discrete discovery set of FTNS patients, but did not identify second hit mutations in *APOL1*.

Keywords: *APOL1*, *MYH9*, Fechtner syndrome, Epstein syndrome

Introduction

Mutations in nonmuscle myosin heavy chain IIa, *MYH9*-cytoskeletal contractile protein, result in a spectrum of macrothrombocytopenia disorders collectively named “the giant-platelet” disorders, or the *MYH9*-related disorders (MYH9RD). These include May-Hegglin anomaly (MHA, MIM #155100), Sebastian platelet syndrome (SBS, MIM #605249), Alport-like syndrome (APSM), Epstein syndrome (EPS, MIM #153650), and Fechtner syndrome (FTNS, MIM #153640).^{1,2} These disorders are generally considered to be rare in the population; however, as they can often be misdiagnosed as immune thrombocytopenia (ITP), the prevalence of MYH9RD may be underestimated.¹ While all patients share the clinical features of macrothrombocytopenia and characteristic leukocyte inclusions (Döhle-like bodies), some will also display high-tone sensorineural deafness and/or the development of pre-senile cataracts.³⁻⁶ Of particular interest, a number of individuals with MYH9RD also develop kidney disease, which can range from glomerulosclerosis to end-stage renal disease.⁷⁻¹⁰ It should also be noted that *MYH9* has been identified to play a role in nonsyndromic hereditary deafness.¹³ In accordance with its role in hearing/deafness, *MYH9* expression was identified in rat cochlea.¹³

MYH9 is expressed in a number of tissues, but of particular interest with regard to disease pathology, it is expressed in platelets, kidney,¹¹ leukocytes,¹² and the cochlea.¹³ The N-terminus of the protein contains the motor domain, which binds to actin and

Correspondence: John A Martignetti
Mount Sinai School of Medicine,
Department of Genetics and Genomic
Sciences, Pediatrics, and Oncological
Sciences, 1425 Madison Avenue, Room
14-26D, New York, NY 10029, USA
Tel +1 212 659-6744
Fax +1 212 360-1809
Email john.martignetti@mssm.edu

ATP, while the C-terminus contains the regulatory region.¹⁴ Genotype-phenotype correlations with specific *MYH9* mutations have been investigated by others.^{15,16} Affected individuals with a mutation within the motor domain, compared to the tail domain, of the MYH9 are at higher risk, but not certain risk, of having a more severe phenotype.¹⁵ For example, a small percentage of patients with the tail domain 1424 mutation have a more severe phenotype.¹⁵ Sekine et al demonstrated that nine patients with R702 mutations developed renal disease,¹⁶ whereas Pecci et al determined that 32% of patients with R702 mutations developed nephritis.¹⁵ Thus, while associations can be made, no universally acceptable genotype-phenotype correlation completely predicts the clinical spectrum of each *MYH9* mutation.

Most recently, *MYH9* was proposed as a potential risk factor in nondiabetic end-stage kidney disease (ESKD) or renal disease (ESRD).⁷ Mapping by admixture linkage disequilibrium (MALD) indicated that *MYH9* maps to a chromosomal region that is associated with ESKD risk variants.^{7,8} African Americans have a 3.7 times increased risk of ESRD, significantly higher than individuals of European descent (US Renal Data System 2007). Various intronic single nucleotide polymorphisms (SNPs) have presented as kidney risk variants, but even after resequencing of the *MYH9* gene, no functional mutations were discovered.^{17,18} A genome wide association study (GWAS) identified a region that contained both *MYH9* and *APOL1* and included a risk allele for focal segmental glomerulosclerosis (FSGS) in the African American population but not the European population.¹⁹ Given these ethnic differences, the authors suggested that *APOL1* represented a possible disease-gene candidate because of its relationship in defining disease resistance to a lethal parasite, the trypanosome, which is common in Africa.¹⁹

APOL1 is a member of a gene family consisting of six APOL genes, L1-L6,^{20,21} which are involved in cellular apoptosis,²² trypanosomal lysis,²³ and lipid metabolism.^{24,25} APOL1 has the ability to remain sequestered as an intracellular protein or to be secreted.²⁵ Freedman and colleagues studied novel SNP frequencies in two populations of African Americans: one containing 205 patients with FSGS, and another containing 180 control patients.²⁶ They presented evidence that mutations in the last exon of *APOL1* – rs73885319 (S342G) and rs60910145 (I384M) – were causative.²⁶

Based on the studies of *APOL1* in the involvement in ESKD, we investigated its role in Fechtner and Epstein syndromes. The initial linkage block that was identified included the genes *APOL1* and *MYH9*,⁸ which raised the possibility that FTNS and EPS are both digenic diseases.

In fact, some forms of renal disease have been suggested result from digenic mutations.^{27,28} Hypothetically, a second hit, a mutation in *APOL1*, could be the causative difference among the macrothrombocytopenia disorders that have renal disease. Given the renal phenotype of MYH9RD and the renal disease now linked to the adjacent gene *APOL1*, we were curious to examine the possibility that mutations in both genes are necessary to give rise to the full spectrum of MYH9RD. We therefore performed mutational analysis of four DNA samples from individuals with Fechtner or Epstein syndromes.

Methods

Patients and patient selection

Genomic DNA from four MYH9RD individuals shown previously to harbor *MYH9* mutations was analyzed for possible concurrent mutations in *APOL1* (Table 1). Information regarding ethnicity and clinical history was obtained. Patient 1 was a pediatric patient of African descent who presented with a diagnosis of Fechtner syndrome and died as a result of end stage renal failure at the age of sixteen. Patient 2 was of French heritage and had been clinically diagnosed with Epstein syndrome. The patient's mother had originally presented with symptoms of deafness and kidney failure requiring dialysis. Patient 3 was of Italian descent and diagnosed with Fechtner syndrome. Patient 4 was of French descent and presented with Fechtner syndrome. All samples collected were approved by the IRB.

DNA sequence, mutation analysis

Primers encompassing the coding regions and flanking intron/exon boundaries of *APOL1* were designed to amplify the *APOL1* transcript variant 2, NM_145343, as well as NM_003661.3, NM_001136540.1, and NM_001136541.1 (Table 2). The *MYH9* primers have been previously described.² For exons 1–6 of *APOL1* and 7B–7E, reactions were performed in 20 µL of a final concentration of 1X PCR Buffer (GoTaq, Promega Corporation, Madison, Wisconsin, USA) containing 0.25 µM of each primer, and 40 ng of genomic DNA. For exon 7A, reactions were performed in 25 µL of a final concentration of 2.5 mM MgCl₂, 1X PCR

Table 1 MYH9 mutations

	MYH9 protein alterations
Patient 1	R702C
Patient 2	S96L
Patient 3	D1424H
Patient 4	R702G

Table 2 PCR primer sequences for the amplification of *APOLI* from genomic DNA

Primer name	Forward primer	Reverse primer	Annealing temperature (°C)	Product size
<i>APOLI</i> EX1	GCTCCTCTAAGGGCAGCAG	TTGCCGCTACTTTCCTGTCT	55	350
<i>APOLI</i> EX2	ATGGGGGCAACAGAATTTT	GGGAGGTGAATTTCCAGAG	58	542
<i>APOLI</i> EX3	AGACAGGTGGTCACGAGGTC	AATTCGGGAAGTGACTTTG	58	497
<i>APOLI</i> EX4/5	CCCTGGTCATTGTCAGAAC	GTTCTTCTGGGGCTCACTCA	55	531
<i>APOLI</i> EX6	ACATCACAGCTGTCCAGGAA	GCCAAGAAGGTCATCCTCAA	58	366
<i>APOLI</i> EX7A	GGTCTCAATCTCCTGACCTTGT	TCACCCAAAACTCCCTCAC	62	589
<i>APOLI</i> EX7B	GGTGGACACAAGCCCAAG	AAAAAGTTTGCATTTTGTCTG	58	564
<i>APOLI</i> EX7C	AGCCACCAGGAGAGATATGC	TCTTCCCATTCCCCACACT	55	542
<i>APOLI</i> EX7D	ATGGGAAGTGGAGAGTGTGG	TCTTCGGAGGACATTGAACC	58	535
<i>APOLI</i> EX7E	AAGAAACAGCGGCTCCACTA	CCATCACCACCATTACGAGA	55	580

Buffer, 0.2 mM dNTPs, 0.064 units of Amplitaq Gold (Applied Biosystems, Carlsbad, California, USA), and 40 ng of genomic DNA.

PCR reactions were carried out using the following conditions: 95°C for 10 minutes, followed by 35 cycles of 95°C for 30 seconds, 30 seconds for annealing at temperatures indicated in Table 2, and elongation at 72°C for 1 minute. This was then followed by a final elongation step at 72°C for 10 minutes. Amplified products were purified using Exosap (Affymetrix, Santa Clara, California, USA) and then Sanger sequenced in both orientations (Applied Biosystems 3730, MSSM Core Facility).

Sequencing data was analyzed using the program Sequencher 4.9 (Gene Codes Corp, Ann Arbor, Michigan, USA). Control DNA, obtained from an individual without evidence of disease, was also amplified and sequenced. Patient DNA was compared to a reference sequence

NG_023228 and the cDNA sequences: NM_145343.2, NM_003661.3, NM_001136540.1, and NM_001136541.1. Sequencing variants were compared to dbSNP135 information tracks (UCSC Genome Browser Hg19 February 2009; <http://www.genome.ucsc.edu/>).

Results

For each MYH9RD patient, the *APOLI* gene was sequenced to include all coding and non-coding exons and their intron and exon boundaries. No novel coding mutations were identified in any of the four patient-derived samples. As shown in Figure 1 and Table 3, a number of previously known SNPs were present. These included 10 intronic SNPs and five within coding exons. The five exonic SNPs predicted three non-synonymous changes (E166K, M244I, and R271K) and were all located in exon 7. Patients 1 and 4 possessed all five of these SNPs. Patient 4 was homozygous for all five, whereas

Figure 1 *APOLI* single nucleotide polymorphisms SNPs.

Notes: This is the *APOLI* transcript variant 2, NM_145343.2. Boxes represent exons, connecting lines represent introns. Translation start site is depicted by "ATG" and translational stop site is depicted by "Stop". The SNP designations shown in red are exonic; those in black are intronic.

Table 3 Single nucleotide polymorphisms (SNPs) in *APOL1*

SNP	Reference sequence	Patient 1 sequence	Patient 2 sequence	Patient 3 sequence	Patient 4 sequence
c.-125-82C>T rs13056427	C	C	C/T	C/T	C/T
c.-43C>T rs6000220	C	C/T	C	C	C
c.147-90G>A rs4820224	G	G/A	G	G	G/A
c.236-14T>G rs136163	T	G	G/T	G/T	G
c.362+21C>T rs136164	C	C/T	C/T	C/T	C
c.363-48G>A rs76406801	A	G	A	A	G
c.363-45G>A rs28480494	G	A	G	G	A
c.496G>A rs2239785 E166K	G	G/A	G	G	A
c.702C>A rs136174 A235A	C	A	C	C	A
c.732G>A rs136175 M244I	G	A	G	G	A
c.812G>A rs136176 R271K	G	A	G	G/A	A
c.1008G>A rs136177 R336R	G	G/A	G/A	G/A	A
c.*658G>C rs151210481	G	G	G/C	G/C	G
c.*888delA rs66950669	A	Deletion A	Deletion A	Deletion A	Deletion A
c.*1169A>G rs78523	A	G	A/G	A/G	G

Notes: All numbering is based on *APOL1* transcript variant 2, NM_145343.2; c. = coding DNA reference sequence; * = after translational stop.

Patient 1 was homozygous for three of them (rs136174, rs136175, and rs136176). Patient 3 was heterozygous for the fourth SNP, rs136176. All patients, except for Patient 4 who was homozygous for the SNP, were heterozygous for the fifth exonic SNP, rs136177.

Discussion

MYH9RD represents a group of five autosomal dominant macrothrombocytopenias, which historically have been distinguished based on the constellation of associated clinical features beyond their commonly shared platelet defects. Gene discovery efforts, which first linked May-Hegglin Anomaly and Fechtner syndrome to the same region of chromosome 22q11-13,^{29,30} originally suggested a shared genetic etiology between these two disorders. Gene discovery then

definitively demonstrated that all five clinical entities were, in fact, one allelic disorder.^{2,31} However, despite the seemingly clear understanding of the gene mutation(s) that causes MYH9RD, the correlation between genotype and phenotype remains unclear. Without this information, providing families and individuals with guidance regarding the natural history of their specific disease and/or designing future therapeutic strategies becomes difficult, if not impossible.

It should be noted that while we uncovered no evidence to support a second mutation influencing the origin of renal disease in this syndrome, a limitation of this study was the small number of patients analyzed and the fact that only one of these, Patient 1, was of African descent. Previous studies, which examined *APOL1* mutations with regard to kidney disease, have been focused on individuals of

African descent.^{26,32,33} Interestingly, Patient 1 in our study had the greatest frequency of SNPs.

Viewing this issue of a “simple” Mendelian trait as a possibly (more) complex disorder, one could also consider the possibility of digenic versus monogenic inheritance. This raises the question of whether the mutation of a second gene, especially an adjacent gene now also linked to a renal phenotype, results in or modifies the renal disease of MYH9RD. Examples of genetic disorders with a suggested digenic inheritance pattern are becoming more frequent and this increase may be a reflection of evolving sequencing technologies capable of sequencing entire genomes and exomes (reviewed by Vockley³⁴).

Conclusion

Based on our original hypothesis, we sequenced the *APOL1* gene in four unrelated individuals with MYH9RD who all had evidence of kidney disease. The *MYH9/APOL1* locus has generated a considerable interest given its association with both a rare monogenetic trait and a common kidney disease (reviewed by Rosset et al³⁵). While our analysis does not provide any evidence to support the requirement for digenic inheritance in MYH9RD renal disorders, the results are consistent with the concept that even “simple” Mendelian disorders can represent complex traits and – as was noted more than a decade ago – for many of these traits, “only a subset of all mutations reliably predicts phenotypes.”³⁶

Disclosure

The authors report no conflicts of interest in this work.

References

- Althaus K, Greinacher A. MYH9-related platelet disorders. *Semin Thromb Hemost*. 2009;35(2):189–203.
- Seri M, Cusano R, Gangarossa S, et al. Mutations in MYH9 result in the May-Hegglin anomaly, and Fechtner and Sebastian syndromes. The May-Hegglin/Fechtner Syndrome Consortium. *Nat Genet*. 2000;26(1):103–105.
- Mhawech P, Saleem A. Inherited giant platelet disorders. Classification and literature review. *Am J Clin Pathol*. 2000;113(2):176–190.
- Kunishima S, Saito H. Advances in the understanding of MYH9 disorders. *Curr Opin Hematol*. 2010;17(5):405–410.
- Seri M, Savino M, Bordo D, et al. Epstein syndrome: another renal disorder with mutations in the nonmuscle myosin heavy chain 9 gene. *Hum Genet*. 2002;110(2):182–186.
- Peterson LC, Rao KV, Crosson JT, White JG. Fechtner syndrome – a variant of Alport’s syndrome with leukocyte inclusions and macrothrombocytopenia. *Blood*. 1985;65(2):397–406.
- Kao WH, Klag MJ, Meoni LA, et al. MYH9 is associated with non-diabetic end-stage renal disease in African Americans. *Nat Genet*. 2008;40(10):1185–1192.
- Kopp JB, Smith MW, Nelson GW, et al. MYH9 is a major-effect risk gene for focal segmental glomerulosclerosis. *Nat Genet*. 2008;40(10):1175–1184.
- Freedman BI, Hicks PJ, Bostrom MA, et al. Polymorphisms in the non-muscle myosin heavy chain 9 gene (MYH9) are strongly associated with end-stage renal disease historically attributed to hypertension in African Americans. *Kidney Int*. 2009;75(7):736–745.
- Freedman BI, Hicks PJ, Bostrom MA, et al. Non-muscle myosin heavy chain 9 gene MYH9 associations in African Americans with clinically diagnosed type 2 diabetes mellitus-associated ESRD. *Nephrol Dial Transplant*. 2009;24(11):3366–3371.
- Simons M, Wang M, McBride OW, et al. Human nonmuscle myosin heavy chains are encoded by two genes located on different chromosomes. *Circ Res*. 1991;69(2):530–539.
- Toothaker LE, Gonzalez DA, Tung N, et al. Cellular myosin heavy chain in human leukocytes: isolation of 5’ cDNA clones, characterization of the protein, chromosomal localization, and upregulation during myeloid differentiation. *Blood*. 1991;78(7):1826–1833.
- Lalwani AK, Goldstein JA, Kelley MJ, Luxford W, Castelein CM, Mhatre AN. Human nonsyndromic hereditary deafness DFNA17 is due to a mutation in nonmuscle myosin MYH9. *Am J Hum Genet*. 2000;67(5):1121–1128.
- Harrington WF, Rodgers ME. Myosin. *Annu Rev Biochem*. 1984;53:35–73.
- Pecci A, Panza E, Pujol-Moix N, et al. Position of nonmuscle myosin heavy chain IIA (NMMHC-IIA) mutations predicts the natural history of MYH9-related disease. *Hum Mutat*. 2008;29(3):409–417.
- Sekine T, Konno M, Sasaki S, et al. Patients with Epstein-Fechtner syndromes owing to MYH9 R702 mutations develop progressive proteinuric renal disease. *Kidney Int*. 2010;78(2):207–214.
- Nelson GW, Freedman BI, Bowden DW, et al. Dense mapping of MYH9 localizes the strongest kidney disease associations to the region of introns 13 to 15. *Hum Mol Genet*. 2010;19(9):1805–1815.
- Winkler CA, Nelson G, Oleksyk TK, Nava MB, Kopp JB. Genetics of focal segmental glomerulosclerosis and human immunodeficiency virus-associated collapsing glomerulopathy: the role of MYH9 genetic variation. *Semin Nephrol*. 2010;30(2):111–125.
- Genovese G, Tonna SJ, Knob AU, et al. A risk allele for focal segmental glomerulosclerosis in African Americans is located within a region containing *APOL1* and *MYH9*. *Kidney Int*. 2010;78(7):698–704.
- Duchateau PN, Pullinger CR, Orellana RE, et al. Apolipoprotein L, a new human high density lipoprotein apolipoprotein expressed by the pancreas. Identification, cloning, characterization, and plasma distribution of apolipoprotein L. *J Biol Chem*. 1997;272(41):25576–25582.
- Duchateau PN, Pullinger CR, Cho MH, Eng C, Kane JP. Apolipoprotein L gene family: tissue-specific expression, splicing, promoter regions; discovery of a new gene. *J Lipid Res*. 2001;42(4):620–630.
- Wan G, Zhaorigetu S, Liu Z, Kaini R, Jiang Z, Hu CA. Apolipoprotein L1, a novel Bcl-2 homology domain 3-only lipid-binding protein, induces autophagic cell death. *J Biol Chem*. 2008;283(31):21540–21549.
- Vanhamme L, Paturiaux-Hanocq F, Poelvoorde P, et al. Apolipoprotein L-I is the trypanosome lytic factor of human serum. *Nature*. 2003;422(6927):83–87.
- Monajemi H, Fontijn RD, Pannekoek H, Horrevoets AJ. The apolipoprotein L gene cluster has emerged recently in evolution and is expressed in human vascular tissue. *Genomics*. 2002;79(4):539–546.
- Vanhollebeke B, Truc P, Poelvoorde P, et al. Human Trypanosoma evansi infection linked to a lack of apolipoprotein L-I. *N Engl J Med*. 2006;355(26):2752–2756.
- Freedman BI, Kopp JB, Langefeld CD, et al. The apolipoprotein L1 (*APOL1*) gene and nondiabetic nephropathy in African Americans. *J Am Soc Nephrol*. 2010;21(9):1422–1426.
- Nozu K, Inagaki T, Fu XJ, et al. Molecular analysis of digenic inheritance in Bartter syndrome with sensorineural deafness. *J Med Genet*. 2008;45(3):182–186.
- Hoefele J, Wolf MT, O’Toole JF, et al. Evidence of oligogenic inheritance in nephronophthisis. *J Am Soc Nephrol*. 2007;18(10):2789–2795.
- Martignetti JA, Heath KE, Harris J, et al. The gene for May-Hegglin anomaly localizes to a <1-Mb region on chromosome 22q12.3-13.1. *Am J Hum Genet*. 2000;66(4):1449–1454.

30. Toren A, Rozenfeld-Granot G, Rocca B, et al. Autosomal-dominant giant platelet syndromes: a hint of the same genetic defect as in Fechtner syndrome owing to a similar genetic linkage to chromosome 22q11-13. *Blood*. 2000;96(10):3447–3451.
31. Kelley MJ, Jawien W, Ortel TL, Korczak JF. Mutation of MYH9, encoding non-muscle myosin heavy chain A, in May-Hegglin anomaly. *Nat Genet*. 2000;26(1):106–108.
32. Genovese G, Friedman DJ, Ross MD, et al. Association of trypanolytic ApoL1 variants with kidney disease in African Americans. *Science*. 2010;329(5993):841–845.
33. Tzur S, Rosset S, Shemer R, et al. Missense mutations in the APOL1 gene are highly associated with end stage kidney disease risk previously attributed to the MYH9 gene. *Hum Genet*. 2010;128(3):345–350.
34. Vockley, Jerry (2011) Digenic Inheritance. In: Encyclopedia of Life Sciences (ELS). John Wiley & Sons, Ltd: Chichester. DOI: 10.1002/9780470015902.a0005560.pub2
35. Rosset S, Tzur S, Behar DM, Wasser WG, Skorecki K. The population genetics of chronic kidney disease: insights from the MYH9-APOL1 locus. *Nat Rev Nephrol*. 2011;7(6):313–326.
36. Dipple KM, McCabe ER. Phenotypes of patients with “simple” Mendelian disorders are complex traits: thresholds, modifiers, and systems dynamics. *Am J Hum Genet*. 2000;66(6):1729–1735.

Advances in Genomics and Genetics

Publish your work in this journal

Advances in Genomics and Genetics is an international, peer reviewed, open access journal that focuses on new developments in characterizing the human and animal genome and specific gene expressions in health and disease. Particular emphasis will be given to those studies that elucidate genes, biomarkers and targets in the development of new or improved therapeutic

interventions. The journal is characterized by the rapid reporting of reviews, original research, methodologies, technologies and analytics in this subject area. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/advances-in-genomics-and-gene-expression-journal>

Dovepress