Incorporation of a selective sigma-2 receptor ligand enhances uptake of liposomes by multiple cancer cells

Yifei Zhang^{1,*}
Yixian Huang^{1,*}
Peng Zhang¹
Xiang Gao¹
Robert B Gibbs²
Song Li¹

¹Center for Pharmacogenetics,

²Department of Pharmaceutical Sciences, University of Pittsburgh School of Pharmacy, Pittsburgh, PA, USA,

*These authors contributed equally to this work

Background: The sigma-2 receptor is an attractive target for tumor imaging and targeted therapy because it is overexpressed in multiple types of solid tumors, including prostate cancer, breast cancer, and lung cancer. SV119 is a synthetic small molecule that binds to sigma-2 receptors with high affinity and specificity. This study investigates the utility of SV119 in mediating the selective targeting of liposomal vectors in various types of cancer cells.

Methods: SV119 was covalently linked with polyethylene glycol-dioleoyl amido aspartic acid conjugate (PEG-DOA) to generate a novel functional lipid, SV119-PEG-DOA. This lipid was utilized for the preparation of targeted liposomes to enhance their uptake by cancer cells. Liposomes with various SV119 densities (0, 1, 3, and 5 mole%) were prepared and their cellular uptake was investigated in several tumor cell lines. In addition, doxorubicin (DOX) was loaded into the targeted and unmodified liposomes, and the cytotoxic effect on the DU-145 cells was evaluated by MTT assay.

Results: Liposomes with or without SV119-PEG-DOA both have a mean diameter of approximately 90 nm and a neutral charge. The incorporation of SV119-PEG-DOA significantly increased the cellular uptake of liposomes by the DU-145, PC-3, A549, 201T, and MCF-7 tumor cells, which was shown by fluorescence microscopy and the quantitative measurement of fluorescence intensity. In contrast, the incorporation of SV119 did not increase the uptake of liposomes by the normal BEAS-2B cells. In a time course study, the uptake of SV119 liposomes by DU-145 cells was also significantly higher at each time point compared to the unmodified liposomes. Furthermore, the DOX-loaded SV119 liposomes showed significantly higher cytotoxicity to DU-145 cells compared to the DOX-loaded unmodified liposomes.

Conclusion: SV119 liposomes were developed for targeted drug delivery to cancer cells. The targeting efficiency and specificity of SV119 liposomes to cancer cells was demonstrated in vitro. The results of this study suggest that SV119-modified liposomes might be a promising drug carrier for tumor-targeted delivery.

Keywords: SV119, targeted delivery, liposomes, cellular uptake, cancer therapy

Introduction

One major problem with conventional anticancer agents is their severe side effects on normal tissues and organs. To solve this issue, targeted drug delivery to solid tumors was developed as an effective strategy to improve the therapeutic effect and minimize toxicity. Many factors affect the success rate of targeted drug delivery, including the choice of targeting ligands. Many types of ligands have been explored, including antibodies, peptides, aptamers, and small molecule ligands.¹ Small molecule ligands have the advantages of minimal immunogenicity, excellent chemical stability, and ease of synthesis.² Some examples of small molecule ligands that have been developed for

Correspondence: Song Li
639 Salk Hall, Center for Pharmacogenetics, School of Pharmacy, University of Pittsburgh, Pittsburgh, PA 15261, USA
Tel +1 412 383 7976
Fax +1 412 648 1664
Email sol4@pitt.edu

tumor targeting include ligands of the folate receptor,³ prostate-specific membrane antigen (PSMA),⁴ and integrin receptors.⁵ There has been a recent interest in exploring sigma receptor ligands for tumor imaging and targeted therapy. Sigma receptors are overexpressed in a variety of human tumors including non-small cell lung carcinoma (NSCLC), prostate cancer, melanoma, and breast cancer.^{6–8} The small molecule ligands of sigma receptors are considered potential tumor imaging agents.^{6–8} The authors of this study and Dr Huang's group have reported that sigma receptor ligands, such as anisamide, effectively mediate the selective delivery of liposomes containing doxorubicin (DOX) to prostate cancer cells in vitro and in vivo.^{9,10} Extensive studies by Dr Huang's group have demonstrated the targeted delivery of oligodeoxynucleotides and siRNA to lung cancer and melanoma using anisamide-decorated nanoparticles.^{11–13} Another sigma receptor ligand, haloperidol, was shown to improve the transfection of MCF-7 breast carcinoma cells by more than 10-fold, compared with the control (unmodified) liposomes.¹⁴

Neither anisamide nor haloperidol distinguishes between the sigma-1 or sigma-2 receptor subtypes with respect to binding. This study determines whether sigma 2-selective ligands can be employed for the targeted delivery of liposomal vectors. There are several potential advantages associated with sigma 2-selective ligands for tumor-targeted delivery. First, there is a higher density of sigma-2 versus sigma-1 receptors in tumor samples and cultured tumor cells.^{15–17} Second, ligand binding and photoaffinity labeling studies have demonstrated a lower expression of sigma-2 versus sigma-1 receptors in normal tissues from the central nervous system, gastrointestinal tract, kidney, liver, and heart.¹⁸ Third, a correlation between sigma-2 receptor expression and tumor proliferation has been established both in vitro and in vivo,^{19,20} but no such correlation was observed for the sigma-1 subtype.

SV119 is a small molecule ligand that binds to sigma-2 receptors with high affinity and specificity.²¹ The tumor-targeting efficiency of SV119 has been demonstrated by conjugating SV119 with apoptosis-inducing peptides, which resulted in the selective killing of tumor cells.²² The present study explores the potential of SV119 to mediate the targeted delivery of liposomes to multiple tumor cell lines. Liposomes are well-established drug carriers for intravenous delivery and several liposomal formulations have been approved for clinical applications.²³ Our results show that the incorporation of a SV119 derivatized PEG-lipid significantly increases the cellular uptake of liposomes by several tumor cell lines, but not by normal cells. In addition, DOX-loaded SV119-modified liposomes showed significantly higher cytotoxicity compared to unmodified liposomes loaded with DOX. These findings suggest that SV119 liposomes may

have potential for the targeted delivery of anticancer agents to tumors that over-express sigma-2 receptors.

Materials and methods

Materials

L- α -phosphatidylcholine (Soy PC), DSPE-PEG(2000)-OCH₃, and 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(lissamine rhodamine B sulfonyl) (ammonium salt) (rhodamine-PE) were purchased from Avanti® Polar Lipids, Inc (Alabaster, AL). Cholesterol (Chol), doxorubicin hydrochloride, FITC-conjugated anti-mouse IgG, 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT), and dimethyl sulfoxide (DMSO) were purchased from Sigma-Aldrich (St Louis, MO, USA). Mouse anti-early endosome antigen 1 (EEA1) monoclonal antibody was purchased from BD Transduction Laboratories. LysoTracker® Green DND-26 was purchased from Life Technologies (Invitrogen®; Carlsbad, CA). A PD-10 desalting column was purchased from GE Healthcare (Little Chalfont, UK). Dulbecco's phosphate-buffered saline (DPBS) was purchased from Life Technologies (Gibco®) and t-Boc Amine PEG NHS Ester (TBOC-PEG-NHS, MW 3500) was purchased from JenKem Technology USA, Inc (Allen, TX). All other chemicals and organic solvents were purchased from Sigma-Aldrich. ¹H nuclear magnetic resonance (NMR) spectra were recorded on a Bruker FT 400 MHz. The synthesized compounds were also identified by matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry (Voyager-DE PRO BioSpectrometry Workstation; Applied Biosystems®, Life Technologies). The mass spectrum was obtained in a linear mode with CHCA as the matrix.

Human prostate adenocarcinoma cell line DU-145, human lung carcinoma cell line A549, and human breast cancer MCF-7 cells were obtained from American Type Culture Collection ([ATCC], Manassas, VA) Rockville, MD. BEAS-2B (human bronchial epithelium cell line) was generously gifted by Dr Y Peter Di from the University of Pittsburgh. The cells were cultured in Dulbecco's modified Eagle's medium (DMEM; ATCC) and supplemented with 10% FBS (Invitrogen®) and 1% penicillin-streptomycin at 37°C in a humidified atmosphere of 5% CO₂ in air. All the reagents were obtained from Sigma-Aldrich unless stated otherwise.

Synthesis and chemical characterization of SV119-PEG-DOA conjugates and intermediates

Figure 1 schematically depicts the synthetic procedure for preparing SV119-PEG-DOA. The experimental details are delineated below.

Compound 2

NaH (0.8 g, 60% in oil, 20 mmol) was added to a solution of 1 (3.04 g, 10 mmol) and benzyl 2-bromoacetate (2.75 g, 12 mmol) in THF (50 mL).²⁴ The mixture was stirred overnight at an ambient temperature. After ether (75 mL) was added, the organic solution was quenched with water (30 mL) and then washed with a saturated NaCl solution (40 mL), and finally, the organic solution was dried over Na₂SO₄. After the solvent was removed, the crude product was purified by flash column chromatography with hexane-ethyl acetate (1:1) to obtain compound 2 (4.33 g, 96%) as a pale yellow oil. ¹H NMR (CDCl₃) δ 7.98 (s, 1H), 7.37–7.21 (m, 6H), 6.77 (m, 2H), 5.17 (s, 2H), 5.00 (m, 1H), 3.83 (s, 3H), 3.42 (m, 2H), 3.27 (m, 2H), 2.33 (m, 2H), 2.30 (s, 3H), 2.13 (m, 1H), 1.98 (m, 2H), 1.51 (m, 3H), 1.17 (m, 2H). ESI-MS *m/z* 453.5 ([M+H]⁺).

Compound 3

Chemical name: 2-(3-((2-methoxy-5-methylphenyl) carbamoyl)-9-aza-bicyclo[3.3.1]nonan-9-yl)acetic acid. Pd/C (10%, 500 mg) was added to a solution of 2 (3.62 g, 10 mmol) and stirred under atmospheric H₂ pressure in MeOH (50 mL). EtOAc (100 mL) was added after the solvents evaporated under a vacuum. The mixture was washed by a saturated NaCl solution (40 mL), and finally, the organic solution was

dried over Na₂SO₄. EtOAc was then removed to obtain compound 3 (2.82 g, 100%) as a white solid. ¹H NMR (CDCl₃) δ 10.9 (brs, 1H), 7.96 (s, 1H), 7.15 (s, 1H), 6.79 (m, 2H), 4.96 (m, 1H), 3.83 (s, 3H), 3.32 (m, 2H), 3.04 (m, 2H), 2.35 (m, 2H), 2.29 (s, 3H), 2.08 (m, 1H), 1.64 (m, 3H), 1.50 (m, 4H). ESI-MS *m/z* 363.4 ([M+H]⁺).

Compound 4

A mixture of Boc-aspartic acid (2.47 g, 10 mmol), oleyl amine (8.01 g, 30 mmol), DCC (6.18 g, 30 mmol), and DMAP (122 mg, 1 mmol) in anhydrous CH₂Cl₂ (60 mL) was stirred vigorously at room temperature for 5 hours. After the reaction was completed, the mixture was filtered and evaporated under reduced pressure, and the residue was purified by flash column chromatography (MeOH: CHCl₃ 1:9) to afford the product (4.40 g, 60%). ¹H NMR (CDCl₃) δ 7.93 (m, 1H), 5.37 (m, 4H), 4.17 (m, 1H), 3.24 (m, 4H), 1.98 (m, 9H), 1.51 (m, 4H), 1.47 (s, 9H), 1.26 (m, 45H), 0.87 (m, 6H). ESI-MS *m/z* 719.1 ([M+H]⁺).

DOA (dioleoyl amido aspartic acid)

Trifluoroacetic acid (6 mL) was added to a solution of 4 (4.4 g, 6 mmol) in anhydrous CH₂Cl₂ (6 mL). After stirring for 2 hours, the reaction mixture was diluted with CH₂Cl₂ (50 mL) and a saturated aqueous solution of Na₂CO₃ (20 mL)

Figure 1 Synthesis scheme for SVI19-PEG3500-DOA.

Abbreviations: DOA, dioleoyl amido aspartic acid; PEG, polyethylene glycol.

was added until the gas evolution ceased. The organic layer was then separated and washed with brine. Next, the organic layers were dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography ($\text{MeOH}:\text{CHCl}_3$ 1:9) to afford the product (3.70 g, 95%). ^1H NMR (CDCl_3) δ 7.53 (m, 1H), 6.33 (m, 1H), 5.37 (m, 4H), 3.67 (m, 1H), 3.22 (m, 4H), 1.98 (m, 9H), 1.50 (m, 4H), 1.26 (m, 45H), 0.87 (m, 6H). ESI-MS m/z 641.0 ($[\text{M}+\text{Na}]^+$).

Compound 6

A mixture of Boc- NH_2 -PEG-NHS (350 mg, 0.1 mmol) and DOA (63.1 mg, 0.1 mmol) in anhydrous CH_2Cl_2 (1 mL) was stirred vigorously at room temperature for 2 hours. The solution was precipitated in cold diethyl ether and the precipitate was washed again by diethyl ether. Anhydrous CH_2Cl_2 (1 mL) and TFA (1 mL) were added to the precipitate and the mixture was stirred for 2 hours. The reaction mixture was diluted with CH_2Cl_2 (10 mL) and a saturated aqueous solution of Na_2CO_3 (2 mL) was then added until the gas evolution ceased. The organic layer was separated and precipitated in cold diethyl ether. The obtained precipitate was washed again by diethyl ether.

Compound 7

A mixture of compound 3 (72 mg, 0.2 mmol), PEG-DOA (300 mg, 0.1 mmol), DCC (41.2 mg, 0.2 mmol), and DMAP (12 mg, 0.1 mmol) in anhydrous CH_2Cl_2 (1 mL) was stirred vigorously at room temperature for 5 hours. The solution was precipitated in cold diethyl ether and washed again with diethyl ether to obtain compound 7 (245 mg, 73%).

Compound 8

A mixture of compound 1 (304 mg, 1 mmol), succinic anhydride (200 mg, 2 mmol), and DMAP (12 mg, 0.1 mmol) in anhydrous CH_2Cl_2 (1 mL) was stirred vigorously at room temperature for 5 hours. The residue was purified by flash column chromatography (ethyl acetate) to afford the product (3.70 g, 95%). ^1H NMR (CDCl_3) δ 10.9 (brs, 1H), 7.96 (s, 1H), 7.15 (s, 1H), 6.79 (m, 2H), 4.96 (m, 1H), 3.83 (s, 3H), 3.04 (m, 2H), 2.72 (m, 2H), 2.35 (m, 2H), 2.29 (s, 3H), 2.24 (m, 2H), 2.08 (m, 1H), 1.64 (m, 3H), 1.50 (m, 4H).

Compound 9

A mixture of compound 8 (75 mg, 0.2 mmol), PEG-DOA (300 mg, 0.1 mmol), DCC (41.2 mg, 0.2 mmol), and DMAP (12 mg, 0.1 mmol) in anhydrous CH_2Cl_2 (1 mL) was stirred vigorously at room temperature for 5 hours. The solution was precipitated in cold diethyl ether and washed again with diethyl ether to obtain compound 9 (212 mg, 65%).

Preparation of rhodamine-PE labeled liposomes

Rhodamine-PE labeled liposomes with or without SV119 were prepared by thin lipid film hydration followed by probe sonication. Unmodified liposomes were composed of SPC:Chol:DSPE-PEG2000:rhodamine-PE in a 7:3:0.5:0.05 molar ratio. A chloroform solution of the lipid components was mixed and evaporated under a gentle stream of N_2 followed by vacuum for at least 4 hours. The dried lipid films were hydrated with DPBS at 4°C overnight with an initial total lipid concentration of 16 mM. After a brief vortex, the suspension was then probe sonicated for 1 hour at a power of 3 watts. The liposomes were then passed through a PD-10 desalting column and used for fluorescence microscopic and quantitative liposome uptake studies. The preparation of the SV119 liposomes followed the same procedures, except the DSPE-PEG2000 was partially substituted by a DOA-PEG(3500)-SV119 conjugate. The size distribution and ζ -potentials of the liposomes were monitored with a dynamic light scattering method using the Malvern Zetasizer (Nano-ZS; Malvern Instruments Ltd, Worcestershire, UK).

Cellular uptake studies

Cells (2×10^4 per well) were seeded onto 48-well plates (Corning Inc, Corning, NY) and allowed to attach overnight. The complete medium was replaced with the serum-free DMEM containing rhodamine-PE labeled liposomes at a final total lipid concentration of 800 μM . Unmodified or SV119 liposomes were incubated with cells at 37°C for 1 hour unless otherwise indicated. After incubation, the cells were washed three times with ice cold PBS and lysed with 200 μl 0.3% Triton X-100 at room temperature for 0.5 hours. The fluorescence intensity of the cell lysate was determined by a PerkinElmer LS 50B luminescence spectrometer (Waltham, MA) ($\text{ex} = 560 \text{ nm}$; $\text{em} = 588 \text{ nm}$). The results were expressed as arbitrary fluorescence intensity/mg of protein lysates. All the numbers were subtracted by the autofluorescence of the cells incubated in the serum-free medium without any formulations. The results are reported as the average values of four wells measured from the same plate on the same day.

To evaluate the cellular uptake by microscopy, the cells were fixed with 4% paraformaldehyde in DPBS at 4°C for 20 minutes and washed three times by DPBS. The fluorescence images were obtained using a Nikon fluorescence-phase contrast-optical microscope.

To investigate the effect of free SV119 on the uptake of SV119 liposomes, SV119 was conjugated with PEG550 to increase its water solubility (Figure S1A). The effect of

SV119 on the uptake of SV119 liposomes was examined by incubating the DU-145 cells with the liposomes in serum-free media for 1 hour at 37°C in the presence of SV119-PEG550 (200 μ M) or PEG550 (200 μ M), and a fluorescence microscopy study was carried out as described above.

Preparation of DOX-loaded liposomes

DOX was encapsulated into unmodified and SV119 liposomes using an ammonium sulphate gradient method as previously described.²⁵ The components of the blank liposomes were similar to the liposomes described in the section titled “Preparation of rhodamine-PE labeled liposomes.” The percentage of SV119-PEG3500-DOA in the total lipid of targeted liposomes was 5 mole%. Instead of DPBS, 250 mM ammonium sulfate solution was used to hydrate the lipid membrane. DOX (10 mg/mL) dissolved in 5% glucose was mixed with the liposomes and incubated at 37°C for 20 minutes. The drug/lipid ratio was kept at approximately 100 μ g/ μ mol to achieve an encapsulation efficiency greater than 90%.²⁵ Nonentrapped DOX was removed by filtration through a PD-10 column. After the solubilization of the liposomes in 1% (w/v) Triton X-100, the DOX concentration of the liposomes was determined by measuring the absorbance at 480 nm using free DOX solution as a standard.

Cytotoxicity assay

The viability of cells treated by DOX encapsulated in SV119 liposomes or unmodified liposomes was measured by MTT colorimetric assay. DU-145 cells were trypsinized and resuspended in serum-free DMEM. The formulations were diluted in a serum-free medium and mixed with DU-145 cell suspension, followed by incubation at 37°C for 1 hour. The medium was then removed by centrifugation. The cell pellets were resuspended in a complete medium and plated on a 96-well plate with a density of 5×10^3 cells per well. The cells were cultured at 37°C for another 72 hours followed by an MTT assay. Twenty microliters of MTT solution (5 mg/mL in PBS) were added to each well and the cells were incubated for 4 hours at 37°C. The media were then removed and 150 μ L DMSO was added to each well. Absorbance at 550/630 nm was measured using an Ultramark Microplate Imaging System (Bio-Rad Laboratories, Hercules, CA). Six wells were used for each sample, and the data were expressed as the percentage of viable cells compared to the untreated control cells.

Intracellular localization

DU-145 cells grown to 60%–70% confluence in glass bottom dishes (In Vitro Scientific, Sunnyvale, CA) were incubated with rhodamine-PE labeled liposomes (with

3 mole% SV119-PEG-DOA) at 37°C for 3 hours or 16 hours. To detect early endosomes, the cells were fixed with 4% paraformaldehyde, permeabilized with 0.2% saponin, and stained with anti-EEA1 (1:100), followed by FITC-conjugated anti-mouse IgG (1:100). To label the lysosomes in live cells, LysoTracker Green® (50 nM; Life Technologies) was added to the medium 30 minutes before the incubation period concluded. The stained cells were rapidly washed with ice-cold PBS and observed using an Olympus FV1000 laser scanning confocal microscope (Olympus America Inc, Melville, NY).

Statistical analysis

Statistical analysis was conducted using a two-tailed unpaired Student's *t*-test with the SPSS software (version 11.5; IBM Corporation, Armonk, NY). The data were expressed as mean $\pm$ SD and differences of *P* < 0.05 were considered significant.

Results and discussion

Several selective sigma-2 receptor ligands have recently been developed and evaluated in vitro and in vivo. For example, Mach and Wheeler reported the synthesis of a series of *N*-substituted 9-azabicyclo[3.3.1]nonan-3a-yl phenylcarbamate analogs.²⁶ Several high-affinity sigma-2-selective ligands such as SV119 were identified through structure-activity relationship (SAR) studies. SV119 has a free amino group that is linked to a bridgehead nitrogen atom via a linker group of six methylene units. SAR studies suggest that the amino group is a preferred substituent for assuring a high affinity for σ_2 receptors and high $\sigma_1 : \sigma_2$ selectivity. Nevertheless, various SV119 conjugates were synthesized by chemically modifying this amino group. For example, coupling SV119 with a fluorescence molecule via this amino group led to the development of a sigma-2-selective probe – SW-120 – which was used to effectively assess the sigma-2 receptor expression levels in cultured tumor cells.²⁷ In addition, a recent study by Spitzer et al showed that coupling SV119 with a proapoptotic peptide via this amino group similarly led to the development of a conjugate that demonstrated selective cytotoxicity towards tumor cells that overexpress the sigma-2 receptor.²⁸

In order to develop lipid-based nanoparticles that incorporate SV119 on the surface, we designed and synthesized a SV119-derivatized PEG-lipid. DOA is a double-chain lipid that is synthesized from oleyl amine and aspartic acid and can be inserted into the lipid bilayer of liposomes. The PEG moiety on liposomes provides steric hindrance, which renders the nanoparticles more stable and minimizes nonspecific binding of the liposomes on the cell surface.²⁹ After replacing the

amine group with a carboxyl group, a SV119-PEG-DOA conjugate was synthesized by covalent linkage between the SV119 carboxyl group and the PEG terminal amine group. Figure 1 illustrates the synthesis scheme of SV119-PEG3500-DOA.

Figure 2A shows the NMR spectrum of synthesized SV119-PEG3500-DOA. The peaks at 7.2–7.3 ppm are attributed to the benzene groups on SV119, the signals at 5.0–5.2 ppm are attributed to the double bond of DOA, the peaks between 3–4 are attributed to the PEG, and the signals at 1–2 ppm are attributed to the carbon chain of DOA. The molecular weight of the synthesized product was determined by MALDI-TOF mass spectrometry. As

shown in Figure 2B, the spectra exhibited a bell-shaped distribution of 44 Da-spaced lines centered at 4754.9 Da for SV119-PEG3500-DOA. A comparison between the mass spectrum of SV119-PEG3500-DOA (Figure 2B) with PEG3500 (Figure S2) and PEG3500-DOA (Figure S3) suggests the successful synthesis of the SV119-PEG3500-DOA conjugate.

The synthesized SV119-PEG-DOA was used to formulate targeted liposomes. Liposomes are mainly composed of phospholipids and cholesterol. These molecules are also components of the cells and, thus, are believed to be biocompatible and non-toxic. The present experiment demonstrates that all the liposomal formulations have a mean

Figure 2 NMR spectrum (A) and MALDI-TOF mass spectrum (B) of SV119-PEG3500-DOA. The peaks in (A) are attributed to the benzene groups on SV119 (a), the double bonds of DOA (b), and the carbon chains of DOA (c).

Abbreviations: DOA, dioleoyl amido aspartic acid; MALDI-TOF, matrix-assisted laser desorption/ionisation-time of flight; NMR, nuclear magnetic resonance; PEG, polyethylene glycol.

diameter of approximately 90 nm (Table 1), which is optimal for penetrating the blood vessels surrounding the tumor and subsequent endocytosis by the tumor cells.³⁰

The targeting efficiency of SV119 liposomes was characterized *in vitro*. Cellular uptake was studied using DU-145 cells as target cells. DU-145 is a human prostate cancer cell line and these cells are reported to overexpress sigma-2 receptors.⁸ Human bronchial epithelial cells – BEAS-2B – were chosen to examine the liposomal uptake by normal cells. SV119-PEG3500-DOA was formulated into rhodamine-PE labeled liposomes to partially substitute PEG2000-DSPE, and the total amount of PEG-conjugated lipid in the liposomes was kept at 5 mole%. A longer spacer (PEG3500) was used between SV119 and DOA to minimize the potential steric hindrance imposed by DSPE-PEG2000 and to increase the targeting efficiency.³¹

Figure 3A shows the fluorescence images of the DU-145 cells that were treated with rhodamine-PE labeled SV119 liposomes. After incubation with SV119 liposomes or unmodified liposomes for 1 hour, fluorescence was observed on the cell membrane and inside the cells. Increasing the amount of SV119 in the liposomes was associated with a significant increase in the intensity of rhodamine fluorescence. This is in contrast to the unmodified liposomes, which showed minimal binding and cellular uptake with the DU-145 cells (Figure 3A). Figure 3B shows the quantitative measurements of the cell-associated fluorescence 1 hour after the DU145 cells were treated with the SV119 liposomes and the unmodified liposomes. Consistent with the fluorescence imaging studies (Figure 3A), the amount of cell-associated fluorescence for the DU-145 cells increased significantly as the amount of ligand in the liposomes increased. In addition, the binding and uptake of SV119 liposomes was competitively blocked by the presence of the SV119-PEG550 conjugate, but was not inhibited by the same concentration of PEG550 (Figure S1B). These results demonstrate that the enhanced uptake of SV119 liposomes was mediated by SV119.

In order to demonstrate the cell-type specificity of the uptake of SV119 liposomes, normal human bronchial epithelial cells (BEAS-2B) were tested as a control. The results showed little binding and cellular uptake of SV119

Table 1 Size and zeta potential of SV119 liposomes

SV119-PEG-DOA	0%	1%	3%	5%
Particle size (nm)	90.0 ± 0.2	90.7 ± 0.3	97.6 ± 0.2	89.2 ± 0.6
Zeta potential (mV)	-2.0 ± 0.6	-2.4 ± 0.5	-2.2 ± 0.3	-3.1 ± 0.4

Note: Data are the mean ± SD for n = 3.

Figure 3 Enhanced intracellular uptake of SV119 liposomes by DU-145 cells, examined by microscopic study (A) and quantitative analysis (B) (n = 4, mean ± SD). In (A), the percentage of SV119-PEG-DOA in the total lipid was 0% (a), 1% (b), 3% (c), and 5% (d), respectively.

Abbreviations: DOA, dioleoyl amido aspartic acid; PEG, polyethylene glycol.

liposomes and unmodified liposomes by the BEAS-2B cells (Figure 4A). No difference was found between the SV119 liposomes and unmodified liposomes in terms of the intensity of cell-associated fluorescence (Figure 4B). These observations suggest that SV119-mediated liposomal uptake is specific to tumor cells.

One of the unique features of the sigma-2 receptor is its overexpression in multiple types of cancers. Therefore, following the demonstration of the specific targeting of DU145 cells, we examined the targeting capacity of SV119 liposomes using several other cancer cell lines, including PC-3 (prostate cancer), MCF-7 (breast cancer), A549, and 201T (lung cancer) cells. Similar to the DU-145 cells, the cell-associated fluorescence intensity of the SV119 liposomes was much higher than that of the unmodified liposomes in all the tumor cell lines tested (Figure 5), which suggests a

Figure 4 Cellular uptake of liposomes by BEAS-2B cells. Microscopic images (A), and quantitative analysis (B) ($n = 4$, mean $\pm$ SD). The percentage of SV119-PEG-DOA in the total lipid was 0% (a), or 3% (b).

Abbreviations: DOA, dioleoyl amido aspartic acid; PEG, polyethylene glycol.

potential for the SV119 liposomes to target multiple types of tumors.

Next, we investigated the time course of liposomal uptake by the DU-145 cells. As shown in Figure 6, the cell-associated fluorescence intensity increased during the prolonged incubation within 6 hours for the SV119 liposomes and the unmodified liposomes. Note that the cellular uptake of the SV119 liposomes was significantly higher at each time point compared to the unmodified liposomes. This observation further supports our conclusion that the incorporation of SV119 significantly enhances the uptake of liposomes into tumor cells.

Like many other sigma-2 receptor ligands,²⁶ SV119 has a tertiary amine group. Therefore, incorporating SV119 onto the surface of liposomes may positively charge the particles, which may increase the nonspecific cellular uptake of the SV119 liposomes. However, the zeta potential analysis showed that the incorporation of up to 5 mol% of lipid-PEG-derivatized SV119 into the liposomes did not significantly alter the surface charge (Table 1), which excluded the possibility that the enhanced uptake was mediated by a

Figure 5 Uptake of SV119 liposomes by MCF-7 (A), PC-3 (B), 201T (C), and A549 cells (D) ($n = 4$, mean $\pm$ SD).

Abbreviations: DOA, dioleoyl amido aspartic acid; PEG, polyethylene glycol.

non-specific charge-charge interaction. Furthermore, to minimize the protonation of the amine nitrogen, a modified SV119 (mSV119) was synthesized by introducing an adjacent carbonyl group, and the mSV119 was similarly conjugated with PEG3500-DOA for incorporation into the liposomes (Figure 7A). This modification did not affect the targeting efficiency of the ligand (Figure 7B). This result may shed new light on the structure-activity relationship, which may aid in the design of new sigma-2 receptor ligands.

Figure 6 Time-course of cellular uptake for SV119 liposomes.

Notes: Rhodamine-PE labeled liposomes with 0% or 3% SV119-PEG-DOA were incubated with DU-145 cells for an indicated period at 37°C. ($n = 4$, mean $\pm$ SD).

Abbreviations: DOA, dioleoyl amido aspartic acid; PEG, polyethylene glycol.

Figure 7 Uptake of liposomes decorated with modified SV119 (mSV119) by DU-145 cells. Scheme of synthesis (**A**) and microscopic images (**B**) ($n = 4$, mean $\pm$ SD). In (**B**), the percentage of mSV119-PEG-DOA in the total lipid was 0% (a), 1% (b), 3% (c), and 5% (d), respectively.

Abbreviations: DOA, dioleoyl amido aspartic acid; PEG, polyethylene glycol.

After demonstrating the efficiency and specificity of SV119 in mediating the uptake of liposomes by tumor cells, we investigated the utility of the SV119 liposomes for the targeted delivery of anticancer agents. We chose DOX because it is a chemotherapeutic drug widely used in clinical therapy. The DU-145 cells were pre-treated with DOX-loaded SV119 liposomes (SV119-L[D]), DOX-loaded unmodified liposomes (L[D]) and free DOX for 1 hour. The cells were then cultured for an additional 72 hours before the MTT assay was conducted. The results show that the cytotoxic effects of all the treatments were dose-dependent at concentrations of 0.2–10 $\mu\text{g/mL}$ (Figure 8). At each of the indicated concentrations, the viability of the cells treated with L[D] showed enhancements of 1.5–3-fold, compared with that of the free DOX, which suggests that the cells were protected by the encapsulation of the drug in the liposomes. In addition, SV119-L[D] exhibited an ~2-fold higher cytotoxicity to the DU-145 cells than L[D] (Figure 8), which suggests improved efficacy of

Figure 8 Cytotoxic effect of free and liposomal DOX on DU-145 prostate cancer cells. Cells were treated with free DOX, DOX-loaded SV119 liposomes or DOX-loaded unmodified liposomes with DOX concentrations of 0.2, 1, or 10 $\mu\text{g/mL}$. Cytotoxicity was assessed by MTT assay. Cells receiving no treatment were defined as the maximal cell viability. Values presented are the mean $\pm$ SD of six replicates.

Note: * $P < 0.05$ compared to the control treatments with unmodified liposomes.

Abbreviations: DOX, doxorubicin; L[D], DOX-loaded unmodified liposomes.

the SV119-modified liposomes versus the unmodified liposomes for delivering DOX to the tumor cells. The increased cytotoxicity of SV119-L[D] is consistent with the enhanced uptake of the SV119 liposomes, as shown in Figure 3. This demonstrates that the SV119 liposome can potentially be used a carrier for the targeted delivery of chemotherapeutics to tumor cells.

Despite the enhanced cellular uptake of the SV119 liposomes, it is crucial to understand the intracellular fate of the liposomes in order to maximize the bioavailability of the loaded drug. Nanoparticles internalized via the endocytosis pathway are often sequestered in endosomes and lysosomes, which represent an intracellular barrier that limits the accessibility of encapsulated agents to reach their molecular targets. In this study, confocal microscopy was used to examine the intracellular localization of rhodamine-PE labeled SV119 liposomes in DU-145 cells. After 3 hours of incubation, the liposomes showed a peri-nuclei punctuated distribution that is substantially co-localized with early endosomes (Figure 9A–C). After 13 hours of additional incubation, the liposomes were largely accumulated in the lysosomes, as evidenced by the colocalization with the lysosome markers (LysoTracker®; Figure 9D–F). These observations are consistent with the internalization via the endocytosis pathway. Future studies are warranted to improve the formulation and to achieve more efficient release from the endosomes and lysosomes.

Figure 9 Localization of SV119 liposomes in DU-145 cells. Cells were incubated with rhodamine-PE labeled SV119 liposomes for 3 hours (A–C) or 16 hours (D–F) in a serum-free medium, stained with markers of early endosomes (anti-EEA1; A–C) or lysosomes (LysoTracker®; Life Technologies, Carlsbad, CA) (D–F), and imaged using confocal microscopy. Red fluorescence represents rhodamine-PE labeled liposomes (A and D), green fluorescence represents early endosomes (B) or lysosomes (E). Yellow color observed in the red + green overlay (C and F) indicates colocalization of red liposomes with the green endosomes (C) or lysosomes (F).

Note: Scale bar = 10 μ m.

Conclusion

We have demonstrated the successful synthesis of the functional lipid, SV119-PEG-DOA, as well as its incorporation into a liposome formulation for targeted delivery to cancer cells. As compared with unmodified liposomes, the SV119 liposomes demonstrated enhanced uptake in multiple types of tumor cell lines including DU-145, MCF-7, PC-3, 201T, and A549 cells. In addition, this enhanced uptake was shown to be dependent on ligand density and incubation time, and the uptake of SV119 liposomes was dramatically inhibited by the presence of free SV119. In contrast, the enhanced uptake of SV119 liposomes was not observed in BEAS-2B normal cells, which suggests that enhanced uptake is tumor cell-specific. Furthermore, results of the MTT assay suggest that incorporating SV119 onto the surface of liposomes significantly enhances the delivery and cytotoxicity of liposomal DOX in vitro. At the time of writing this manuscript, Xia and colleagues reported that SV119 mediated the selective delivery of SV119-gold nanocage conjugates to tumor cells.³² Collectively, these data suggest that SV119-conjugated nanoparticles may represent a promising delivery system for the targeted delivery of anticancer therapeutics to tumor cells.

Acknowledgment

This work was supported in part by NIH grants (R21CA128415 and R21CA155983) and a DOD grant (BC09603).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Wang AZ, Gu F, Zhang L, et al. Biofunctionalized targeted nanoparticles for therapeutic applications. *Expert Opin Biol Ther*. 2008;8(8):1063–1070.
2. Wang X, Li J, Wang Y, et al. A folate receptor-targeting nanoparticle minimizes drug resistance in a human cancer model. *ACS Nano*. 2012;5(8):6184–6194.
3. Lee RJ, Low PS. Folate-mediated tumor cell targeting of liposome-entrapped doxorubicin in vitro. *Biochim Biophys Acta*. 1995;1233(2):134–144.
4. Moffatt S, Papasakelariou C, Wiehle S, Cristiano R. Successful in vivo tumor targeting of prostate-specific membrane antigen with a highly efficient J591/PEI/DNA molecular conjugate. *Gene Ther*. 2006;13(9):761–772.
5. Zhang YF, Wang JC, Bian DY, Zhang X, Zhang Q. Targeted delivery of RGD-modified liposomes encapsulating both combretastatin A-4 and doxorubicin for tumor therapy: In vitro and in vivo studies. *Eur J Pharm Biopharm*. 2010;74(3):467–473.
6. Maurice T, Su TP. The pharmacology of sigma-1 receptors. *Pharmacol Ther*. 2009;124(2):195–206.
7. Cobos EJ, Entrena JM, Nieto FR, Cendan CM, Del Pozo E. Pharmacology and therapeutic potential of sigma(1) receptor ligands. *Curr Neuropharmacol*. 2008;6(4):344–366.
8. John CS, Vilner BJ, Geyer BC, Moody T, Bowen WD. Targeting sigma receptor-binding benzamides as in vivo diagnostic and therapeutic agents for human prostate tumors. *Cancer Res*. 1999;59(18):4578–4583.
9. John CS, Bowen WD, Saga T, et al. A malignant-melanoma imaging agent – synthesis, characterization, in-vitro binding and biodistribution of Iodine-125-(2-piperidinylaminoethyl)4-Iodobenzamide. *J Nucl Med*. 1993;34(12):2169–2175.
10. Banerjee R, Tyagi P, Li S, Huang L. Anisamide-targeted stealth liposomes: A potent carrier for targeting doxorubicin to human prostate cancer cells. *Int J Cancer*. 2004;112(4):693–700.
11. Nakagawa O, Ming X, Huang L, Juliano RL. Targeted intracellular delivery of antisense oligonucleotides via conjugation with small-molecule ligands. *J Am Chem Soc*. 2010;132(26):8848–8849.
12. Li SD, Huang L. Targeted delivery of antisense oligodeoxynucleotide and small interference RNA into lung cancer cells. *Mol Pharm*. 2006;3(5):579–588.
13. Li SD, Chono S, Huang L. Efficient oncogene silencing and metastasis inhibition via systemic delivery of siRNA. *Mol Ther*. 2008;16(5):942–946.
14. Mukherjee A, Prasad TK, Rao NM, Banerjee R. Haloperidol-associated stealth liposomes: a potent carrier for delivering genes to human breast cancer cells. *J Biol Chem*. 2005;280(16):15619–15627.
15. Vilner BJ, John CS, Bowen WD. Sigma-1 and sigma-2 receptors are expressed in a wide variety of human and rodent tumor-cell lines. *Cancer Res*. 1995;55(2):408–413.
16. Wheeler KT, Wang LM, Wallen CA, et al. Sigma-2 receptors as a biomarker of proliferation in solid tumours. *Br J Cancer*. 2000;82(6):1223–1232.
17. Matsumoto RR. Sigma receptors: historical perspective and background. In: Matsumoto RR, Bowen WD, Su TP, editors. *Sigma Receptors: Chemistry, Cell Biology and Clinical Implications*. New York: Springer; 2007:1–24.
18. Mach RH, Smith CR, AlNabulsi I, Whirrett BR, Childers SR, Wheeler KT. Sigma-2 receptors as potential biomarkers of proliferation in breast cancer. *Cancer Res*. 1997;57(1):156–161.
19. Colabufano NA, Berardi F, Contino M, et al. Correlation between sigma2 receptor protein expression and histopathologic grade in human bladder cancer. *Cancer Lett*. 2006;237(1):83–88.

21. Kashiwagi H, McDunn JE, Simon PO, et al. Selective sigma-2 ligands preferentially bind to pancreatic adenocarcinomas: applications in diagnostic imaging and therapy. *Mol Cancer*. 2007;6:48.
22. Spitzer D, Simon PO, Kashiwagi H, et al. Use of multifunctional sigma-2 receptor ligand conjugates to trigger cancer-selective cell death signaling. *Cancer Res*. 2011;72(1):201–209.
23. Di Paolo D, Pastorino F, Brignole C, et al. Drug delivery systems: application of liposomal anti-tumor agents to neuroectodermal cancer treatment. *Tumori*. 2008;94(2):246–253.
24. Vangveravong S, Xu J, Zeng C, Mach RH. Synthesis of N-substituted 9-azabicyclo[3.3.1] nonan-3 α -yl carbamate analogs as σ 2 receptor ligands. *Bioorg Med Chem*. 2006;14(20):6988–6997.
25. Eliaz RE, Szoka FC Jr. Liposome-encapsulated doxorubicin targeted to CD44: a strategy to kill CD44-overexpressing tumor cells. *Cancer Res*. 2001;61(6):2592–2601.
26. Mach RH, Wheeler KT. Development of molecular probes for imaging sigma-2 receptors in vitro and in vivo. *Cent Nerv Syst Agents Med Chem*. 2009;9(3):230–245.
27. Xu J, Zeng C, Chu W, et al. Identification of the PGRMC1 protein complex as the putative sigma-2 receptor binding site. *Nat Commun*. 2011;2:380.
28. Spitzer D, Simon PO, Kashiwagi H, et al. Use of multifunctional sigma-2 receptor ligand conjugates to trigger cancer-selective cell death signaling. *Cancer Res*. 2012;72(1):201–209.
29. Lasic DD, Martin FJ, editors. *Stealth Liposomes*. Boca Raton: CRC Press; 1995.
30. Gabizon AA. Liposome circulation time and tumor targeting: implications for cancer-chemotherapy. *Adv Drug Deliv Rev*. 1995; 16(2–3):285–294.
31. Zhou W, Yuan X, Wilson A, et al. Efficient intracellular delivery of oligonucleotides formulated in folate receptor-targeted lipid vesicles. *Bioconjug Chem*. 2002;13(6):1220–1225.
32. Wang Y, Xu J, Xia X, et al. SV119-gold nanocage conjugates: a new platform for targeting cancer cells via sigma-2 receptors. *Nanoscale*. 2012;4(2):421–424.

Supplementary materials

Synthesis of SV119-PEG550

SV-119 succinate

A solution of SV-119 (2.0 g, 5 mmol) in CH_2Cl_2 (20 mL) was treated with DMAP (1.2 g, 10 mmol) and succinic anhydride (1 g, 10 mmol). The reaction was refluxed and stirred overnight. After the reaction was completed, the crude product was purified by flash chromatography with silica gel (EtOAc: PE, 3:1) to obtain SV-119 succinate.

SV119-PEG550

A solution of SV-119 succinate (2.5 g, 5 mmol) in CH_2Cl_2 (10 mL) was treated with DCC (0.82 g, 4 mmol), DMAP (0.12 g, 1 mmol), and MeO-PEG550-OH (2 g, 3.6 mmol). The reaction was stirred at room temperature overnight. After the reaction was completed, 100 mL of Et_2O was added and the mixture was filtered to afford the crude product. The crude product was purified by flash chromatography with silica

gel (MeOH: CH_2Cl_2 , 1:10) to afford pure SV119-PEG550. ^1H NMR δ 7.14 (s, 1H), 6.73–6.80 (m, 2H), 5.14 (m, 1H), 3.85 (s, 3H), 3.60–3.70 (m, 50H), 3.36 (s, 3H), 3.05–3.10 (m, 2H), 2.66–2.71 (m, 2H), 2.55–2.60 (m, 2H), 2.40–2.50 (m, 6H), 2.30 (s, 3H), 2.08–2.24 (m, 1H), 1.82–1.94 (m, 2H), 1.20–1.55 (m, 14H).

Method for testing the effect of SV119 on the uptake of SV119-liposomes

DU-145 cells were treated with SV119-liposomes (with 3% SV119-PEG-DOA) in a serum-free Dulbecco's modified Eagle's medium (DMEM; American Type Culture Collection [ATCC], Manassas, VA) for 1 hour in the presence of PEG550 (200 μM) or SV119-PEG550 (200 μM). The cells were treated with unmodified liposomes or SV119-liposomes alone as control groups. Fluorescence images were taken, as described in the section titled "Cellular uptake studies."

Figure S1 Scheme of synthesis for SV119-PEG550 (**A**) and effect of SV119 on the uptake of SV119-liposomes by DU-145 cells (**B**).
Abbreviation: PEG, polyethylene glycol.

Figure S2 MALDI-TOF mass spectrometry of PEG3500.

Abbreviations: MALDI-TOF, matrix-assisted laser desorption/ionisation-time of flight; PEG, polyethylene glycol.

Figure S3 MALDI-TOF mass spectrometry of PEG3500-DOA.

Abbreviations: DOA, dioleoyl amido aspartic acid; MALDI-TOF, matrix-assisted laser desorption/ionisation-time of flight; PEG, polyethylene glycol.

International Journal of Nanomedicine

Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-nanomedicine-journal>

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress