

Associations between functional ability and life satisfaction in the oldest old: results from the longitudinal population study Good Aging in Skåne

Åsa Enkvist
Henrik Ekström
Sölve Elmståhl

Department of Health Sciences, Lund University, Malmö University Hospital, Malmö, Sweden

Objectives: To describe change in functional ability in the oldest-old population during 3 years and examine its relation to life satisfaction (LS). A total of 681 individuals aged 78 and older from the population-based study Good Aging in Skåne took part.

Methods: Functional ability was assessed using Sonn and Åsberg's Activities of Daily Living (ADL) scale and related to LS assessed by Neugarten et al's Life Satisfaction Index A (LSI-A).

Results: Fifty-one percent of 87–93-year-olds reported ADL decline during 3 years. Individuals reporting impaired ADL had a mean LSI-A value of 23.0 compared to 26.4 in those unchanged. ADL decline had a stronger negative effect on LS in the younger group (78–84 years), $r = 0.207$, $P < 0.001$. In a multiple regression model, one score's decline in ADL capacity corresponded to 1.5 scores lower LS ($P < 0.001$).

Discussion: Effort put into keeping the oldest old on a high level of functional ability has the potential to maintain the LS of this population.

Keywords: life satisfaction, functional ability, longitudinal, oldest old

Introduction

During the last few decades, there has been great optimism in gerontology due to increased longevity and improved health and quality of life in the elderly population. Researchers such as Baltes and Smith, though, have highlighted the vast difference in prerequisites of the young old and the oldest old, and called for caution in optimism concerning the latter group: the more vulnerable, frail and those “at the limits of their functional ability.”¹ Different definitions are in use when it comes to the term “oldest old”; some use it for individuals aged 85 and above, some use 80 as the lower limit. In this study, we chose the latter limit to increase power in the study.

Baltes is well known for his aging theory of selective optimization with compensation (SOC).² Very briefly, selection implies focusing on areas of high priority as age-related losses in functioning put a limit to previous possibilities. Optimization means investing in behaviors that augment and maximize reserve capacities, and *compensation* means finding ways – for example, psychological or technological – to compensate for losses. In this framework, he concludes that as people age, the discrepancy between subjective and objective health increases. The elderly individual assesses his/her health as much better than an outside observer would as a psychological means of nurturing well-being and life satisfaction (LS) facing decreasing functional ability and other signs of declining health. Psychological coping strategies have been shown to mediate the effect between declining health and well-being.³

Correspondence: Åsa Enkvist
Department of Health Sciences, Division of Geriatric Medicine, Lund University, CRC, Malmö University Hospital, Malmö SE-20502, Sweden
Tel +46 40 391 000
Fax +46 40 391 313
Email asa.enkvist@med.lu.se

In the oldest old, the capacity to adapt and apply strategies to compensate is pushed to the limit as functional ability decreases; LS is thereby threatened.¹

We hypothesized that the ability to compensate for decreasing functional ability measured as diminished activities of daily living (ADL) capacity is exceeded in the oldest old and LS is threatened. This would be true even after controlling for factors such as age, number of symptoms, depressive mood, and personality factors previously known to affect LS^{4,5} and interact with functional ability. Studies in this area on the oldest old are rare, which is why we had to look for references in studies on somewhat younger individuals.

Asakawa et al⁶ followed 692 elderly people in Japan for 2 years. All were functionally intact at baseline. Among those, 12.3% experienced functional decline during the follow-up. Those subjects also reported a shrinking social network, increasing depressive symptoms, and lower LS than those that remained functionally intact. Avlund et al⁷ confirmed that people with decreased functional ability also have a diminished social network, something that is often considered a buffer against the negative impact of declining health.

Bowling and Grundy⁸ found that worse overall health and functional decline were the most significant predictors of lower LS and more depressive symptoms among people aged 65–84 years in a 3-year follow-up. Studies have also pointed to the fact that the sense of security and the feeling that help (from relatives and/or society) is available reduces the negative influence that functional ability has on LS.⁹

A paper studying people aged 65 and above in six European countries living with reduced ADL capacity¹⁰ concluded that it is not the reduced ADL capacity in itself that influences LS, but rather personality factors such as self-esteem, feeling worried, and the person's perception of their health. These three factors came out as significant in all six countries taking part in the study, whereas financial and social resources were significant in four out of six countries. But in a Swedish study by Hellström et al,¹¹ a reduced ADL capacity per se led to significantly lower LS among Swedish people aged 75 and above.

The rapid growth of the oldest-old population will mean increasing numbers of people living with some kind of functional disability. The search for clues of how to maintain LS while facing declining functional ability in the oldest old is of great importance.

The aim of this study was to describe the change in functional ability in the oldest-old population during 3 years. The aim was also to analyze possible associations between

changes in functional ability during the 3-year follow-up and LS adjusted for possible covariates.

Methods

Study population

The material in this study is from a longitudinal, randomized, population-based study of the elderly called Good Aging in Skåne (GÅS), which is part of the Swedish National Study on Aging and Care.¹² It started in 2001 and includes 2931 subjects aged 60 and above. Those aged 78 and above were chosen for this paper. Subjects were reexamined after a 3-year interval.

Five municipalities in southern Sweden were included. Participants underwent medical examination and functional tests, and filled in a questionnaire with identical study protocols at both examinations. Examination took place either at the clinic or in the participant's home. Informed consent was obtained from the participant and when needed from relatives.¹²

At baseline, 1253 individuals 78–93 years old participated. Of those, 314 died, eight moved from the area, and 250 chose not to take part in the reexamination. The study population consisted of the 681 individuals who came to the reexamination, including six age cohorts: 78, 81, 84, 87, 90, and 93 years. There was no difference in attrition between men and women, but the participants aged 87 and above had significantly higher attrition for the Life Satisfaction Index A (LSI-A) variable ($P < 0.001$). A total of 109 individuals did not answer the LSI-A questionnaire in the reexamination (15.9%); those individuals had significantly lower LSI-A in baseline examination (22.6 compared to 26.4 in the group that answered LSI-A at reexamination, $P < 0.001$). For further details on attrition, refer to Enkvist et al.¹³

Questionnaires

Neugarten's LSI-A was used to assess LS.¹⁴ It consists of five components of life satisfaction: zest versus apathy; resolution and fortitude; congruence between desired and achieved goals; positive self-concept; and mood tone. It includes 20 questions, scored from 0 to 2, for which participants had three possible answers to choose from: "disagree," "doubtful," and "agree." A high score thus indicates high LS (range 0–40). Cronbach's alpha reliability for internal consistency was established by Neugarten.¹⁴ In our material, Cronbach's alpha was 0.78.

Functional ability was assessed through self-report according to Sonn and Åsberg's ADL scale, the revised version.¹⁵ The nine activities included in the ADL staircase

are divided into four instrumental activities (i-ADL): transportation, cleaning, grocery shopping, and cooking; and five personal activities (p-ADL): bathing, dressing and undressing, toileting, mobility, and food intake. In our study, transportation was excluded due to high attrition for this item. Incontinence was excluded, since this has been considered an ability rather than an activity. In their study, Sonn and Åsberg also tested the scale and found good validity, and reliability was estimated at 0.9 using the Kuder–Richardson formula (KR-20). KR-20 is analogous to Cronbach's alpha, and a value above 0.65 is considered acceptable. In our study, Cronbach's alpha for ADL at baseline was 0.84.

The ADL staircase refers to the fact that the activities in the instrument can be arranged cumulatively in the form of a comprehensive nine-step ordinal scale (0–8), where 0 is given for a completely independent individual and 8 to someone dependent in all activities. Individuals needing assistance in activities such as food intake, the highest step, are likely to be dependent in all activities further down the staircase.¹⁶ Change in the ADL staircase was calculated as ADL at baseline minus ADL at reexamination. ADL was assessed with identical protocols at baseline and reexamination.

The Montgomery–Åsberg Depression Rating Scale is a subscale of the Comprehensive Psychopathological Rating Scale (CPRS),¹⁷ and was used to assess depressive mood. It includes ten questions, each graded in six steps concerning anxiety, sleep, appetite, concentration, initiative, emotional involvement, content of thoughts, and basic mood during the last 3 days. The score ranges from 0 to 60, where 0–6 indicates mental well-being, 7–34 mild to moderate depressive mood, and 35–60 severely depressive mood.¹⁸ In our population, no subject scored above 35, so this variable could be split in two groups: 1 for nondepressive ($n = 507$) and 2 ($n = 117$) for mildly to moderately depressive mood. The first group had a mean score of 1.7 (standard deviation [SD] 1.9), while the second group's mean score was 11.3 (SD 4.9).

The number of symptoms was rated with a modified version of the symptom scale from the Göteborg Quality of Life Instrument.¹⁹ Originally, the instrument included 30 symptoms; in the GÅS project, the following seven symptoms were added: difficulty walking, difficulty expressing myself verbally, difficulty swallowing, memory impairment, urinary and fecal incontinence, and slow-healing wounds. It thus contained 37 common physical and psychiatric symptoms, such as headache, obstipation, musculoskeletal pain, fatigue, restlessness, low-spiritedness, and sleeping disturbances. Participants responded with “yes” or “no” to whether they had

had symptoms during the last 3 months. The score thus ranged from 0 to 37, with a mean value of 11.8 (SD 6.3).

The health locus of control (LoC) scale²⁰ contains three subscales, one assessing internal LoC, the perception that the individual himself has the main control over his health. “LoC power” represents the notion that health status is mainly in the hands of powerful others, family members, or health-care professionals. “LoC chance” stands for the perceived importance of luck or fate as the main determinant of personal health. The score of each subscale ranges from 6 to 30, with a higher number indicating stronger beliefs in the aspect of LoC represented by the subscale.

Financial resources were assessed by the question: “Do your financial means cover your needs?” There were five possible answers: “very well,” “well,” “neither well nor insufficient,” “poorly,” and “not at all”. The first two alternatives were categorized as sufficient, and the last three as insufficient.

Married or cohabiting subjects were categorized as married, and singles, widows/widowers, and divorced subjects as singles.

Activities were subjected to a principal component analysis. The correlation matrix revealed many coefficients above 0.3. The Kayser–Meyer–Olkin value was 0.82, and Bartlett's test of sphericity reached significance ($P < 0.001$), supporting factorability of the correlation matrix. After inspection of the scree plot, three factors were used for further analysis: social, cultural, and leisure-time activities. Meeting friends/relatives, participation in study circles and nonprofit organizations, and going to restaurants were categorized as social activities. Cultural activities included going to concerts, theaters, movies, museums, art exhibitions, shows, musicals, and attending church services. Finally, leisure-time activities included long walks, gardening, repairing of cars or mechanical equipment, and traveling. Participants that had taken part at least once a year were categorized as active.

Domestic aid was assessed using the questions: “Do you receive help from relatives?” and “Do you receive help from the community?”

The subject was asked if he/she used any walking aids in- or outdoors. Answers were categorized as “yes” or “no.”

The subjects were categorized into two groups depending on whether their level of education was elementary school or high school and/or university.

Ethics

The study was approved by the regional ethics committee at Lund University, registration no LU 744-00. All subjects provided written consent for participation in the study.

Statistical methods

Differences in proportions in ADL change between men and women, the younger and the older age-group, higher/lower education, and married/single were tested with the chi-squared test (Table 1).

The population was stratified by gender, age, marital status, and education. Analysis of variance (ANOVA) was used to calculate mean LSI values for functionally impaired, unchanged, and improved in each group (Table 2).

Differences in mean values of LSI-A at reexamination regarding sex, age, marital status, education, walking aids, economic sufficiency, help from relatives, social services, and social, cultural, and leisure-time activities were tested with Student's *t*-test (Table 3). The LSI-A was normally distributed.

Correlations between LSI-A at reexamination and difference in ADL, depressive mood, and number of symptoms in the baseline examination were calculated using Spearman's correlation coefficient rho due to skewed distribution (Table 3). A linear multiple regression model was constructed with LSI as dependent variable, and regression coefficients were calculated. Independent variables included in the regression model were those that showed significant

Table 1 Changes in activities of daily living (ADL) functioning broken down by sex, age, marital status, education, and included subitems

	ADL-function			P-value
	Unchanged (%)	Impaired (%)	Improved (%)	
Sex				
Male/female	131/236 (58/65)	72/105 (32/29)	21/23 (9/6)	0.209
Age				
78–84/ 87–93 years	309/58 (70/40)	103/74 (23/51)	31/13 (7/9)	<0.001
Marital status				
Married/single	152/215 (64/61)	65/111 (28/32)	19/25 (8/7)	0.557
Education				
Elementary/high school or university	236/230 (62/63)	117/59 (31/28)	25/19 (7/9)	0.488
ADL items				
Cleaning	515 (76)	105 (15)	15 (2)	
Grocery shopping	521 (77)	102 (15)	16 (2)	
Cooking	515 (76)	92 (14)	30 (4)	
Bathing	572 (84)	63 (9)	4 (1)	
Dressing	599 (88)	33 (5)	6 (1)	
Toileting	593 (87)	41 (6)	5 (1)	
Mobility	630 (93)	24 (3)	0	
Food intake	641 (94)	5 (1)	0	
ADL scale	367 (62)	177 (30)	44 (8)	

Note: Differences in proportions tested with chi-squared test.
Abbreviation: ADL, activities of daily living.

Table 2 Comparison of mean Life Satisfaction Index A (LSI-A) value at reexamination for unchanged, impaired, and improved activities of daily living (ADL) stratified by sex, age, marital status and education

ADL	LSI-A value			P-value
	Unchanged	Impaired	Improved	
Sex				
Male	27.3 (6.6)	24.4 (7.3)	27.6 (7.2)	0.017
Female	25.9 (6.1)	22.0 (6.7)	25.4 (5.5)	<0.001
Age (years)				
78–84	26.4 (6.4)	23.5 (7.6)	27.5 (6.0)	<0.001
87–93	26.5 (6.0)	22.2 (6.0)	23.7 (7.3)	0.002
Marital status				
Married	27.3 (6.6)	25.6 (6.4)	27.2 (7.1)	0.233
Single	25.7 (6.0)	21.1 (6.9)	25.9 (5.8)	<0.001
Education				
Elementary	26.0 (6.3)	22.2 (7.1)	24.6 (5.7)	<0.001
High school/university	27.3 (6.3)	24.3 (6.8)	28.8 (6.7)	0.008

Note: Differences in means tested with analysis of variance, standard deviations in parentheses.

differences associated with levels of LSI-A when the level of significance was set to $P < 0.2$. The higher *P*-value, compared to the more common value of <0.05 , was chosen not to exclude variables that might have a decisive influence on LS.

All statistical tests were two-sided. Data analysis was performed using SPSS software version 18.0 (IBM, Armonk, NY).

Results

A total of 177 individuals (30%) reported impaired ADL function 3 years later, 82 of which were impaired by one step, 44 two steps, 24 three steps, 13 four steps, 8 five steps, 4 six steps, and 2 seven steps. On average, this group had declined by 1.7 steps in ADL-function. A total of 367 individuals (62%) reported unchanged ADL, while 44 (8%) scored higher in the reexamination, of whom 40 had increased in ADL function by one step, three subjects by two, and one by three steps. The whole study population was impaired by a mean value of 0.6 steps in the ADL staircase during the 3 years. The younger group – individuals aged 78–84 years – were impaired by 0.4 steps compared to the older group – individuals aged 87–93 – which decreased by 1.0 step ($P < 0.001$). The greatest decline in ADL was, as might be expected, in the older group, in which 51% reported a decline during the 3 years, compared to 23% among the younger subjects ($P < 0.001$). Even if results were not significant, the tendency was that more men reported impaired ADL compared to women: 32.1% compared to 28.8% (Table 1).

Clinical Interventions in Aging downloaded from https://www.dovepress.com/ by 3.238.130.97 on 14-May-2021 For personal use only.

Table 3 Comparison of means of Life Satisfaction Index A (LSI-A, range 0–40) at reexamination and variables at baseline examination

Variables	n	LSI at reexamination		P-value
		Mean ^a /standard deviation		
Sex				
Men	226	26.3/7.0		0.006
Women	346	24.7/6.5		
Age (years)				
78–84	430	25.7/6.8		0.005
87–93	142	23.8/6.3		
Marital status				
Married/cohabiting	246	26.7/6.6		<0.001
Single/widowed/divorced	323	24.2/6.6		
Education				
Elementary school	364	24.7/6.6		0.003
High school/university	203	26.4/6.7		
Walking aids				
Yes	161	23.6/6.6		<0.001
No	410	26.0/6.6		
Economical sufficiency				
Yes	421	25.9/6.5		<0.001
No	144	23.6/6.9		
Help from relatives				
Yes	112	22.7/6.3		<0.001
No	426	26.1/6.6		
Social services				
Yes	63	23.6/6.7		0.003
No	495	25.6/6.6		
Social activities				
Yes	500	25.7/6.6		0.003
No	55	22.7/7.0		
Cultural activities				
Yes	427	25.9/6.5		0.003
No	130	23.8/7.0		
Leisure-time activities				
Yes	472	25.8/6.6		<0.001
No	86	22.9/6.8		
Depressive mood (comprehensive psychopathological rating scale)				
No	507	25.1/6.4		<0.001
Yes	117	22.2/6.8		

Notes: Significance tested with Student's *t*-test; ^aSpearman's correlation coefficient *r* (rho).

When the different items in the ADL staircase were separated, the greatest declines were reported for cleaning, grocery shopping, and cooking.

Differences in mean LSI value stratified for sex, age, marital status, and education were compared for unchanged, impaired, and improved ADL function (Table 2). The group that reported impaired ADL had a mean LSI value of 23.0 compared to 26.4 in the group that was unchanged ($P < 0.001$) (Table 2). There were significant differences in LSI-A between all groups except for the married group. Post hoc tests (Tukey) revealed significant differences at the <0.05 level between unchanged and impaired ADL for

men, women, 87–93-year-olds, and the group with lower education. For 78–84-year-olds, singles, and the higher educated, differences were significant between unchanged and impaired ADL, as well as for impaired and improved ADL. For married subjects, there were no significant differences depending on change in ADL.

A higher mean value of LSI-A was noted for male gender, lower age, being married, higher education, being without walking aids, better economy, managing without help from family or social services, partaking in different activities, and no depressive mood (Table 3). A negative correlation was found between LSI-A and impaired ADL, locus of control chance, and a higher number of symptoms and depressive symptoms, in contrast to a positive correlation for locus of control internal and power (Table 4).

A linear multiple regression model was constructed in order to control for factors previously known to affect LS, such as age, number of symptoms, depressive mood, and personality factors, and factors such as walking aids and help from society or relatives interacting with functional ability. In the adjusted regression model difference in ADL, depressive mood, number of symptoms, health locus of control, education, and marital status in the baseline examination explained levels of LS 3 years later (Table 5).

Associations between changes of LSI from baseline to reexamination and change in ADL were analysed and stratified for age (Figure 1), and the younger group showed a positive association ($r = 0.207$ [$P < 0.001$]) in contrast to the older group ($r = -0.009$ [$P = 0.921$]).

Discussion

As we hypothesized, decline in ADL function predicted lower LS, as did more symptoms independently. Higher education, more internal locus of control, and cohabiting were also related to higher LS. This is in line with results from our previous study.⁵ Decline in ADL capacity was significantly

Table 4 Correlations between LSI-A at reexamination and parameters from baseline examination

	n	Correlation (r)	P-value
Change in activities of daily living scale	537	-0.219 ^a	<0.001
Locus of control			
Chance	556	-0.046 ^b	0.276
Power	554	0.134 ^b	0.002
Internal	557	0.210 ^b	<0.001
Number of symptoms	554	-0.366 ^b	<0.001
Depressive mood	534	-0.285 ^a	<0.001

Notes: ^aSpearman's correlation coefficient *r*; ^bPearson's correlation coefficient *r*.

Abbreviations: LSI-A, Life Satisfaction Index A; *r*, rho.

Table 5 A multiple linear regression model with independent variables from baseline examination and Life Satisfaction Index 3 years later as dependent variable

Variable	B-coefficient	P-value
Sex		
Female/male	0.438	0.474
Age	-0.128	0.086
Married/cohabiting vs single	-1.370	0.028
Education elementary/secondary	1.182	0.035
Economical sufficiency, no/yes	1.225	0.054
Depressive mood, yes/no	1.826	0.014
Number of symptoms	-0.336	<0.001
Locus of control, internal	0.239	0.001
Locus of control, power	0.153	0.019
Social activities, no/yes	1.178	0.234
Cultural activities, no/yes	1.155	0.111
Leisure-time activities, no/yes	0.808	0.361
Help from relatives, no/yes	-0.304	0.692
Help from social services, no/yes	-1.942	0.058
Walking aids, no/yes	-1.205	0.099
Change in activities of daily living scale	-1.316	<0.001

associated with lower LS, even after adjusting for partaking in social, cultural, and leisure-time activities, age, and other independent factors like economical sufficiency and social support. Reconnecting with Baltes' SOC theory discussed in the introduction,² we conclude that in today's society, the ability to apply SOC strategies to overcome the threat that declining functional ability constitutes on LS is limited.

Twenty-three percent of the 78–84-year-old group showed a decline in ADL function compared to 51% in the

older group. This represents an increased vulnerability in the higher ages, even if this group constitutes “survivors.” The main decline was seen in the I-ADL items. (One step's decline in the ADL staircase means losing independence in one item; for example, not being able to do one's own grocery shopping or cooking anymore.)

Interestingly, when comparing the older group in our population to the somewhat younger one, it seems the oldest group is less affected in LS when ADL-function decreases. This could reflect lower expectations and more readiness to accept functional decline in the 87–93-year-old group. Other factors, such as an increasing burden from symptoms or losing a spouse or close friends, could have an even higher impact than declining functional ability on LS. However, the 78–93-year-old group with impaired ADL did show lower LS than the same group with unchanged ADL. In the 78–84-year-old group, there was a decline in LS as ADL declined. This could reflect a life crisis in individuals having enjoyed a totally independent life suddenly having to face needing help from spouses or formal care from society.

Functional ability in terms of number of symptoms is one of the most tangible aspects by which the elderly person perceives his health. In our previous work^{5,13} conducted on the same population, the burden of symptoms had significant influence on LS, especially symptoms in the following categories: depression, tension, gastrointestinal, and musculoskeletal. Specific diagnoses such as heart failure, dementia, stroke, or fracture did not have as strong an influence on LS as the burden of symptoms. The elderly individual seems to perceive his health based on his experiences of daily life rather than objective medical findings, and the elderly person's perceived health has been shown to influence his LS.²¹

Asakawa et al⁶ noted an increase in depressive symptoms with declining functional ability, but it has also been discussed that preventing and treating depression in old age might reduce functional decline.²² Here lies the potential for interdisciplinary interaction to reach results that increase the LS of the elderly.

Our results suggest that a lower functional ability results in lower LS, regardless of the availability of help from community or family caregivers. Though not significant, there was a negative association between receiving help and LS. This might seem contradictory, but the impact on the elderly of having no choice but to receive public services has been related to psychological strain,²³ potentially negatively influencing LS. The negative association was there even for receiving help from relatives, and should probably be interpreted as an effect of the elderly persons'

Clinical Interventions in Aging downloaded from https://www.dovepress.com/ by 3.238.130.97 on 14-May-2021 For personal use only.




Figure 1 Change in Life Satisfaction Index A (LSI-A) from baseline to reexamination depending on change in activities of daily living (ADL) in the same period between the younger and the older group ($P = 0.0273$).
Note: The correlation between change in LSI-A and changes in ADL for the younger group (81–87 years) was $r = 0.207$ ($P < 0.001$).

loss of independence rather than having a relative prepared to help. Though the comfort of the presence of loved ones was a positive aspect, feelings of guilt for being a burden on relatives were also evident in Janlöv et al's study,²³ and could further explain the negative association with LS.

In Asakawa et al's study,⁶ those reporting impaired functional ability also reported a shrinking social network and lower LS than those that remained functionally intact. We included partaking in different activities as a measure of the subjects' social life and ability to be a part of the world around them. Impaired functional ability resulted in lower LS even after this adjustment, but being married or cohabiting was associated with higher LS. Living with a spouse could, apart from providing social anchorage, buffer a decline in functional ability, as some degree of help is more available than to those living alone. The need for social services might thus be less pronounced.

Economical sufficiency did not come out as a significant factor in the regression model. Perhaps a reason for this is that the Swedish health-care system provides health care, social services, and different kinds of aids at little or no cost, buffering the impaired functional ability without the extra burden of financial worries.

The study population comprised a sample from the general population covering urban as well as rural areas. It was randomly selected using the Population Registry, thereby increasing generalization of results to the older general population. Higher ages were also included, and no exclusion was made for subjects living in residential care or homes for the elderly. Home visits were made in almost every third case to reduce selection bias. Social, medical, and psychological aspects were included in the model. Further strengths in this study are the longitudinal design and that it comprises a broad context of health-related and psychosocial factors. The instruments used in this study (such as Neugarten, ADL, and CPRS) were standardized and internationally well-known instruments that have shown good reliability in previous validations.^{14–17} Proxy information was retrieved from next of kin and ward staff to ensure validity and completeness of information. Cognitive assessment by medical examination was done for all participants in a standardized way.

In the attrition analysis, it was noted that the younger participants who were more satisfied with life and had fewer depressive and physical symptoms were those that came back to the reexamination.⁵ Respondents from the 87–93-year-old group constitute the main attrition in LSI-A. If these had been included, the correlation between age and decline in ADL would have been even stronger, since functional ability

declines with higher age. Furthermore, the lack of association between changes of ADL and changes in LS among the oldest old might be influenced by selection bias in this group. People with severe dementia were not able to complete the questionnaire, and this condition is more prevalent with age and related to decline in functioning, thereby introducing possible bias. Efforts were made to include proxy information if possible.

One advantage of using the ADL scale is that each step is represented by a defined categorization, which makes it easy to compare different individuals' improvement or decline in ADL. It is important to remember that the steps of the scale are not equivalent, and that differences in the experience of the various steps will vary among individuals. In addition, individual differences exist between individuals on the same step.¹⁶

The ADL scale is a rather coarse instrument in measuring functional ability, not able to capture milder limitations that might nevertheless be experienced as a great hindrance to the person affected. In a Swedish investigation of almost 1300 people aged 60–89 years, half of the respondents reported feeling hindered by health problems in spite of having no decline in the ADL scale.⁹ Even milder declines in functional ability, not measurable by ADL, have been shown to be connected with lower LS.⁶

Conclusion

High LS throughout the life span is an attractive goal to the elderly individual as well as to anyone privately or professionally engaged with individuals in this age-group. The above results confirm the hypothesis that reduced functional ability leads to lower LS, even after controlling for confounders. The results also suggest that investing in the maintenance and/or restoration of ADL capacity in the age-group 78–84 years has the potential to preserve or even increase the LS of the oldest old. If this association is not as strong for the older group, aged 87–93 years, this has consequences for how far different kinds of treatment strategies should be pushed. More research is needed to confirm and further explore these results. Another interesting area for further research would be an intervention study, testing effects on LS after rehabilitation efforts in this age-group.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Baltes P, Smith J. New frontiers in the future of aging: from successful aging of the young old to the dilemmas of the fourth age. *Gerontology*. 2003;49(2):123–135.

2. Baltes P, Baltes M, editors. *Successful Aging: Perspectives from the Behavioral Sciences*. Cambridge: Cambridge University Press; 1990.
3. Jonker AA, Comijs HC, Kees KC, Deeg DJ. The role of coping resources on change in well-being during persistent health decline. *J Aging Health*. 2009;21(8):1063–1082.
4. Berg AI, Hassing LB, Mclearn GE, Johansson B. What matters for life satisfaction in the oldest-old? *Aging Ment Health*. 2006;10(3):257–263.
5. Enkvist Å, Ekström H, Elmståhl S. What factors affect life satisfaction (LS) in the oldest-old? *Arch Gerontol Geriatr*. 2012;54(1):140–145.
6. Asakawa T, Koyano W, Ando T, Shibata H. Effects of functional decline on quality of life among the Japanese elderly. *Int J Aging Hum Dev*. 2000;50(4):319–328.
7. Avlund K, Due P, Holstein BE, Heikkinen RL, Berg S. Changes in social relations in old age. Are they influenced by functional ability? *Aging Clin Exp Res*. 2002;14(Suppl 3):56–64.
8. Bowling A, Grundy E. Activities of daily living: changes in functional ability in three samples of elderly and very elderly people. *Age Ageing*. 1997;26(2):107–114.
9. Fagerström C, Holst G, Hallberg IR. Feeling hindered by health problems and functional ability at 60 years and above. *Arch Gerontol Geriatr*. 2007;44(2):181–201.
10. Borg C, Fagerström C, Balducci C, et al. Life satisfaction in 6 European countries: the relationship to health, self-esteem and social and financial resources among people (aged 65–89) with reduced functional ability. *Geriatr Nurs*. 2008;29(1):48–57.
11. Hellström Y, Andersson M, Hallberg IR. Quality of life among older people in Sweden receiving help from informal and/or formal helpers at home or in special accommodation. *Health Soc Care Community*. 2004;12(6):504–516.
12. Ekström H, Dahlin Ivanoff S, Elmståhl S. Restriction in social participation and lower life satisfaction among fractured in pain. Results from the population study “Good aging in Skåne”. *Arch Gerontol Geriatr*. 2008;46(3):409–424.
13. Enkvist Å, Ekström H, Elmståhl S. Life satisfaction and symptoms among the oldest-old: results from the population study called Good Ageing in Skåne. *Arch Gerontol Geriatr*. 2012;54(1):146–150.
14. Neugarten BL, Havighurst RJ, Tobin SS. The measurement of life satisfaction. *J Gerontol*. 1961;16:134–143.
15. Sonn U, Åsberg KH. Assessment of activities of daily living in the elderly. A study of a population of 76-year-olds in Gothenburg, Sweden. *Scand J Rehab Med*. 1991;23(4):187–191.
16. Åsberg KH, Sonn U. The cumulative structure of personal and instrumental ADL. A study of elderly people in a health service district. *Scand J Rehab Med*. 1988;21(4):171–177.
17. Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry*. 1979;134:382–389.
18. Snaith RP, Harrop FM, Newby D, Teale C. Grade scores of the Montgomery-Asberg Depression and the Clinical Anxiety Scales. *Br J Psychol*. 1986;148:599–601.
19. Tibblin G, Tibblin B, Peciva S, Kullman S, Svärdsudd K. “The Göteborg quality of life instrument” – an assessment of well-being and symptoms among men born 1913 and 1923. Methods and Validity. *Scand J Prim Health Care Suppl*. 1990;1:33–38.
20. Walston BS, Walston KA, Kaplan GD, Maides SA. Development and validation of the Health Locus of Control (HLC) Scale. *J Consult Clin Psychol*. 1976;44(4):580–585.
21. Hillerås P, Jorm AF, Herlitz A, Winblad B. Life satisfaction among the very old: a survey on a cognitively intact sample aged 90 years or above. *Int J Aging Hum Dev*. 2001;52(1):71–90.
22. Palmore EB, Burchett BM. Predictors of disability in the final year of life. *J Aging Health*. 1997;9(3):283–297.
23. Janlöv AC, Hallberg IR, Petersson K. Older persons’ experience of being assessed for and receiving public home help: do they have any influence over it? *Health Soc Care Community*. 2006;14(1):26–36.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, the American Chemical Society’s ‘Chemical Abstracts Ser-

vice’ (CAS), Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress