Coronary heart disease is not significantly linked to acute kidney injury identified using Acute Kidney Injury Group criteria

Josef Yayan

Department of Internal Medicine,
Vinzentius Hospital, Landau, Germany

Background: Patients with unstable angina or myocardial infarction are at risk of acute kidney injury, which may be aggravated by the iodine-containing contrast agent used during coronary angiography; however, the relationship between these two conditions remains unclear.

Objective: The current study investigated the relationship between acute kidney injury and coronary heart disease prior to coronary angiography.

Methods: All patients were evaluated after undergoing coronary angiography in the cardiac catheterization laboratory of the Vinzentius Hospital in Landau, Germany, in 2011. The study group included patients with both acute coronary heart disease and acute kidney injury (as defined according to the classification of the Acute Kidney Injury Group); the control group included patients without acute coronary heart disease. Serum creatinine profiles were evaluated in all patients, as were a variety of demographic and health characteristics.

Results: Of the 303 patients examined, 201 (66.34%) had coronary artery disease. Of these, 38 (18.91%) also had both acute kidney injury and acute coronary heart disease prior to and after coronary angiography, and of which in turn 34 (16.91%) had both acute kidney injury and acute coronary heart disease only prior to the coronary angiography. However, the occurrence of acute kidney injury was not significantly related to the presence of coronary heart disease ($P = 0.95$, Chi-square test).

Conclusion: The results of this study indicate that acute kidney injury is not linked to acute coronary heart disease. However, physicians should be aware that many coronary heart patients may develop kidney injury while hospitalized for angiography.

Keywords: acute kidney injury, acute coronary heart disease, myocardial infarction, angina pectoris, incidence, association

Introduction

Physicians often ignore increases – especially minor ones – in the serum creatinine levels of patients with angina pectoris or myocardial infarction.¹ However, such increases may be indicative of acute kidney injury, and may therefore be important to investigate in further detail. Unfortunately, there has been little, if any, research on either the relationship between acute kidney injury and acute coronary heart disease or the frequency of acute kidney injury in patients with acute coronary syndrome and acute myocardial infarction.¹

Because the diagnosis of acute renal failure has not been standardized, it is difficult to study the outcomes of this condition.¹ Rodrigues et al recently reported finding more than 30 definitions of “acute renal failure” in the medical literature;¹ this variety causes confusion and makes it difficult to compare the results of multiple

Correspondence: Josef Yayan
Department of Internal Medicine,
Vinzentius Hospital, Cornichonstraße 4,
D-76829, Landau, Germany
Tel +49 6341 17 6128
Fax +49 6341 14 2254
Email josef.yayan@hotmail.com

studies by different authors.¹ For example, although acute renal failure has been described as a complex, multicausal dysfunction with clinical features ranging from small changes in serum creatinine to anuric renal failure, it has also been defined using absolute values or percentile increases in serum creatinine levels or different rates of reduction in urine output.¹ Recently, new definitions have been proposed to standardize the diagnosis of acute renal failure.^{1,2} These establish the concept of acute kidney injury¹ by emphasizing the subtle but important differences between the terms “acute kidney injury” and “acute renal failure.”¹ One important distinction between these is that injury precedes failure.¹ Several studies have shown that small increases in serum creatinine levels during hospitalization are associated with a worse prognosis for the patient.¹

The Acute Kidney Injury (AKIN) group developed a classification system that allows the observation of small changes in serum creatinine over a short period of time to be used in the early diagnosis and treatment of acute kidney injury.^{1,2} Specifically, the group defined acute kidney injury as a state marked by one of the following three criteria: an absolute increase in serum creatinine level, to >0.3 mg/dL (≥ 26.4 $\mu\text{mol/L}$); an increase of more than 50% (or 1.5 times) from baseline over a 48-hour period; or documented oliguria of <0.5 mL/kg/h for >6 hours.^{1,2} One important feature of this new definition is the concept of staging the severity of acute kidney injury.¹

The purpose of the present investigation was to determine the association between the onset of acute kidney injury and acute coronary heart disease. All patients underwent coronary angiography and provided multiple blood samples that could be subjected to a variety of tests, including evaluation of serum creatinine levels, to find out acute kidney injury prior coronary angiography.

Methods

I conducted a retrospective study of all patients ($N = 303$) admitted to the Vinzentius Hospital (Landau, Germany) in 2011 after undergoing coronary angiography in the cardiac catheterization laboratory. In all, 201 patients were diagnosed with coronary artery disease. The remaining 102 individuals served as the control population; only 19 of these patients suffered from acute kidney injury. Among individuals in the study population, 38 had acute coronary heart disease and were diagnosed with acute kidney injury prior or after to their coronary angiography.

For this study, acute kidney injury was defined in one of two ways, both according to the AKIN classification. For

patients whose baseline creatinine levels were known, acute kidney injury was defined as an increase of more than 50% (or 1.5 times) within a 48-hour time frame. For patients with unknown baseline creatinine levels, acute kidney injury was defined as an absolute increase of ≥ 0.3 mg/dL (≥ 26.4 $\mu\text{mol/L}$) from the upper-limit value of the normal range of serum creatinine (0.6–1.2 mg/dL) stated in the Clinical Laboratory and Standards Institute guidelines.³ Prior to normalization, serum creatinine was measured several times in patients in whom levels were elevated. The number of different measurements of serum creatinine obtained for each patient was compared in all patients. Thus, the profile of the serum creatinine was controlled for all patients from admission to discharge from the hospital. By monitoring the course of serum creatinine, I was able to determine whether acute kidney injury occurred before or after coronary angiography.

Chronic renal failure was defined and classified based on the Guidelines of the National Kidney Foundation.⁴ The glomerular filtration rate was used to calculate the level of kidney function and determine the stage of kidney disease. When possible, the glomerular filtration rate was evaluated three times per patient: before hospitalization, on admission, and prior to discharge from the hospital. Patients with poor kidney values received intravenous infusion therapy with 1000 mL 0.9% sodium chloride both before and after coronary angiography.

Iopamidol (Solutrast[®], ALTANA Pharma AG, Konstanz, Germany) was used as a contrast medium during coronary angiography. Both the amount of contrast agent used and the duration of the coronary angiography were measured for each patient. The most recent version of the International Classification of Diseases (ICD) was used to define the symptoms and treatments of stable angina, acute coronary syndrome, ST segment elevation myocardial infarction (STEMI), non-STEMI, and chronic coronary heart disease. Patient electrocardiograms were recorded in the emergency room. Changes to these patterns, such as ST-segment elevation, ST-segment depression, left bundle branch block, R-regression, T-negativity, and presence or absence of cardiac markers in the blood, were used to identify myocardial infarctions.

Coronary angiography, which most often combines diagnosis with percutaneous coronary intervention (ie, angioplasty and stenting), was examined in both groups after past and present coronary angiography with the number of stents in each patient and former bypass surgery. The indication for stent or bypass surgery was placed during coronary angiography by the examiner. Coronary heart disease lesions were

classified as being 1-, 2-, or 3-vessel; further, examiners visually estimated the extent of stenosis diameter (as a percentage of the cardiac catheter), as per the stenosis morphology classification recommendations of the American College of Cardiology/American Heart Association.⁵

For all patients, possible adverse events were documented during the coronary angiography. Angiography was obtained urgently for patients with STEMI, persistent chest pain despite maximal medical therapy, or complications (eg, markedly elevated cardiac markers, the presence of cardiogenic shock, acute mitral regurgitation, and unstable arrhythmias). Patients with uncomplicated non-STEMI or unstable angina whose symptoms have resolved typically undergo angiography within the first 24 to 48 hours of hospitalization to detect lesions that may require treatment. After initial evaluation and therapy, coronary angiography was used in patients with stable angina after a positive ergometric evaluation or for assessment of valvular heart diseases. A cardiac ultrasound was used to assess systolic left ventricular function in all patients.

Basic demographic and health information was collected for each patient at the time of admission and of discharge from the hospital; this included age, gender, whether the patient was taking an angiotensin-converting enzyme inhibitor (ACE), duration of hospital stay, and door-to-needle time for the coronary angiography. Because oliguria was not a problem for any of the patients included in this study, this symptom was not closely examined. However, data were collected on the presence of specific cardiovascular risk factors (CVRFs) such as elevated blood pressure, diabetes mellitus, hyperlipidemia, obesity, smoking, and family history of cardiovascular disease.

For the statistical analysis, I calculated the odds ratio and prior probability (odds) that a patient would have both acute kidney and acute coronary syndrome. All results are expressed as means and standard deviations (SDs). A Mann-Whitney *U*-test was used to investigate differences between the two treatment groups (in age, total number of diagnoses, indications for coronary angiography, heart rhythm, changes in electrocardiogram, use of ACE inhibitors, serum creatinine levels, total number of CVRFs, length of hospital stay, duration of coronary angiography, amount of contrast medium, door-to-needle time, total number of pretreatments, heart failure, ventricular dysfunction, stenting, presence coronary heart disease, and presence of acute kidney injury).

A Chi-square test was used to investigate the differences in gender distribution, number of diagnoses, indications for coronary angiography, heart rhythm, changes in

electrocardiogram, use of ACE inhibitors, CVRF, stages of chronic renal failure, number of pretreatments, heart failure, ventricular dysfunction, stenting, coronary heart disease, and acute kidney injury between study and control group.

I also calculated the overall incidence of acute kidney injury associated with coronary artery disease, as well as associated 95% confidence intervals. Survival rates for both treatment groups were calculated using the Kaplan-Maier method. Significance was defined as $P < 0.05$.

Results

There were 303 patients included in the current study: 191 (63.04%) men and 112 (36.96%) women. The average age across all study participants was 69.64 ± 12.12 years.

The study group comprised 38 (12.54%) patients who were, on average, 76.90 ± 7.64 years old and who had both acute kidney injury and acute coronary disease during the whole hospital stay. Thirty-four patients (16.91%) had both acute kidney injury and acute coronary disease prior to undergoing a coronary angiography ($N = 201$). Over half the patients (163, or 53.80%) had acute coronary disease without acute kidney injury; 108 (66.26%) of these were men and 55 (33.74%) were women.

A total of 102 patients (or 33.66%) composed the control group ($N = 303$). The average age of the control patients was 69.08 ± 13.09 . The average age was significantly higher in the study group than in the control group ($P = 0.002$), but the gender distribution was similar in both groups ($P = 0.18$) (Figure 1).

The prior probability (odds) for the occurrence of acute kidney injury in coronary heart disease was found to be 66% (95% CI, 64%–69%). The odds ratio for the occurrence of acute kidney in acute coronary syndrome was 1.0184 (95% CI, 0.5529–1.8759; $P = 0.95$). No statistical association was found between acute kidney injury and acute coronary syndrome ($\chi^2 = 0.0034$, $df = 1$, $P = 0.95$).

Stable angina pectoris, sinus rhythm, and T-negativity were more commonly found on the electrocardiograms of patients in the control group than on those of patients in the study group ($P = 0.0002$, $P = 0.02$, and $P = 0.013$, respectively) (Table 1). There was only a marginal statistical difference in the number of CVRFs reported for the two groups ($P = 0.041$) (Table 2).

Patients in both groups had similar numbers of pretreatments, heart failure, ventricular function, stenting, and acute kidney injury (all $P > 0.11$) (Table 3). However, acute coronary heart disease was more common in the study group ($P < 0.01$), while patients in the control group were more

Figure 1 Gender distribution in the study and control groups.

Note: Lavender indicates male patients and burgundy indicates female patients.

Table 1 Diagnoses and clinical and medical characteristics of patients in the study and control groups

	Study group N (%)	Control group N (%)	P
Diagnosis			
Stable angina pectoris	10 (15.63)	55 (42.97)	0.0002
Acute coronary syndrome	29 (45.31)	43 (33.59)	0.11
NSTEMI	20 (31.25)	27 (21.09)	0.12
STEMI	5 (7.81)	3 (2.34)	0.07
Total number of diagnoses	64	128	0.49
Other indications for coronary angiography			
Vitium	5 (71.43)	10 (41.67)	0.17
Ergometry	2 (28.57)	14 (58.33)	0.17
Total number	7	24	0.33
Heart rhythm			
Sinus rhythm	27 (71.05)	89 (87.25)	0.02
Atrial fibrillation	8 (21.05)	9 (8.82)	0.049
Atrial fluttering	1 (2.63)	0	0.10
Ventricular fibrillation	2 (5.26)	4 (3.92)	0.73
Total number	38	102	0.89
Changes in electrocardiogram			
T-negativity	4 (26.67)	15 (68.18)	0.013
ST-segment depression	1 (6.67)	2 (9.09)	0.79
ST-segment elevation	3 (20.00)	1 (4.55)	0.14
Left bundle branch block	4 (26.67)	1 (4.55)	0.053
R-regression	3 (20.00)	3 (13.64)	0.61
Total number	15	22	0.55
ACE inhibitor			
ACE-taking before this stay	9 (29.03)	23 (30.67)	0.87
ACE-taking during this stay	22 (70.97)	52 (69.33)	0.87
Total number	31	75	0.33

Note: P-values in bold font are those indicating significant differences between the two groups.

Abbreviations: ACE, angiotensin converting enzyme; STEMI, myocardial infarction with ST elevation; NSTEMI, non-ST segment myocardial infarction.

likely to have chronic coronary heart disease or no heart disease at all ($P < 0.01$) (Table 3).

Patients in the study group had significantly higher levels of serum creatinine (1.98 ± 0.92 mg/dL) than those in the control group (1.23 ± 0.57 mg/dL) ($P < 0.001$) (Figures 2 and 3). The mean value of the number of controls of serum creatinine was 5.67 (SD = 3.89) times in the study group and 3.97 (SD = 4.93) times in the control group. More controls of serum creatinine for each patient were conducted in the study group ($P < 0.01$).

Chronic renal failure was equally common in the control and study groups at admission (Stage 1: $P = 0.43$; Stage 2: $P = 0.06$; Stage 3: $P = 0.25$; Stage 4: $P = 0.32$; Stage 5: $P = 0.10$) and during previous hospital stays with statistically significant difference in Stage 2 (Stage 1: $P = 0.40$; Stage 2: $P = 0.03$; Stage 3: $P = 0.11$; Stage 4: $P = 0.97$; Stage 5: $P = 0.40$). Statistically significant difference between

Table 2 Cardiovascular risk factors (CVRF) and their distribution across the study population

CVRF	Study group N (%)	Control group N (%)	P
Hypertension	34 (35.42)	79 (34.05)	0.81
Diabetes	24 (25.00)	38 (16.38)	0.07
Hyperlipidemia	18 (18.78)	48 (20.69)	0.69
Obesity	6 (6.25)	18 (7.76)	0.63
Nicotine habit	11 (11.46)	26 (11.21)	0.95
Family history of cardiovascular disease	3 (3.13)	23 (9.91)	0.038
Total cases of diagnosis (N = 328)	96	232	0.041

Note: P-values in bold font are those indicating a significant difference between the study and control groups.

Table 3 Heart and kidney characteristics of patients in the study and control groups

	Study group N (%)	Control group N (%)	P
Pretreatments			
Coronary angiography previously	16 (53.33)	43 (52.44)	0.93
Coronary angiography prior to this stay	11 (36.67)	24 (29.27)	0.45
Bypass surgery	3 (10.00)	15 (18.29)	0.29
Total cases	30	82	0.2
Ventricular function			
Heart failure	14 (26.92)	26 (20.31)	0.33
Good systolic left ventricular function	24 (46.15)	76 (59.38)	0.11
Slightly reduced left ventricular function	6 (11.54)	8 (6.25)	0.23
Moderately reduced left ventricular function	7 (13.46)	13 (10.16)	0.52
Greatly reduced left ventricular function	1 (1.92)	5 (3.91)	0.50
Total cases	52	128	0.42
Percutaneous interventional angiography	21 (47.73)	6 (50.00)	0.89
1-stent	17 (38.64)	4 (33.33)	0.74
2-stent	4 (9.09)	0	0.28
Indication for bypass surgery	2 (4.55)	2 (16.67)	0.15
Total cases	44	12	0.34
Presence/absence of coronary heart disease			
Acute coronary heart disease	38 (84.44)	0	0.0001
No coronary heart disease	0	48 (48.48)	0.0001
Chronic coronary heart disease	7 (15.56)	51 (51.52)	0.0001
Total cases	45	99	0.4
Type of coronary heart disease			
1-vessel coronary heart disease	8 (20.51)	6 (12.00)	0.27
2-vessel coronary heart disease	11 (28.21)	14 (28.00)	0.98
3-vessel coronary heart disease	20 (51.28)	30 (60.00)	0.41
Total cases	39	50	1.0
Acute kidney injury	38 (39.58)	19 (39.58)	0.99
Acute kidney injury prior coronary angiography	34 (35.42)	17 (35.42)	0.99
Acute kidney injury after coronary angiography	24 (25.00)	12 (25.00)	0.99
Total cases	96	48	0.1

Note: P-values in bold are those indicating significant differences between the two groups.

the groups was found at discharge from the hospital; specifically, the study group had higher levels of Stage 4 chronic renal failure (Stage 1: $P = 0.40$; Stage 2: $P = 0.23$; Stage 3: $P = 0.72$; Stage 4: $P = 0.022$; Stage 5: $P = 0.97$) (Figure 4). No patients suffered an oliguria of <0.5 mL/kg/h for more than 6 hours.

Length of hospital stay was significantly greater for patients in the study group (9.45 ± 6.28 days) than for those in the control group (7.24 ± 6.03 days) ($P = 0.01$). Likewise, door-to-needle time was longer in the study

group (52.30 ± 66.24 hours) than in the control group (22.84 ± 39.18 hours) ($P = 0.0016$).

No adverse events were observed in any of the patients examined. Although more contrast agent was required for patients in the study group (108.37 ± 53.37 mL) than in the control group (93.19 ± 33.67 mL), this difference was not significant ($P = 0.32$). Also nonsignificant was the difference in duration of coronary angiography ($P = 1.0$), which was the same, up to 120 minutes, in both groups.

Three patients died during the study; one of these was from the study group (2.63% of the total), and the other two were from the control group (1.96% of the total). Mortality rates were similar in the two groups ($P = 0.81$). However, survival rates were better in the control group (98.04% [95% CI, 95.32–100.76%]) than in the study group (97.37% [95% CI, 92.21%–102.53%]).

Discussion

Among the patients examined in the current study, one in five suffered from both acute coronary heart disease and acute kidney injury. Nearly all individuals were hypertensive, and many patients displayed multiple risk factors for acute coronary heart disease. Patients in the study group were more likely to have myocardial infarction and acute coronary heart disease. Further, study individuals possessed significantly higher levels of serum creatinine, though the only significance in chronic renal failure was found at hospital discharge and previous hospital stay, and only for individuals with Stage 2 and 4 failure. Cumulatively, these results indicate that kidney damage is a real threat for angina pectoris and heart attack patients, and that care must be taken to minimize risk to these individuals during medical treatment.

In a prospective study of STEMI patients, Goldberg et al found that 9.6% of individuals suffered worsening renal injury (defined arbitrarily as a >0.5 mg/dL elevation of the serum creatinine level at any point during the hospital stay) during their hospital stay.⁶ In the current study, however, the incidence of STEMI was only slightly higher among patients with acute kidney injury and coronary heart disease, indicating that there is no association between acute kidney injury and acute coronary heart disease. This apparent lack of association between the two diseases, one linking acute renal injury and acute heart disease, was also suggested by the results of a Chi-square analysis ($P = 0.95$). The discrepancies between this and the Goldberg et al study⁶ may stem from the fact that the previous work did not use AKIN criteria for classifying acute kidney injury and, further, there was no specific period of time during which worsening renal function or acute kidney injury were evaluated. The absence of a defined focal period

Figure 2 Serum creatinine profiles of patients in the study group, from hospital admission to discharge.

increases the potential for other, unrelated, causes of acute renal injury (such as sepsis during hospitalization) to affect the results.¹ In addition, Goldberg et al⁶ did not routinely collect creatinine measurements after the third day of hospitalization, which may have led to an underestimation of acute kidney injury.¹ Goldberg et al⁶ reported a higher incidence of kidney problems than was found in the current study; thus, they underestimated kidney injury. They collected additional creatinine measurements – and therefore they potentially identified more kidney injuries – so actually their results are even more different from this study than they already seem.

Among elderly patients discharged after hospitalization for acute myocardial infarction, small changes in serum creatinine levels have previously been associated with a higher risk of end-stage renal disease and death.⁷ Unfortunately, as in the Goldberg et al study,⁶ the authors of this work did not use AKIN criteria for diagnosing acute kidney injury.¹ Thus, it is difficult to compare these previous results with those found in the current study. Further, while all patients in the current study underwent an immediate coronary angiography and tended to be elderly, neither of these conditions was true in the research performed by

Figure 3 Serum creatinine profiles of patients in the control group, from admission to discharge from hospital.

Figure 4 Stages of chronic renal failure in the study (A) and control (B) groups, from admission to discharge from hospital.

Newsome et al.⁷ Another group evaluated the clinical and prognostic relevance of acute kidney injury (defined as a 25% increase over baseline serum creatinine levels) in STEMI complicated by cardiogenic shock. In their study, 55% of patients developed acute kidney injury during their stay in the intensive care unit.⁸ Again, it is difficult to compare these results to those of the current study, as AKIN criteria were not used to diagnose acute kidney injury.¹ Additionally, cardiogenic shock was not observed in any of the patients studied here.

Rather than employing AKIN criteria, Parikh et al defined acute kidney injury as an absolute change in serum creatinine levels;⁹ they classified the resulting injury as mild (0.3–0.4 mg/dL), moderate (0.5–0.9 mg/dL), or severe (>1.0 mg/dL). Using these criteria, they identified acute kidney injury in 19.4% (7.1% mild, 7.1% moderate, 5.2% severe) of their patients.⁹ However, the validity of these results has been questioned because the left ventricular ejection fraction was unavailable in 39.4% of the patients without acute kidney injury; multivariate analysis has

demonstrated that the ventricular ejection fraction is a strong predictor of long-term survival after an acute myocardial infarction.^{1,10} In the present investigation, heart failure was more commonly observed in patients with both acute kidney injury and acute coronary heart disease.

Acute kidney injury increases morbidity and mortality, particularly in patients who are critically ill.¹¹ Recognition of this has led to recent efforts (eg, AKIN and RIFLE¹² [risk, injury, failure, loss, and end-stage renal failure]) to standardize definitions of acute kidney injury; one particularly important goal has been the identification of biomarkers of early kidney injury, as these should facilitate early interventions and prevention of progression to the acute stage of kidney injury.¹⁰ Unfortunately, there are currently no data comparing the use of RIFLE and AKIN criteria in patients with acute coronary syndrome or myocardial infarction; further, it is difficult to compare these classification schemes, since the RIFLE criteria, unlike those in the AKIN classification, involve five stages.¹ It is also important to note that, while the RIFLE and AKIN criteria are easy to use in a variety of clinical and research settings, they have several limitations.¹¹ For instance, both sets of criteria focus on increases in serum creatinine levels from a hypothetical baseline value, in conjunction with a decrease in urine output. However, these surrogate markers of renal impairment manifest relatively late, after injury has occurred; further, they do not reflect the nature or site of the kidney injury.¹²

Conclusion

This study shows that acute kidney injury occurs relatively frequently in patients with coronary heart disease, but that the association between these diseases is not statistically significant. However, physicians should still monitor their patients – especially those with angina pectoris and myocardial infarction – for acute kidney injury after the administration of contrast medium.

Disclosure

The author reports no conflicts of interest in this work.

References

- Rodrigues FB, Bruetto RG, Torres US, et al. Effect of kidney disease on acute coronary syndrome. *Clin J Am Soc Nephrol*. 2010;5:1530–1536.
- Bellomo R, Ronco C, Kellum JA, et al. Acute renal failure – definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care*. 2004;8:R204–R212.
- Hahn JM. *Checkliste Innere Medizin*. 4th ed. Stuttgart: Thieme;2003.
- Levey AS, Coresh J, Balk E et al. National Kidney Foundation. National Kidney Foundation practice guidelines for chronic kidney disease evaluation, classification, and stratification. *Ann Intern Med*. 2003;139:137–147.
- Levine GN, Bates ER, Blankenship JC et al. 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *Circulation*. 2011;124:2574–609.
- Goldberg A, Hammerman H, Petcherski S, et al. In-hospital and 1-year mortality of patients who develop worsening renal function following acute ST-elevation myocardial infarction. *Am Heart J*. 2005;150:330–337.
- Newsome BB, Warnock DG, McClellan WM, et al. Long-term risk of mortality and end-stage renal disease among the elderly after small increases in serum creatinine level during hospitalization for acute myocardial infarction. *Arch Intern Med*. 2008;168:609–616.
- Marenzi G, Assanelli E, Campodonico J, et al. Acute kidney injury in ST-segment elevation acute myocardial infarction complicated by cardiogenic shock at admission. *Crit Care Med*. 2010;38:438–444.
- Parikh CR, Coca SG, Wang Y, et al. Long-term prognosis of acute kidney injury after acute myocardial infarction. *Arch Intern Med*. 2008;168:987–995.
- Bouzas-Mosquera A, Vázquez-Rodríguez JM, Peteiro J, et al. Acute kidney injury and long-term prognosis after acute myocardial infarction. *Arch Intern Med*. 2009;169:87.
- Choudhury D. Acute kidney injury: current perspectives. *Postgrad Med*. 2010;122:29–40.
- Ricci Z, Cruz DN, Ronco C. Classification and staging of acute kidney injury: beyond the RIFLE and AKIN criteria. *Nat Rev Nephrol*. 2011;7:201–208.

International Journal of General Medicine

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-general-medicine-journal>

Dovepress

A key focus is the elucidation of disease processes and management protocols resulting in improved outcomes for the patient. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.