Mitochondrial K_{ATP} channels in skeletal muscle: are protein kinases C and G, and nitric oxide synthase involved in the fatigue process?

Elizabeth Sánchez-Duarte¹
Xóchitl Trujillo¹
Miguel Huerta¹
Mónica Ortiz-Mesina¹
Christian Cortés-Rojó²
Salvador Manzo-Ávalos²
Alfredo Saavedra-Molina²
Rocío Montoya-Pérez³

¹Unidad de Investigación Enrico Stefani Bonfanti, Centro Universitario de Investigaciones Biomédicas, Universidad de Colima, Colima, México; ²Instituto de Investigaciones Químico-Biológicas, ³Coordinación General de Estudios de Posgrado Universidad Michoacana de San Nicolás de Hidalgo, Morelia, Michoacán, México

Background: Fatigue in skeletal muscle is defined as a reduction in the physical power needed to execute a function or as an inability to maintain mitochondrial ATP production. The mitochondrial potassium channel (mito K_{ATP}) participates in combating fatigue in skeletal muscle. In this work, we evaluated the role of the mito K_{ATP} channel activator (diazoxide) and inhibitors of the signaling routes (protein kinase C, staurosporine; protein kinase G, KT5823; and nitric oxide synthase, metil N^G-Nitro-L-arginine ester, L-NAME), on muscle fatigue tension. In addition, we evaluated the main signaling routes used by the nitric oxide synthase protein and protein kinase C and G, in the presence of their specific activators.

Methods: We used the anterior latissimus dorsi skeletal muscle of 2–3-week-old chicks. This muscle consists of slow muscle fibers. Tension was achieved by applying repetitive electrical stimulation that induced fatigue in an in vitro model.

Results: Diazoxide significantly reduced muscle fatigue ($P = 0.0002$ in peak tension, $P = 0.000002$ in maximum tension) by increasing post-fatigue tension, in spite of the fact that 5-hydroxydecanoate, a selective inhibitor of mito K_{ATP} did not suppress post-fatigue tension.

Conclusion: Our results suggest a lack of direct interaction in inhibition of the signaling routes during fatigue-induced mito K_{ATP} activation. This effect is possibly due to the type of skeletal muscle fibers (slow), the stimulation protocols (twitch), and the animal (avian) model used in the study.

Keywords: fatigue, skeletal muscle, mitochondrial ATP-sensitive potassium channels

Introduction

ATP-dependent potassium channels (K_{ATP}) were described in patches or membrane derived from guinea pig ventricular myocytes.¹ These channels have been reported in several tissues, including β -pancreatic cells,² skeletal muscle fibers of amphibians,³ mammals,^{4–6} and birds,^{7,8} and in the liver-derived internal membrane mitochondria of the rat.⁹ K_{ATP} channels connect cell excitability with its metabolism and play an important role in several cellular functions by detecting the intracellular ATP/ADP relationship.¹⁰

Mitochondrial K_{ATP} (mito K_{ATP}) has been isolated and partially purified, and its function in mitochondrial swelling has been demonstrated.^{9,11} It is composed of a sulfonylurea receptor, coupled with an inward-rectifying potassium channel (Kir 6.x), and configured in a stoichiometry of 4:4.¹² The skeletal muscle K_{ATP} channel phenotype varies in its molecular composition, biophysical properties, and pharmacological response depending on the type of muscle.^{5,6} Garlid et al¹³ demonstrated that mito K_{ATP} activation by diazoxide, a specific activator, was greater than sarcolemmal K_{ATP} activation.

Correspondence: Rocío Montoya-Pérez
Coordinación General de Estudios de Posgrado, Universidad Michoacana de San Nicolás de Hidalgo, Francisco J Mújica s/n Ciudad Universitaria, Edif C5, Col Felicitas del Río 58030, Morelia, Michoacán, México
Tel +52 44 3322 3500 ext 4157
Email biochio@yahoo.com.mx

Some of the activators or inhibitors of the mitoK_{ATP} channel affect the activity of protein kinase C,¹⁴ such as 5-hydroxydecanoate, chelerythrine, or calphostine C, which inhibit the beneficial effect of diazoxide on pretreated mitochondria during preconditioning to ischemia.¹⁵ Protein kinase C modifies excitation-contraction coupling by phosphorylation of Ca²⁺ L-type channels and/or ryanodine receptors,¹⁶ whereas activation of mitoK_{ATP} activates the downstream protein kinase C pathway, leading to production of reactive oxygen species.¹⁶ These signaling routes could be involved in the activity of mitoK_{ATP} in relation to muscle fatigue, because there is a functional level of interaction between mitoK_{ATP} and protein kinase C.¹⁷ Direct phosphorylation of the channel results in a conformational change that opens the channel and fluxes K⁺ to the mitochondria. Costa et al¹⁸ and Costa and Garlid¹⁹ reported on protein kinase G activation and mitoK_{ATP} channel opening, suggesting that there was an increase in production of reactive oxygen species. Light scattering measurements in mitochondria swelling tests indicate that cGMP-dependent protein kinase (protein kinase G) induces a similar opening in mitoK_{ATP} as do K_{ATP} activators (diazoxide and cromakalim) in the heart, liver, and brain-derived mitochondria. Costa et al¹⁸ also reported inhibition of this effect by the K_{ATP} inhibitor (5-hydroxydecanoate), the lipophilic quaternary ion (tetraphenyl phosphonium), sulfonylurea (glibenclamide), and the protein kinase G inhibitor (KT5823). These results suggest that protein kinase G could be the cytosolic component of the mitoK_{ATP} activation route, transmitting the protection signal from the cytosol to the inner mitochondrial membrane through a route that involves one of the kinase proteins (the PKC-ε).¹⁹

On the other hand, there is evidence indicating direct mitoK_{ATP} activation by nitric oxide,^{20,21} and it has been suggested that protein kinase C and nitric oxide synthase are intermediaries in the signaling route that activates mitoK_{ATP} in fatigue-induced mammalian skeletal muscle.²² Therefore, we conducted a study using an *in vitro* model to determine the participation of mitoK_{ATP} in fatigue of avian slow skeletal muscle fibers using activators and inhibitors of this channel and evaluated the proposed signalization routes related to this channel.

Materials and methods

Ethical approval

Chickens were used in accordance with the Institute for Laboratory Animal Research²³ Guide for the Care and Use of Laboratory Animals and the Ethics Committee of the Centro Universitario de Investigaciones Biomédicas of

the Universidad de Colima, which approved the protocol. To minimize animal pain and distress and to perform muscle extraction, the chicks were anesthetized beforehand with chloroform, followed by cervical dislocation and decapitation, ensuring a fast and complete separation of the head from the body, according to the American Veterinary Medical Association guidelines on euthanasia.²⁴

Dissection

The anterior latissimus dorsi muscle was used for this investigation because it is exclusively composed of slow muscle fibers. The muscles were carefully dissected out, together with a piece of the humerus bone and a portion of the proximal tendon, from 2–3-week-old chicks. The spinal cord was pinned to the bottom of the experimental chamber and 3.0 surgical silk threads were tied around the humerus bone to attach the distal end of the muscle bundle (1–2 mm thick) to a force transducer (Grass FT03, West Warwick, RI) by means of a lightweight wire. To record the force, a mechanical transducer was wired to an amplifier (Cyberamp 320; Axon Instruments, Foster City, CA), and connected to an analog-to-digital converter (Digidata 1322A; Axon Instruments) with a 5 Hz sampling rate. Data were acquired using the Axoscope subroutine (pClamp 9.2; Axon Instruments) on a desktop computer.

Fatigue protocols

We used different electrical stimulation frequencies to produce twitches to establish a fatigue protocol in the slow latissimus dorsi muscle, as previously described.²⁵ In short, supramaximal twitches at 300 ms and 0.2 Hz (Grass S-48 stimulator and SIU-5B stimuli isolation unit) were applied until bundle contractions reduced the control contraction by approximately 50% (at about 60 minutes). Pulses were delivered across platinum wire electrodes situated on both sides of the muscle. Each protocol consisted of an initial stimulation period to induce fatigue (about 50% of initial tension), after which drugs were applied to the bath for 6 minutes to maintaining stimulation. Drug withdrawal was then carried out by passing normal Ginsborg saline²⁶ through the muscle and continuing electrical stimulation for 15 minutes to record post-fatigue tension.

Solutions

The muscles were immersed in normal Ginsborg saline^{26,27} composed (in mM) of: 167 NaCl, 5 KCl, 2 MgCl₂, 5 CaCl₂, and 2 imidazole-chloride, with pH adjusted to 7.4. Finally, normal saline containing dextrose 2 g/L was added.

The solutions contained a specific inhibitor of the mitoK_{ATP} channel (5-hydroxydecanoate, 500 μM ; Sigma, St Louis, MO), the mitoK_{ATP} channel activator, diazoxide (10, 30, and 100 μM ; Sigma), a potent selective inhibitor of the cGMP-dependent protein kinase (KT5823, 1 μM ; Sigma), a nitric synthase oxide protein inhibitor (N ω -Nitro-L-arginine methyl ester hydrochloride, L-NAME, 100 μM ; Tocris, Bristol, UK), and a protein kinase C inhibitor (staurosporine, 1 μM ; Calbiochem, La Jolla, CA). All probe solutions were prepared by adding the correct volume of a stock solution dissolved in water or in dimethylsulfoxide (0.1% v/v for diazoxide and staurosporine) to the saline bath. Solution exchange was carried out via a three-way tap located at one end of the central channel of the experimental chamber.

Statistical analysis

Analysis of the experimental data took into consideration the following twitch parameters: amplitude to twitch peak tension expressed in g (from the basal resting tension to the maximal tension), and total tension (area under the tension-time curve). The control twitch was used as 100% and the probe twitch was expressed as the percentage in relation to the control. Twitch was obtained by electrical stimulation (see above).

Tension was measured as the maximum tension from the basal line to the peak before, during, and after the addition of the experimental drug (peak tension), and the tension-time integral was obtained from the area under the twitch profile. We compared each experimental condition. Data analyses were carried out using the Clampfit subroutine of the pClamp 9.2 software (Axon Instruments) and graphs were elaborated using Sigmaplot 10.0 software (Systat Software Inc. Erkrath, Germany). Results were expressed as the mean $\pm$ standard error of the mean, followed by the *n* value. Comparison of means was done using the Student's *t*-test, accepting a significant effect when *P* was <0.05 .

Results

Diazoxide increases twitch tension of fatigued slow skeletal muscle fibers

To determine the possible participation of mitoK_{ATP} during post-fatigue tension in the slow skeletal muscle of the chicken, the effect of diazoxide (10, 30, and 100 μM), considered to be a selective activator of the mitoK_{ATP} channel, was explored (Figure 1). Diazoxide at 30 μM strongly increased the tension of the fatigued muscle.

Figure 2 shows that basal tension (considered as 100%) was reduced to $33.71\% \pm 8.62\%$ in peak tension during fatigue; nevertheless, in the presence of diazoxide 30 μM ,

Figure 1 Normalized tension of the control twitch.

Notes: The third and fourth columns in the graph correspond to the peak and total tension of the twitch in fatigued muscle. The following columns show the post-fatigue effect of diazoxide (10, 30, and 100 μM). At 10 and 30 μM , diazoxide increased the post-fatigue tension in both peak tension and total tension. Normalization was done to compare tension in percentages for different bundles, given that each bundle possesses a different number of muscle fibers.

tension was increased to $103.38\% \pm 7.97\%$ indicating post-fatigue (peak tension). Figure 2B also shows the effect of diazoxide on total tension, which increased the post-fatigue tension to $163.82\% \pm 11.42\%$ in relation to fatigued muscle tension ($n = 4$; $P < 0.05$). These results are consistent with previous reports in which diazoxide antagonizes fatigue.

Figure 2 Effects of diazoxide 30 μM on twitch at almost 60% of muscle fatigue. (A) Representative recorded tension at the beginning of the experiment is represented by (a). In trace (b), the recorded tension approximately one hour after stimulating the bundle at 0.2 Hz is shown. Trace (c) displays the increase in the tension of the fatigued muscle 6 minutes after application of diazoxide. Trace (d) shows the tension recorded 10 minutes after withdrawal of diazoxide. (B) Effects of diazoxide on single twitch post-fatigue tension.

Notes: Diazoxide produced a significant increase in the total and maximal tension of twitches in fatigued muscle fibers ($n = 4$). * $P < 0.05$.

Effect of 5-hydroxydecanoate on twitch tension in fatigued slow skeletal muscle fibers

To determine the possible participation of $\text{mitoK}_{\text{ATP}}$ in post-fatigue tension, the action of the $\text{mitoK}_{\text{ATP}}$ channel-selective inhibitor, 5-hydroxydecanoate, was studied at the previously reported concentration range (500 μM).^{22,28} The maximum tension obtained prior to adding the drug was the control tension (100%). A 50% reduction in control tension was considered to be fatigue. The 5-hydroxydecanoate was added once fatigue was reached, and tension was recovered at $11.91\% \pm 1.6\%$ in relation to muscle fatigue tension. We observed a tension recovery ($15.3\% \pm 4.8\%$) in the presence of 5-hydroxydecanoate. Nevertheless, this difference was not statistically significant ($n = 4$; $P > 0.05$).

Effect of diazoxide and 5-hydroxydecanoate on tension in slow muscle fibers

Previous reports have shown that diazoxide antagonizes fatigue by opening $\text{mitoK}_{\text{ATP}}$ channels, which was corroborated by adding 5-hydroxydecanoate and blocking the effect of diazoxide.²² Figure 3 shows the results when diazoxide 30 μM and 5-hydroxydecanoate 500 μM were added. According to these results, there was an increase by $5.91\% \pm 4.99\%$ with respect to control (standardized 100%). However, there was an increase of

$26.0\% \pm 4.99\%$ in peak tension, whereas the total tension increment was $44.87\% \pm 7.90\%$ with respect to fatigue tension; nevertheless, the effect was statistically significant ($P = 0.025$ and $P = 0.023$, respectively). These results suggest that 5-hydroxydecanoate in combination with diazoxide acts as an agonist during fatigue, although it is possible that 5-hydroxydecanoate does not act as a selective inhibitor of $\text{mitoK}_{\text{ATP}}$ in the slow skeletal muscle fibers of the chicken, due to substrate conversion by β -oxidation.²⁹

Effect of diazoxide and staurosporine on tension in slow muscle fibers

Previous studies have implicated protein kinase C in $\text{mitoK}_{\text{ATP}}$ channel function. García et al²² suggested that diazoxide has an antagonistic effect on muscle fatigue through several signaling routes. We studied the action of staurosporine (1 μM), a selective inhibitor for protein kinase C, in combination with diazoxide. Diazoxide and staurosporine caused an increase in post-fatigue tension in relation to muscle fatigue tension when added to the bath (Figure 4). The maximum tension (control) dropped to $75.77\% \pm 2.47\%$ in fatigued muscle. Nevertheless, there was an increase in peak tension of $120.75\% \pm 3.54\%$ (post-fatigue) and in total tension of $158.82\% \pm 8.60\%$. Both of these increases were statistically significant ($P = 0.007$ and $P = 0.021$, respectively). These results differ from those previously reported by García et al,²² in which the effect of

Figure 3 Effect of the combination of 5-hydroxydecanoate 500 μM and diazoxide 30 μM on the fatigued skeletal muscle tension, along with peak tension ($P = 0.023$) and total tension ($P < 0.05$).

Note: *Represents statistical significance. There is a significant increase in post-fatigue tension after application of both agents. First columns indicate controls standardized to 100%, second columns indicate a comparison of reduction of force from controls and third columns indicate the effect of used drugs compared against fatigue as 100%.

Figure 4 Effect of diazoxide and staurosporine on maximum and total tension.

Notes: *Represents statistical significance. The combination of drugs produced a great increase in post-fatigue tension in comparison with the tension of fatigued muscle without the drugs. Data correspond with those from other bundles following the same protocol. First columns indicate controls standardized to 100%, second columns indicate a comparison of reduction of force from controls and third columns indicate the effect of used drugs compared against fatigue.

diazoxide on post-fatigue tension was stronger when protein kinase C was blocked with chelerythrine and the two drugs were combined.

Effect of diazoxide and KT5823 on tension in slow muscle fibers

Previous reports have indicated that protein kinase G induces opening of the mito K_{ATP} channel in a manner similar to diazoxide. In addition, blockade of the channel by selective inhibition of protein kinase G with KT5823 has been reported.¹⁹ To determine the possible participation of mito K_{ATP} during post-fatigue tension via this mechanism, KT5823 was used in combination with diazoxide. The combination of KT5823 1 μ M with diazoxide resulted in an increase in post-fatigue tension with respect to the effect of diazoxide alone (Figure 5). The increment in maximal tension was $16.17\% \pm 6.04\%$, whereas the increase in total tension was $63.88\% \pm 4.74\%$. Both of these increases in tension were statistically significant ($P = 0.002$ and $P = 0.0001$, respectively). Our results show that KT5823 did not antagonize the effect of diazoxide, therefore promoting an increase in post-fatigue tension.

Effect of diazoxide and L-NAME on post-fatigue tension in slow muscle fibers

The effect of the combination of L-NAME, a selective nitric oxide synthase inhibitor, and diazoxide on mito K_{ATP}

Figure 5 Effect of diazoxide and KT5823 on post-fatigued tension (maximum and total).

Notes: *Represents statistical significance. The combination of both drugs increased post-fatigue tension. Data correspond with those from the other bundles following the same protocol. First columns indicate controls standardized to 100%, second columns indicate a comparison of the reduction of the force respecting controls, and third columns indicate the effect of the used drugs compared against fatigue tension.

Figure 6 Effect of diazoxide and metil N^G-Nitro-L-arginine ester on the post-fatigue tension (maximum and total tension).

Note: *Represents statistical significance. The figure illustrates the tension twitches in different electrically stimulated bundles after the application of this drug combination, which significantly increased post-fatigue tension. First columns indicate controls standardized to 100%, second columns indicate a comparison of the reduction of the force respecting controls, and third columns indicate the effect of the used drugs compared against fatigue tension.

channels was investigated.²² Figure 6 shows the effect of diazoxide 30 μ M and L-NAME 100 μ M on maximum and total tension. Comparing the average for muscle fatigue in relation to the control (100%), there was a decrease of $69.35\% \pm 5.18\%$, whereas the combination of diazoxide and L-NAME increased post-fatigue tension to $62.31\% \pm 12.38\%$ ($P = 0.0003$) and produced a higher increase in total tension ($91.22\% \pm 15.06\%$ with respect to fatigue, $P = 0.0004$). García et al²² reported inhibition when diazoxide plus L-NAME was present. However, our results showed just the opposite, ie, there was a remarkable increment in post-fatigue tension with a combination of the two drugs, and it was even higher with diazoxide alone.

Discussion

The ADP/ATP relationship is a reflection of the metabolic necessities of the cell. Therefore, a decrease in ATP has an important role in the development of muscle fatigue, and any event that alters the production of this metabolite can determine the tendency of the muscle to become fatigued.³⁰

In the present work, the effects of mito K_{ATP} channel agonists and antagonists were analyzed to clarify their possible participation during the fatigue process in slow skeletal muscle fibers of the chicken. Previously, García et al²² reported activation of mito K_{ATP} during fatigue in fast ELD skeletal mammalian muscle. More recently, we suggested

the participation of K_{ATP} channels in avian muscle (anterior latissimus dorsi),²⁵ composed mainly of slow muscle fibers (type I).³¹

It has been reported that diazoxide 100 μ M increased post-fatigue tension, and it was proposed that $mitoK_{ATP}$ has a dominant role in this action. In the proposed analysis, diazoxide reduced fatigue without affecting twitches and/or tetanus, which suggests a direct influence on $mitoK_{ATP}$ discarding any effect on sarcolemmal K_{ATP} in addition to the specificity previously reported for this channel in isolated heart mitochondria.¹³ It is important to point out that a diazoxide concentration was used and reported to have the maximum effects in these experiments.²⁸ In our results, diazoxide 30 μ M had a statistically significant effect on post-fatigue tension (Figure 1), consistent with the results of García et al²² but at a lower dose.

Activation of $mitoK_{ATP}$ channels reconstituted in liposomes or in rat skeletal muscle bundles is inhibited by 5-hydroxydecanoate.^{13,32} In contrast, we did not observe any effect on post-fatigue tension using 5-hydroxydecanoate 500 μ M (data not shown). Perhaps this is because an intracellular mediator^{13,29} or an activator could be required in order to guarantee $mitoK_{ATP}$ activation.²² Nevertheless, it is questionable to consider 5-hydroxydecanoate as a selective inhibitor because it is a potential metabolite for β -oxidation.²⁹ Recently, we have reported no modification in the respiratory rate of mitochondria from chicken skeletal muscle as a result of treatment with 5-hydroxydecanoate in relation to different respiratory substrates.³² In this regard, Sarre et al³³ suggested the existence of other factors involved in the ischemic process implied during fatigue, upstream activation of protein kinase C, and inhibiting diazoxide activation of $mitoK_{ATP}$ with chelerythrine. Costa et al¹⁸ and Costa and Garlid¹⁹ suggested that protein kinase G induces the opening of $mitoK_{ATP}$ in a way similar to that achieved by the channel activators, diazoxide and cromakalim, in heart, liver, and brain mitochondria. Furthermore, they verified the suggested route through inhibition of channel opening by KT5823, a selective inhibitor of protein kinase G. García et al²² proposed that protein kinase C and mitochondrial nitric oxide synthase act as intermediaries in the signaling route where $mitoK_{ATP}$ is involved. Another proposed factor for $mitoK_{ATP}$ activation is production of reactive oxygen species due to nitric oxide production and protein kinase C activation.

Our present results differ from those previously described for the heart and fast skeletal muscle of the rat. Nevertheless, it is important to mention that we worked with a different

in vitro model of fatigue (avian slow skeletal muscle fibers) and with a different stimulation protocol (twitches). In our protocol, we successfully induced fatigue, and we considered that the fatigued muscle kept up its strength under ischemic conditions.^{34,35} Protein kinase C has been reported to have a central role during ischemic preconditioning.¹⁸

According to previous studies, staurosporine inhibits the protective effect of ischemic preconditioning via protein kinase C inhibition in the rat heart.³⁶ Chelerythrine, a protein kinase C inhibitor, eliminates the inhibitory effect of diazoxide in fatigue.²² In the present report, the combination of staurosporine with diazoxide showed an increase in post-fatigue tension (Figure 4). Simultaneously, we used a selective inhibitor of protein kinase G (KT5823) to confirm whether protein kinase G takes part in the upstream mitochondrial signaling route of $mitoK_{ATP}$ through its phosphorylation.^{18,19} Thus, we observed an increase in post-fatigue tension. These results do not concur with those reported for the rat heart (Figure 5). We suggest that KT5823 could not have effects on protein kinase G due to the experimental model (complete muscle) used. This is why it is necessary to perform experiments on single fibers and isolated mitochondria of skeletal muscle to confirm that KT5823 has an effect on protein kinase G in skeletal muscle.

In addition, the effects of the nitric oxide synthase inhibitor, L-NAME, were analyzed to determine whether $mitoK_{ATP}$ channels are involved in the effect of diazoxide on the signaling cascade that involves nitric oxide. Nitric oxide, an important messenger molecule in cells, is formed during skeletal muscle activity³⁷ and regulates $mitoK_{ATP}$ function.²⁸ L-NAME blocked the cardioprotective effects of diazoxide.³⁸ There is evidence indicating that nitric oxide directly activates $mitoK_{ATP}$.^{21,39} García et al²² maintain that a combination of diazoxide with L-NAME does not have any effect on post-fatigue tension, which could indicate that L-NAME does indeed inhibit fatigue. In our case, we observed an increase in the presence of both drugs. It is possible that $mitoK_{ATP}$ could be activated by mitochondrial nitric oxide synthase,³³ a place the nitric oxide synthase inhibitor would not have access to.

In conclusion, the effects of K_{ATP} channel inhibitors on muscle fatigue are controversial. In some studies, no significant effects on fatigue have been found,^{40,41} while others have reported increases in fatigue.⁴² Further studies are needed in order to understand better the pharmacological properties of this type of channel in this type of tissue, because they could be the key to identifying the cellular mechanisms involved in slow muscle fiber fatigue. The importance of this study involves the use of a model of

fatigue composed of exclusively of slow muscle fibers, ie, the anterior latissimus dorsi of the chicken.

Acknowledgments

We wish to thank Humberto Cruzblanca for his kind donation of the staurosporine used in this study. ESD has a scholarship from CONACyT, México. This paper was partially financed by grant RMP-PROMEP-PTC311.

Disclosure

The authors declare they have no conflict of interests in this work.

References

- Noma A. ATP-regulated K^+ channels in cardiac muscle. *Nature*. 1983;305:147–148.
- Cook DL, Hales CN. Intracellular ATP directly blocks K^+ channels in pancreatic β -cells. *Nature*. 1984;311:271–273.
- Spruce AE, Standen NB, Stanfield PR. Voltage-dependent ATP-sensitive potassium channels of skeletal muscle membrane. *Nature*. 1985;316:736–738.
- Burton F, Dorstelmann U, Hutter OF. Single-channel activity in sarcolemmal vesicles from human and other mammalian muscles. *Muscle Nerve*. 1988;11:1029–1038.
- Tricarico D, Mele A, Camerino GM, et al. The K_{ATP} channel is a molecular sensor of atrophy in skeletal muscle. *J Physiol*. 2010;588:773–784.
- Tricarico D, Mele A, Lundquist AL, et al. Hybrid assemblies of ATP-sensitive K^+ channels determine their muscle-type-dependent biophysical and pharmacological properties. *Proc Natl Acad Sci U S A*. 2006;103:1118–1123.
- Thomas SA, Hume RI. Single potassium channel currents activated by extracellular ATP in developing chick skeletal muscle: a role for second messengers. *J Neurophysiol*. 1993;69:1556–1566.
- Fosset M, Allard B, Lazdunski M. Coexistence of two classes of glibenclamide-inhibitable ATP-regulated K^+ channels in avian skeletal muscle. *Pflugers Arch*. 1995;431:117–124.
- Inoue I, Nagase H, Kishi K, et al. ATP-sensitive K^+ channel in the mitochondrial inner membrane. *Nature*. 1991;352:244–247.
- Inagaki N, Seino S. ATP-sensitive potassium channels: structures, functions, and pathophysiology. *Jpn J Physiol*. 1998;48:397–412.
- Paucek P, Mironova G, Mahdi F, et al. Reconstitution and partial purification of the glibenclamide-sensitive, ATP-dependent K^+ channel from rat liver and beef heart mitochondria. *J Biol Chem*. 1992;267:26062–26069.
- Barrett-Jolley R, McPherson GA. Characterization of K_{ATP} channels in intact mammalian skeletal muscle fibres. *Br J Pharmacol*. 1998;123:1103–1110.
- Garlid KD, Paucek P, Yarov-Yarovoy V, et al. Cardioprotective effect of diazoxide and its interaction with mitochondrial ATP-sensitive K^+ channels. *Circ Res*. 1997;81:1072–1082.
- Wang Y, Ashraf M. Activation of mitochondrial ATP-sensitive K^+ channel for cardiac protection against ischemic injury is dependent on protein kinase C activity. *Circ Res*. 1999;84:1156–1165.
- Wang Y, Ashraf M. Role of protein kinase C in mitochondrial K_{ATP} channel-mediated protection against Ca^{2+} overloads injury in rat myocardium. *Circ Res*. 1999;84:1156–1165.
- Rybin VO, Steinberg SF. Protein kinase C isoform expression and regulation in the developing rat heart. *Circ Res*. 1994;74:299–309.
- Jaburek M, Costa ADT, Burton JR, et al. Mitochondrial PKC ϵ and mito K_{ATP} copurify and core constitutes to form a functioning signaling module in proteoliposomes. *Circ Res*. 2006;99:878–883.
- Costa AD, Garlid KD, West IC, et al. Protein kinase G transmits the cardioprotective signal from cytosol to mitochondria. *Circ Res*. 2005;97:329–336.
- Costa AD, Garlid KD. Intramitochondrial signaling: interactions among mito K_{ATP} , PKC ϵ , ROS, and MPT. *Am J Physiol Heart Circ Physiol*. 2008;295:H874–H882.
- Sasaki N, Sato T, Ohler A, et al. Activation of mitochondrial ATP-dependent potassium channels by nitric oxide. *Circulation*. 2000;101:439–445.
- Ljubkovic M, Shi Y, Cheng Q, et al. Cardiac mitochondrial ATP-sensitive potassium channel is activated by nitric oxide in vitro. *FEBS Lett*. 2007;581:4255–4259.
- García MC, Hernández A, Sánchez JA. Role of mitochondrial ATP-sensitive potassium channels on fatigue in mouse muscle fibers. *Biochem Biophys Res Commun*. 2009;385:28–32.
- Institute for Laboratory Animal Research. *Guide for the Care and Use of Laboratory Animals*. National Research Council. Washington, DC: National Academic Press; 1996.
- American Veterinary Medical Association. Guidelines on Euthanasia Formerly Report of the AVMA Panel on Euthanasia, 2007. Available from: http://www.avma.org/issues/animal_welfare/euthanasia.pdf. Accessed May 2, 2012.
- Andrade MF, Trujillo X, Sánchez E, et al. Glibenclamide increases post-fatigue tension in slow skeletal muscle fibers of the chicken. *J Comp Physiol B*. 2011;181:403–412.
- Ginsborg BL. Some properties of avian skeletal muscle fibres with multiple neuromuscular junctions. *J Physiol*. 1960;154:581–598.
- Page SG. Structure and some contractile properties of fast and slow muscles of the chicken. *J Physiol*. 1969;205:131–145.
- Debska G, Kicinska A, Skalska J, et al. Opening of potassium channels modulates mitochondrial function in rat skeletal muscle. *Biochim Biophys Acta*. 2002;1556:97–105.
- Hanley PJ, Drose S, Brandt U, et al. 5-Hydroxydecanoate is metabolized in mitochondria and creates a rate-limiting bottleneck for beta-oxidation of fatty acids. *J Physiol (Lond)*. 2005;562:307–318.
- Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: mechanisms. *Physiol Rev*. 2008;88:287–332.
- Burke RE, Levine DN, Zajac FE. Mammalian motor units: physiological-histochemical correlation in three types in cat gastrocnemius. *Science*. 1971;174:709–712.
- Montoya-Pérez R, Saavedra-Molina A, Trujillo X, et al. Inhibition of oxygen consumption in skeletal muscle-derived mitochondria by pinacidil, diazoxide, and glibenclamide, but not by 5-hydroxydecanoate. *J Bioenerg Biomembr*. 2010;42:21–27.
- Sarre A, Lange N, Kucera P, et al. Mito K_{ATP} channel activation in the postanoxic developing heart protects excitation-contraction coupling via NO-, ROS-, and PKC-dependent pathways. *Am J Physiol Heart Circ Physiol*. 2005;288:H1611–H1619.
- Oxman T, Arad M, Klein R, et al. Limb ischemia preconditions the heart against reperfusion tachyarrhythmia. *Am J Physiol*. 1997;273:H1707–H1712.
- Moses MA, Addison PD, Neligan PC, et al. Mitochondrial K_{ATP} channels in hindlimb remote ischemic preconditioning of skeletal muscle against infarction. *Am J Physiol Heart Circ Physiol*. 2004;288:H559–H567.
- Speechly-Dick ME, Mocanu MM, Yellon DM. Its role in ischemic preconditioning in the rat. *Circ Res*. 1994;75:586–590.
- Silveira LR, Pereira-Da-Silva L, Jue CL, et al. Formation of hydrogen peroxide and nitric oxide in rat skeletal muscle cells during contractions. *Free Radic Biol Med*. 2003;35:455–464.
- Ockaili R, Emani VR, Okubo S, et al. Opening of mitochondrial K_{ATP} channel induces early and delayed cardioprotective effect: role of nitric oxide. *Am J Physiol Heart Circ Physiol*. 1999;277:2425–2434.
- Comtois A, Light P, Renaud JM, et al. Tolbutamide, but not glyburide, affects the excitability and contractility of unfatigued frog sartorius muscle. *Eur J Pharmacol*. 1993;242:65–73.
- Light PE, Comtois A, Renaud JM. The effect of glibenclamide on frog skeletal muscle: evidence for K_{ATP} channel activation during fatigue. *J Physiol*. 1994;475:495–507.

41. Matar W, Nosek TM, Wong D, et al. Pinacidil suppresses contractility and preserves energy but glibenclamide has not effect during muscle fatigue. *Am J Physiol Cell Physiol*. 2000;278:C404–C416.
42. Duty S, Allen DG. The effects of glibenclamide on tetanic force and intracellular calcium in normal and fatigued mouse skeletal muscle. *Exp Physiol*. 1995;80:529–541.

Open Access Animal Physiology

Dovepress

Publish your work in this journal

Open Access Animal Physiology is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of animal physiology. The manuscript management system is completely online and includes a very

quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/open-access-animal-physiology-journal>