Expression of novel tumor markers of pancreatic adenocarcinomas in intrahepatic cholangiocarcinomas

Meijuan Zong¹

Lei Jia¹

Liang Li²

¹Zibo Vocational Institute,

²Department of General Surgery,
Zibo Central Hospital, Zibo, China

Abstract: Intrahepatic cholangiocarcinomas (IHCCs) are morphologically and biologically similar to pancreatic ductal adenocarcinomas (PDACs), so newly identified PDAC-associated genes or proteins could provide clues for screening novel biomarkers for IHCC. In this study, the expression of three novel PDAC tumor markers (T-box transcription factor-4 [TBX4], heat shock protein-60 [HSP60], and Parkinson protein-7 [DJ-1]) identified in previous proteomic studies in IHCC tumors were immunohistochemically detected. The current study confirmed that three novel pancreatic cancer biomarkers TBX4, HSP60, and DJ-1 were also overexpressed in IHCC tumors, but with a relatively lower expression level than PDAC. No significant association was found between tumor marker expression and the clinicopathological characteristics of IHCC patients except that TBX4 expression correlated with tumor grades. Moreover, DJ-1 was demonstrated to be an independent prognostic factor for these patients. The current findings suggest that DJ-1 might play an important role in the malignant progression of IHCC, and its exact mechanism during IHCC progression deserves further investigation.

Keywords: intrahepatic cholangiocarcinomas, biomarker, TBX4, HSP60, DJ-1

Introduction

Intrahepatic cholangiocarcinomas (IHCCs) arising from the epithelia of the intrahepatic bile duct rank second in the primary malignant liver tumors. The incidence and mortality of IHCCs have increased in recent years in China, and this tendency has also been observed in Western countries.¹ Due to the lack of tumor markers for early diagnosis and effective optional therapeutic approaches, the prognosis of IHCC patients is very poor. Therefore, there is a great need to mine novel tumor markers as diagnostic, prognostic, and therapeutic targets for IHCC patients.

IHCCs are morphologically and biologically similar to pancreatic ductal adenocarcinomas (PDACs), most of the tumor markers of PDAC show similar expression patterns in biliary carcinomas.² Proteomics technology has become a powerful and efficient means for mining candidate tumor biomarkers with altered expression patterns in human malignancies. The authors and other groups have recently used multiple proteomic platforms to identify and validate a panel of novel candidate protein tumor markers that are overexpressed in PDAC tissues.^{3,4} As relatively fewer biomarkers have been identified in IHCCs, it was speculated that these PDAC proteins could provide a source for screening novel tumor markers for IHCC.

In a previous study using two-dimensional electrophoresis-based proteomic tools, three candidate biomarkers (T-box transcription factor-4 [TBX4], heat shock protein-60 [HSP60], and Parkinson protein-7 [DJ-1]) for PDAC were identified.³⁻⁶

Correspondence: Meijuan Zong
Zibo Vocational Institute, Liantong Road,
Zibo, Shandong 255314, China
Tel +86 533 282 8157
Fax +86 533 282 8157
Email meijuanzong@163.com

These three proteins were overexpressed in PDAC tumors compared with normal pancreas tissues or benign tumors; furthermore, their expression levels were demonstrated to be associated with clinicopathological traits such as tumor grades. In particular, high expression of TBX4 was also an independent prognostic factor for PDAC.⁵ In this study, the expression of the above three novel PDAC-associated proteins in the IHCC tumors were immunohistochemically assessed in order to validate if these proteins are overexpressed in IHCCs. Their clinicopathological and prognostic significance were also evaluated.

Material and methods

Patients

A total of 72 consecutive IHCC patients who underwent standard curative resection at the Department of General Surgery of Zibo Central Hospital (Zibo, China) and Qilu Hospital (Jinan, China) from 1998 to 2007 were reviewed retrospectively. All the patients routinely received postoperative gemcitabine-based chemotherapy regimens, and no patient received preoperative chemotherapy and/or radiation. The overall survival time of IHCC patients was defined as the time from operation to cancer-caused death. The clinicopathological data of these patients are summarized in Table 1. An additional 55 patients with PDAC who

underwent standard curative resection at the Department of General Surgery of Zibo Central Hospital and Qilu Hospital from 2005 to 2007 were also included in this study. The study was approved by the local research ethics committee at the authors' institution. Informed written consents were obtained from all patients.

Immunohistochemistry

Immunohistochemical staining was performed on 4 µm formalin-fixed, paraffin-embedded archival tissue sections of 72 IHCC primary tumors, their paired adjacent noncancerous tissues, and 55 PDAC tumors. Briefly, deparaffinized and rehydrated sections were boiled in sodium citrate buffer (pH 6.0) to perform antigen retrieval. Endogenous peroxidase activity was blocked by incubating the slides in methanol containing 3% hydrogen peroxide. Sections were incubated overnight at 4°C with the primary antibodies – anti-TBX4, 1:200 (Abcam, Cambridge, MA, USA), anti-HSP60, 1:100 (Santa Cruz Biotechnology, Santa Cruz, CA, USA), and anti-DJ-1, 1:150 (Santa Cruz Biotechnology) – and subsequently developed with an avidin–biotin–peroxidase complex method according to the manufacturer's recommendations (EnVision™ system; Dako Denmark A/S, Glostrup, Denmark). The specificity of antibodies used in this study had been validated in

Table 1 Relationship between tumor markers and clinicopathological characteristics in intrahepatic cholangiocarcinomas

Characteristics	Number of cases	TBX4		P-value	HSP60		P value	DJ-1		P-value
		Low (n = 34)	High (n = 38)		Low (n = 34)	High (n = 38)		Low (n = 33)	High (n = 39)	
Age										
<60 years	54	24	30	0.413	25	29	0.785	28	26	0.076
≥60 years	18	10	8		9	9		5	13	
Gender										
Female	21	8	13	0.320	13	8	0.320	11	10	0.474
Male	51	26	25		25	26		22	29	
Tumor size										
T1–2	25	12	13	0.923	13	12	0.923	14	11	0.207
T3–4	47	22	25		25	22		19	28	
Lymph node metastasis										
Negative	40	21	19	0.316	19	21	0.316	18	22	0.874
Positive	32	13	19		19	13		15	17	
Tumor grade										
G1–2	35	11	24	0.009	22	13	0.096	19	16	0.162
G3	37	23	14		16	21		14	23	
Venous invasion										
Negative	30	14	16	0.936	13	17	0.175	12	18	0.401
Positive	42	20	22		25	17		21	21	
Complication										
No	41	19	22	0.863	21	20	0.761	18	23	0.705
Yes	31	15	16		17	14		15	16	

Abbreviations: DJ-1, Parkinson protein-7; HSP60, heat shock protein-60; TBX4, T-box transcription factor-4.

previous studies.^{3–6} Diaminobenzidine was used as a chromogen, and hematoxylin was used as a counterstain. Nonspecific immunoglobulin G was used as a negative control. The expression level of each tumor marker was measured as high expression when moderate or strong staining was present in more than 25% of the cancer cells, which is accordance with previously established criteria on PDACs.

Statistics

The association of tumor biomarkers with the clinicopathological characteristics of IHCC patients was evaluated with Chi-squared or Fisher's exact test as appropriate. Overall survival in relation to TBX4, HSP60, and DJ-1 expression patterns and other clinicopathological characteristics was evaluated by Kaplan–Meier survival curve and log-rank test. Multivariable analyses were analyzed using a Cox proportional hazards model. Only statistically significant prognostic factors identified by univariate analysis were entered in the multivariate analysis. Statistical analyses were performed using SPSS® 16.0 software for Windows (IBM Corporation, Armonk, NY). The level of statistical significance was set at $P < 0.05$.

Results

The expression of three recently identified PDAC markers (TBX4, HSP60, and DJ-1) in IHCC tissues was determined (Figure 1). TBX4 showed mainly

nuclear staining, and cytoplasmic staining was also frequently observed. Meanwhile, DJ-1 staining was mainly located in the cytoplasm, occasionally nuclear staining was also detected. HSP60 immunostaining was primarily in the cytoplasm and the membrane of neoplastic epithelium in IHCC tumors. The percentage of carcinomas demonstrating either focal or diffuse labeling with these markers varied greatly among IHCC samples. According to established criteria for immunostaining, 52.8%, 52.8%, and 54.2% of the IHCC tumors were defined as positive for TBX4, HSP60, and DJ-1, respectively. PDAC samples presented similar immunostaining patterns for TBX4, HSP60, and DJ-1 in IHCC tumors; however, all of these markers showed lower frequencies of expression in the IHCC samples in comparison to the PDAC samples (69.1% for TBX4, 83.6% for HSP60, and 76.3% for DJ-1). There were also significant differences in the levels of TBX4, HSP60, and DJ-1 expression between neoplastic and adjacent nonneoplastic intrahepatic duct epithelia (2.78% for TBX4, 6.94% for HSP60, and 2.78% for DJ-1) in IHCC tumors.

Because the IHCC tumors showed variable labeling with HSP60, DJ-1, and TBX4, their clinicopathological and prognostic significance were further evaluated. As seen in Table 1, no significant association was identified between tumor markers expression and the clinicopathological characteristics of IHCC patients except that a high level of TBX4 expression had a positive correlation with better differentiation types (grades 1–2). Kaplan–Meier analysis

Figure 1 Representative images of the high expression of TBX4, HSP60, and DJ-1 in IHCC and PDAC tumors, and low or negative expression of them in the normal liver tissues (containing intrahepatic duct) and negative controls (IHCC tumors).

Abbreviations: DJ-1, Parkinson protein-7; HSP60, heat shock protein-60; IHCC, intrahepatic cholangiocarcinoma; PDAC, pancreatic ductal adenocarcinoma; TBX4, T-box transcription factor-4.

revealed a tendency towards shorter overall survival in patients with high DJ-1 expression, but not with TBX4 and HSP60 overexpression (Figure 2). In this study, three factors including tumor size, lymph node metastasis, and DJ-1 expression were identified as significant prognostic factors by univariate analysis. Cox proportional hazard model analysis using these three factors confirmed that DJ-1 expression was an independent predictor for reduced overall survival of IHCC patients (Table 2).

Discussion

IHCCs are similar to PDACs based on the morphological and molecular backgrounds. A panel of tumors such as carbohydrate antigen-19-9, Dupan-2, carbohydrate antigen-50, Span-1, carcinoembryonic antigen, and pancreatic elastase-1 has been used in the diagnosis of both PDACs and IHCCs.² Compared with PDACs, less frequent studies focus on the mining of tumor markers in IHCC. Therefore, it is proposed that previously identified overexpressed genes or proteins in PDACs could provide clues for screening novel biomarkers for IHCC. For example, Swierczynski et al analyzed eight novel PDAC markers (fascin, mucin-4, 14-3-3 σ , prostate stem cell antigen, topoisomerase-II α , and cdc2/p34) in IHCC tumors.⁷ The results of that study indicated that most PDAC markers show similar, yet not identical, expression patterns in biliary carcinomas, suggesting novel PDAC markers are potentially useful in developing diagnostic and therapeutic targets for tumors involving the biliary system.

The current study compared the performance of three novel PDAC markers TBX4, HSP60, and DJ-1 in

Table 2 Univariate and multivariate analysis of overall survival in patients with intrahepatic cholangiocarcinomas

Variables	Univariate		Multivariate	
	HR (95% CI)	P-value	HR (95% CI)	P-value
Age (years)	1.279 (0.729–2.241)	0.391		
Gender	1.059 (0.614–1.826)	0.836		
Tumor size	1.842 (1.083–3.133)	0.024	1.012 (0.494–2.076)	0.973
Lymph node status	2.075 (1.254–3.433)	0.005	2.381 (1.177–4.817)	0.016
Tumor grade	0.978 (0.600–1.593)	0.928		
Venous invasion	1.169 (0.711–1.923)	0.538		
Complication	1.484 (0.913–2.411)	0.111		
TBX4	1.016 (0.623–1.656)	0.951		
HSP60	0.967 (0.593–1.575)	0.892		
DJ-1	1.673 (1.021–2.743)	0.041	1.980 (1.161–3.375)	0.012

Abbreviations: CI, confidence interval; DJ-1, Parkinson protein-7; HR, hazard ratio; HSP60, heat shock protein-60; TBX4, T-box transcription factor-4.

IHCC samples. These PDAC markers were identified in previous proteomic studies, and have not been investigated in IHCC in previous studies. The results indicate that analogous to PDACs, IHCCs demonstrated overexpression of each of the markers examined, but their expression levels were relatively lower than those in PDACs. This trend is similar to that found in previous studies.

In a previous comparative proteomic study on pancreatic cancer and normal pancreas tissues, for the first time TBX4 and HSP60 were identified to be upregulated in PDACs.³ TBX4 belongs to a transcription factor family characterized by a highly conserved DNA-binding motif (T-box) and plays a key role during organogenesis and pattern formation in both vertebrate and invertebrate embryos.^{8,9} HSP60 is a chaperonin protein involved in the folding and assembly of multiple proteins.¹⁰ It was further validated that TBX4 was a differentiation-associated marker for PDAC, and low expression of TBX4 also predicted worse prognosis for patients with PDAC.⁵ However, in this study, no association was observed between these two markers with the clinicopathological traits and prognosis of IHCC patients.

In another quantitative proteomics study on pancreatic juices, DJ-1 was identified to be overexpressed in PDAC tissue and pancreatic juice samples, and it was further confirmed that DJ-1 correlated with tumor differentiation in PDAC.⁴ DJ-1 is a novel mitogen-dependent oncoprotein

Figure 2 Kaplan-Meier survival curves performed according to DJ-1 expression in IHCC patients.

Notes: Follow-up was finished in April 2009. Patients lost to follow-up were censored at the time of their last visit.

Abbreviations: cum, cumulative; DJ-1, Parkinson protein-7; IHCC, intrahepatic cholangiocarcinoma.

involved in Ras-related signal transduction pathway,¹¹ which has been confirmed to be upregulated in multiple cancers.^{12,13} In the current study, it was found that high DJ-1 expression level is an independent prognostic factor for IHCC patients. This finding suggests that DJ-1 might play an important role in the malignant progression of IHCC, and could be used as a potential candidate predictive factor for IHCC patients undergoing surgical resection.

Conclusion

In conclusion, the current study confirmed that three novel pancreatic cancer biomarkers TBX4, HSP60, and DJ-1 were also overexpressed in IHCC tumors, but with a relatively lower expression level than PDACs. DJ-1 might be associated with more progressive behaviors and worse prognosis for IHCC patients; its exact mechanism during IHCC progression deserves further investigation.

Disclosure

The authors report no conflicts of interest in this work.

References

- Shin HR, Oh JK, Masuyer E, et al. Comparison of incidence of intrahepatic and extrahepatic cholangiocarcinoma – focus on East and South-Eastern Asia. *Asian Pac J Cancer Prev*. 2010;11(5):1159–1166.
- Takezako Y, Okusaka T, Ueno H, Ikeda M, Morizane C, Najima M. (Tumor markers for pancreatic and biliary tract cancer). *Gan To Kagaku Ryoho*. 2004;31(9):1443–1446. Japanese.
- Qi T, Han J, Cui Y, Zong M, Liu X, Zhu B. Comparative proteomic analysis for the detection of biomarkers in pancreatic ductal adenocarcinomas. *J Clin Pathol*. 2008;61(1):49–58.
- Tian M, Cui YZ, Song GH, et al. Proteomic analysis identifies MMP-9, DJ-1 and A1BG as overexpressed proteins in pancreatic juice from pancreatic ductal adenocarcinoma patients. *BMC Cancer*. 2008;8:241.
- Zong M, Meng M, Li L. Low expression of TBX4 predicts poor prognosis in patients with stage II pancreatic ductal adenocarcinoma. *Int J Mol Sci*. 2011;12(8):4953–4963.
- Cui Y, Tian M, Zong M, et al. Proteomic analysis of pancreatic ductal adenocarcinoma compared with normal adjacent pancreatic tissue and pancreatic benign cystadenoma. *Pancreatol*. 2009;9(1–2):89–98.
- Swierczynski SL, Maitra A, Abraham SC, et al. Analysis of novel tumor markers in pancreatic and biliary carcinomas using tissue microarrays. *Hum Pathol*. 2004;35(3):357–366.
- Agulnik SI, Garvey N, Hancock S, et al. Evolution of mouse T-box genes by tandem duplication and cluster dispersion. *Genetics*. 1996;144(1):249–254.
- Naiche LA, Papaioannou VE. Loss of Tbx4 blocks hindlimb development and affects vascularization and fusion of the allantois. *Development*. 2003;130(12):2681–2693.
- Rothman JE, Schekman R. Molecular mechanism of protein folding in the cell. *Cell*. 2011;146(6):851–854.
- van der Brug MP, Blackinton J, Chandran J, et al. RNA binding activity of the recessive parkinsonism protein DJ-1 supports involvement in multiple cellular pathways. *Proc Natl Acad Sci U S A*. 2008;105(29):10244–10249.
- Oda M, Makita M, Iwaya K, et al. High levels of DJ-1 protein in nipple fluid of patients with breast cancer. *Cancer Sci*. 2012;103(6):1172–1176.
- Chen Y, Kang M, Lu W, et al. DJ-1, a novel biomarker and a selected target gene for overcoming chemoresistance in pancreatic cancer. *J Cancer Res Clin Oncol*. 2012;138(9):1463–1474.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress