



## Bimatoprost ophthalmic solution 0.03% lowered intraocular pressure of normal-tension glaucoma with minimal adverse events. [Corrigendum]

Tsumura T, Yoshikawa K, Suzumura H, Kimura T, Sasaki S, Kimura I, Takeda R. *Clinical Ophthalmology*. 2012;6:1547–1552.

On page 1549 the significant difference in MD values was stated as:

$(-5.22 \pm 4.07 \text{ dB and } -2.72 \pm 4.06 \text{ dB, } P = 0.0385)$

It should have been:

$(-5.47 \pm 4.78 \text{ dB and } -5.26 \pm 5.22 \text{ dB, } P = 0.6757)$

Figure 2 legend was missing the words:

Left axis relates to the bars and right axis relates to the line chart.

### Clinical Ophthalmology

#### Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on

Submit your manuscript here: <http://www.dovepress.com/clinical-ophthalmology-journal>

PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.