

Letter to editor

James P Reichmann

Sales and Marketing, American
HomePatient, Brentwood, TN, USA

Dear editor

I would like to congratulate the authors of the article entitled, “Optimal management of nausea and vomiting of pregnancy” for the comprehensive review of treatments for a frequent complication of pregnancy.¹ Although this letter arrives well after publication I think it is important for researchers in the future to have clarification on one seemingly minor but important point. In order to avoid confusion it must be pointed out that in the metoclopramide section the statement “subcutaneous metoclopramide is effective in many women with NVP and HG” is supported by one observational trial on continuous subcutaneous metoclopramide.² This distinction is very important because that study reported patients were receiving the drug for 26.7 days on average at a cost of \$4432 per patient. Unfavorable cost comparisons have been previously published.³ The recommendation that more studied and less expensive alternatives are explored prior to subjecting the patient to this extreme measure seems prudent.⁴ Perhaps the use of the reference to continuous subcutaneous metoclopramide may have been accidental but the suggestion that “these findings are reassuring, especially for women who experience excessive sleepiness...” implies that women on Diclectin® who report being tired are advanced to continuous subcutaneous metoclopramide seems like a waste of health care resources and a little reckless.¹

References

1. Ebrahimi N, Maltepe C, Einarson A. Optimal management of nausea and vomiting of pregnancy. *Int J Womens Health*. 2010;2:241–248.
2. Lombardi DG, Istwan NB, Rhea DJ, O'Brien JM, Barton JR. Measuring outpatient outcomes of emesis and nausea management in pregnant women. *Manag Care*. 2004;13(11): 48–52.
3. Reichmann JP, Kirkbride MK. Nausea and vomiting of pregnancy: cost effective pharmacologic treatments. *Manag Care*. 2008;17(12):41–45.
4. Reichmann JP, Kirkbride MK. Reviewing the evidence for using continuous subcutaneous metoclopramide and ondansetron to treat nausea & vomiting during pregnancy. *Manag Care*. 2012;21(5):44–47

Correspondence: James P Reichmann
1322 Garrick Way
Marietta, GA 30068, USA
Tel +1 615 948 5910
Email jim.reichmann@ahom.com

Authors' response

Neda Ebrahimi^{1,2}
 Caroline Maltepe²
 Adrienne Einarson²

¹Pharmaceutical Sciences, University of Toronto, Toronto, Ontario, Canada; ²Motherisk Program, The Hospital for Sick Children, Toronto, Ontario, Canada

Correspondence: Neda Ebrahimi
 Hospital for Sick Children,
 555 University Avenue, Toronto,
 Ontario M5G 1X8, Canada
 Tel +1 416-813-7654-x4928 mailbox 2
 Email neda.ebrahimi@utoronto.ca

Our statement given with respect to metoclopramide, is meant to be an alternative or add-on therapy for patients who don't respond to Diclectin[®], or suffer from its side effects. In Figure 1 of our article, the algorithm for treatment of nausea and vomiting of pregnancy (NVP) is provided, in which

5–10mg/intramuscular or by mouth, of metoclopramide every 8 hours is suggested. Orally disintegrating tablets, and subcutaneous metoclopramide are both acceptable formulations as well. With respect to the cost/benefit analysis, it was not our intention to encourage the use of the most expensive therapy, but rather the most effective in relieving symptoms and reducing hospitalization and ER visits due to NVP, especially hyperemesis gravidarum (HG). In hue of the costs associated with prolonged hospitalization of HG patients, in some extenuating circumstances, there may even be a favorable cost-benefit ratio. As outlined in your Nausea and Vomiting of Pregnancy Cost Effective Pharmacologic Treatments,¹ we acknowledge that length of therapy may dictate the cost benefit of more expensive drugs, however, the duration of treatment does vary on an individual basis, and hence the cost benefit may still endure.

Reference

1. Reichmann JP, Kirkbride MS. Nausea and vomiting of pregnancy: cost effective pharmacologic treatments. *Manag Care*. 2008;17(12): 41–45.

International Journal of Women's Health

Publish your work in this journal

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. The manuscript management system is completely online and includes

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-womens-health-journal>

Dovepress

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.