

Efficacy of musculoskeletal injections by primary care providers in the office: a retrospective cohort study

Anjali Bhagra¹
Husnain Syed¹
Darcy A Reed¹
Thomas H Poterucha¹
Stephen S Cha²
Tammy J Baumgartner¹
Paul Y Takahashi¹

¹Department of Internal Medicine,

²Department of Health Sciences
Research, Mayo Clinic, Rochester,
MN, USA

Background: Musculoskeletal joint pain of varied etiology can be diagnosed and treated with joint and soft-tissue corticosteroid injections.

Purpose: The purpose of our study was to compare patients' bodily pain and quality of life (QOL), in addition to the procedural benefit and patient satisfaction, before and after musculoskeletal injections in the office setting.

Patients and methods: Patients were eligible for recruitment if they were over age 18 and had an injection for musculoskeletal pain from a primary care provider in an office procedural practice. Included in our analysis were knee joint/bursa, trochanteric bursa, and shoulder joint/bursa injection sites. The variables measured were pain, benefit from the injection, QOL physical and mental components, and patient satisfaction. This was a retrospective cohort study approved by the institutional review board.

Results: Patients' pain was assessed by the patients using a six-point Likert scale (none, very mild, mild, moderate, severe, and very severe). We noted that self-perception of pain decreased from 3.10 (\pm standard deviation at baseline 0.96) before to 2.36 (\pm standard deviation after the injection 1.21) ($P = 0.0001$) after the injection. In terms of the impact on QOL, our patients had a pre-injection physical score of 37.25 ± 8.39 and a mental score at 52.81 ± 8.98 . After the injections, the physical score improved to 42.35 ± 9.07 ($P = 0.0001$) and the mental to 53.54 ± 8.20 ($P = 0.0001$) for the overall group. Ninety-six percent of the patients reported they were satisfied or extremely satisfied in the procedure clinic.

Conclusion: In this study, we found significant pain relief and improved physical QOL in patients undergoing an injection in the knee joint/bursa, shoulder joint/bursa, or trochanteric bursa by primary care providers in the office setting.

Keywords: injections, musculoskeletal, quality of life, joints, efficacy

Introduction

Musculoskeletal injections are an effective treatment for joint and soft-tissue pain, due to a variety of causes. Typically, providers initially address musculoskeletal and, in particular, joint pain with oral systemic anti-inflammatory agents; however, studies have highlighted the effectiveness of local injections as monotherapy in certain settings.¹⁻³ Injection therapy is performed by a variety of providers, including orthopedic surgeons, rheumatologists, physiatrists, podiatrists, and primary care physicians (family medicine, internal medicine). While studies have shed some light on the efficacy of these injections⁴ and have compared variations in utilization patterns of corticosteroids and local anesthetics between different specialty and primary care providers,⁵ it is unclear how these procedures improve patients' quality of life (QOL) when performed by primary care physicians in a purely outpatient setting.

Correspondence: Anjali Bhagra
Mayo Clinic, 200 First Street SW,
Rochester, MN, 55905, USA
Tel +1 507 284 2511
Fax +1 507 284 4959
Email bhagra.anjali@mayo.edu

In addition to improving pain, patients often undergo local injection therapy as a means of improving their QOL. The use of quality as a benchmark of superior medical care is becoming increasingly important. Self-assessment is an often-used means of examining the effects of an intervention on a patient's QOL. One such self-survey is the Short Form-8 (SF-8), which has been previously used in studies of injection therapy.^{6,7} There are a few studies from primary care practices where a patient database was used to track the effectiveness of injection-therapy procedures. However, most databases are of studies with small patient numbers or are from mainly rheumatology practices, some of which examine the utility of ultrasound-guided injections.^{8,9} To our knowledge, this is the first study of its size to examine patients' QOL outcomes, following local injection therapy in a primary care setting, using a simple self-assessment tool such as the SF-8 as part of a provider-maintained patient database.

The primary purpose of this study was to determine if patients experienced an improvement in pain and QOL parameters after joint and soft-tissue injections. The secondary outcome of interest involved determining patient satisfaction with the procedure itself. To answer these questions, we developed and reviewed a registry of patients undergoing injections in an outpatient, primary care, multidisciplinary procedural practice, which involved physicians, registered nurses, and licensed nurse practitioners.

Material and methods

Study design

This was a retrospective cohort study of 1188 adults, over age 18, seen in Mayo Clinic's Primary Care Internal Medicine practice, in Rochester, Minnesota. The study was approved by Mayo Clinic's Institutional Review Board and was conducted according to the principles of the Declaration of Helsinki.¹⁰

Setting

The study was conducted in an outpatient procedural practice within the larger outpatient practice in Rochester. The decision to proceed with the injection and the injection site were determined by the patients' primary care providers. Patients were injected by select primary care providers proficient in musculoskeletal injections who formed the outpatient procedural practice within the larger outpatient primary care practice. Patients gave their clinical consent for injection in accordance with standard clinical practice. The records of the participants were evaluated from July 2010 to June 2011. Data were collected from the same time period.

Participants

Patients were over the age of 18. Patients were eligible for recruitment if they had an injection for musculoskeletal pain in the procedural practice during the period mentioned above. In concordance with Minnesota law, participants were excluded if they did not give authorization for medical record use.

Technique

We included injections around the knee (joint and bursa), shoulder (joint and bursa), and hip (trochanteric bursa) in the analysis. The technique for most of these injections was landmark-guided appropriate for the region. Some providers aspirated joint and bursal fluid prior to injections; however, this information was not collected. All primary care physicians performing the procedures had direct-observation training experience for 6 to 8 weeks before independently performing the procedures. One provider with ultrasound training did selected procedures around the knee and shoulder by utilizing ultrasound guidance.

Variables

There were four primary outcome variables: pain, benefit from the injection, QOL, and patient satisfaction. We measured the mean pain scores before and after injection. In addition, we obtained and compared dichotomized pain data (significant pain – severe or very severe, versus no significant pain – none to moderate) before and after injection.

Patients' QOL was measured using the SF-8, a validated measure of QOL. The SF-8 QOL survey consists of physical and mental QOL measures. Each component is scored from 0 to 100, with higher scores reflecting a better QOL. All of the information was obtained through patient-completed pre-injection surveys in the office, prior to the procedure, and by nursing-staff follow-up telephone calls, 4 weeks post-procedure. Additionally, information regarding patients' satisfaction and perceived benefit from the injection was collected on the follow-up phone calls. Patients' satisfaction was measured using a four-point Likert scale: extremely satisfied, satisfied, somewhat satisfied, or not satisfied. Likewise, patients' benefit from the injection was measured using another four-point Likert scale: no benefit, very little benefit, some benefit, or a lot of benefit. All of the above data, in addition to demographic information, were recorded in a procedural clinic registry database. The primary predictor variables were age and gender, which were obtained from the electronic medical record.

Data sources and measurement

The primary data sources were the electronic medical record and the procedural clinic registry. The effectiveness of the procedures was collected as noted above.

Bias

The primary source of bias was the potential placebo response to the injection. The act of using a needle and having a specialized visit might have predisposed some participants to feel that the pain should improve after the procedure. There was also a potential for bias from the pre-injection survey, which was filled out by patients in the office, and from the post-injection information, which was collected through follow-up phone calls made by nursing staff. This was done because a majority of the patients did not need to present in person for another clinical evaluation.

Study size

All eligible participants were enrolled in the study.

Statistical methods

The demographic aspects of the cohort, using mean \pm standard deviation for continuous variables or frequency (percentage) for categorical variables, were reported. The Student's paired *t*-test was used to compare changes from pre- to posttesting in pain scores, and from SF-8 physical and mental scores. Pearson's Chi-squared test was used to compare patient satisfaction with benefit from the procedure,

following the injection. Any *P*-value of <0.05 was considered statistically significant. All analyses were performed by SAS version 9.3 software (SAS Institute Inc, Cary, NC, USA).

Results

Patients

A total of 1456 records from 1188 participants were accumulated using the REDCAP database in this study from July 2010 to June 2011. All participants had authorization for participation in this research. There were 138 records with no follow-up information; therefore, they were excluded from analysis. Out of the remaining 1318 records, 97 had injections in locations other than a knee (joint and bursa), shoulder (joint and bursa), or hip (trochanteric bursa), and 33 had injections at multiple locations; hence, these records were also excluded (Figure 1).

Out of 1188 records, 839 (71%) were women and 349 (29%) were men. Seventy-seven percent of the patients studied were over 60 years of age, with an average age of 68.98 (20–99). The 968 participants studied had the following individual injections: 598 on the knee, 311 on the shoulder, and 279 on the hip. Forty-five (3.78%) of the injections used ultrasound and were not subcategorized. The average Charlson comorbidity index was 1.44 (0, 14) (Tables 1 and 2).

Outcomes

Overall, our patients had a pre-injection physical score of 37.25 ± 8.39 and a mental score of 52.81 ± 8.98 . After the injections, the physical score improved to 42.35 ± 9.07 ($P = 0.0001$)




Figure 1 Patient demographics (from July 2009 to July 2011).

Table 1 Demographics

Variable	Overall (n = 1188)
Age	68.98 ± 13.41
Charlson comorbidity index	1.44 ± 2.30
Age, n (%)	
20–49	96 (8%)
50–59	181 (15%)
60–69	284 (24%)
70–79	342 (29%)
80+	285 (24%)
Gender, n (%)	
Male	349 (29%)
Female	839 (71%)
Procedure, n (%)	
Knee, n (%)	598 (50%)
Shoulder, n (%)	311 (26%)
Hip, n (%)	279 (23%)

Abbreviation: n, number.

and the mental to 53.54 ± 8.20 ($P = 0.0001$) for the overall group. Self-perception of pain also decreased from $3.10 (\pm 0.96)$ to $2.36 (\pm 1.21)$ ($P = 0.0001$). Thus, the group, as a whole, felt improvement after the procedure. By gender, men had a significant reduction in body pain scores and improvement in the physical score, but no change in the mental score. Women had improvement in the physical and mental scores, as well as reductions in body pain scores (all P values were <0.001). Subjects over age 60 had improvement in physical and mental scores and reduction in overall body pain. Those under 60 had improvements in physical scores, but not mental scores (Table 3).

There were improved physical scores and reduced body pain across all three injected sites. Physical scores improved

significantly for all three injection sites (knee, shoulder, and hip). Mental QOL improved with knee injections in females and those over 60 years of age; however, no change was noted for shoulder and hip injections.

Patient satisfaction with the injection was generally good overall, with 1142 patients (96%) reporting they were satisfied or extremely satisfied with the procedure clinic and 905 patients (76%) reporting some or a lot of benefit from the injection. There were no differences noted across age and gender. The perceived benefit of injections differed by joint site: the perceived benefit from knee injection was at 80% – higher than that from the shoulder injection, at 73%, and from the hip, at 70% ($P = 0.014$ and $P = 0.0008$, respectively). The variables of gender, age, and injection site all showed the same trend (Table 4).

Discussion

In our study, we found a significant improvement in bodily pain scores 4 weeks after injections at the knee, shoulder, and the hip overall, and in all groups. The administration of a local steroid injection therapy, regardless of location or patient age, led to significant reduction in pain levels and improvement of physical QOL measures. These results support the notion that primary care physicians, well trained and experienced in the administration of local injection therapy, can reduce patients' joint-pain levels and improve their physical QOL. Probably the most common uses for intra-articular corticosteroid injections are in the knees and shoulders of patients with degenerative joint diseases, including osteoarthritis. The benefit of local corticosteroid

Table 2 Disease severity

Variable	Overall (n = 1188)	Knee (n = 598)	Shoulder (n = 311)	Hip (n = 279)	P value
Charlson index	1.4 ± 2.30	1 ± 2.11	1.6 ± 2.52	1.6 ± 2.42	0.1
MI, n (%)	57 (5%)	23 (4%)	24 (8%)	10 (4%)	0.019
CHF, n (%)	81 (7%)	43 (7%)	22 (7%)	16 (6%)	0.71
PVD, n (%)	84 (7%)	38 (6%)	25 (8%)	21 (8%)	0.61
CVD, n (%)	118 (10%)	57 (10%)	30 (10%)	31 (11%)	0.75
Dementia, n (%)	3 (0%)	2 (0%)	1 (0%)	0 (0%)	0.63
C_pulm, n (%)	200 (17%)	91 (15%)	52 (17%)	57 (20%)	0.16
Ulcer, n (%)	58 (5%)	26 (4%)	17 (5%)	15 (5%)	0.69
M_liver, n (%)	29 (2%)	14 (2%)	10 (3%)	5 (2%)	0.52
Diabetes, n (%)	163 (14%)	78 (13%)	46 (15%)	39 (14%)	0.76
C_diabetes, n (%)	58 (5%)	23 (4%)	20 (6%)	15 (5%)	0.21
Hemiplegia, n (%)	7 (1%)	3 (1%)	1 (0%)	3 (1%)	0.45
Renal, n (%)	86 (7%)	42 (7%)	25 (8%)	19 (7%)	0.81
S_liver, n (%)	2 (0%)	1 (0%)	1 (0%)	0 (0%)	0.64
Metastatic, n (%)	30 (3%)	10 (2%)	10 (3%)	10 (4%)	0.16
AIDS, n (%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
Rheum, n (%)	62 (5%)	27 (5%)	18 (6%)	17 (6%)	0.54
Cancer, n (%)	186 (16%)	90	50	46	0.84

Note: If a person has all variables, the Charlson index can be 33.

Abbreviations: MI, myocardial infarction; CHF, congestive heart failure; CVD, cerebrovascular disease; PVD, peripheral vascular disease; DEMEN, dementia; CPULM, chronic pulmonary disease; M_liver, mild liver disease; C_diabetes, diabetes with end organ disease; Renal, moderate or severe liver disease; S_liver, moderate or severe liver disease; Metastatic, metastatic solid tumor; Rheum, connective tissue or rheumatologic disease.

Table 3 Pre- and posttesting by injection location

Group	Mean ± SD		P-value
	Pre	Post	
Physical score (higher better)			
Overall (n = 1188)	37.25 ± 8.39	42.35 ± 9.07	0.0001
Knee (n = 598)	36.51 ± 7.98	42.06 ± 8.94	0.0001
Shoulder (n = 311)	39.56 ± 8.67	43.30 ± 8.99	0.0001
Hip (n = 279)	36.28 ± 8.48	41.93 ± 9.40	0.0001
Mental score (higher better)			
Overall (n = 1188)	52.81 ± 8.98	53.54 ± 8.20	0.0001
Knee (n = 598)	52.88 ± 9.14	53.84 ± 7.97	0.0001
Shoulder (n = 311)	53.37 ± 8.32	53.64 ± 8.10	0.3609
Hip (n = 279)	52.05 ± 9.31	52.79 ± 8.76	0.0471
Body pain score (0–5/none–severe)			
Overall (n = 1188)	3.10 ± 0.96	2.36 ± 1.21	0.0001
Knee (n = 598)	3.05 ± 0.98	2.27 ± 1.18	0.0001
Shoulder (n = 311)	3.01 ± 1.00	2.33 ± 1.24	0.0001
Hip (n = 279)	3.29 ± 0.88	2.58 ± 1.21	0.0001

Abbreviation: SD, standard deviation.

injections into the knee has been well studied and reported in patients with osteoarthritis.^{11,12} This study reinforces the importance of these injections in the primary care office as a viable mechanism to reduce pain in all types of patients. Our study additionally highlights the positive impact of

pain reduction on the overall QOL of people undergoing injection therapy.

In our evaluations, we found a difference in mental QOL when using a subgroup analysis. Men demonstrated a greater improvement in physical QOL measures, compared to mental QOL. Gender-related differences in response to injections have been reported in previous studies.¹³ The above finding in our study may reflect a lack of pain impact on men's mental QOL, or it may reflect a greater pain impact on females' mood. In the older adults, one likewise saw a greater impact of injections on mental QOL scores, compared to younger adults.

Across different injections sites, we found universal improvement in pain scores and physical QOL. It is further interesting to note that the results across the different injection sites were similar across different age groups and genders. Certain studies have shown that more than 20% of injections are, in fact, not intra-articular.^{14,15} Thus, one might expect some differences between different injection types, depending upon the ease of intra-articular or intrabursal injection. This might be explained by some reports suggesting that a precise intra-articular injection of a steroid may not be necessary for clinical benefit.^{15,16} By some reports, up

Table 4 Patient satisfaction ratings

Variable	Overall (n = 1188) (%)	Knee (n = 598) (%)	Shoulder (n = 311) (%)	Hip (n = 279) (%)	P-value
Did you follow aftercare instructions: limitation of physical activity, after the musculoskeletal injection?					0.59
Yes	964 (81)	484 (81)	254 (82)	226 (81)	
No	67 (6)	29 (5)	22 (7)	16 (6)	
Partly	154 (13)	82 (14)	35 (11)	37 (13)	
How much benefit did you derive from the joint/bursa injection?					0.022
No benefit	120 (10)	50 (8)	35 (11)	35 (13)	
Very little benefit	163 (14)	67 (11)	48 (15)	48 (17)	
Some benefit	317 (27)	166 (28)	88 (28)	63 (23)	
A lot of benefit	588 (49)	315 (53)	140 (45)	133 (48)	
If a second injection was performed, how much benefit did you derive from that joint/bursa injection?					0.28
No benefit	22 (8)	12 (7)	4 (8)	6 (13)	
Very little benefit	34 (13)	18 (11)	6 (12)	10 (21)	
Some benefit	77 (29)	53 (32)	16 (31)	8 (17)	
A lot of benefit	132 (50)	83 (50)	26 (50)	23 (49)	
How would you rate your satisfaction in the procedure clinic?					0.47
Extremely satisfied	849 (71)	434 (73)	212 (68)	203 (73)	
Satisfied	293 (25)	144 (24)	81 (26)	68 (24)	
Somewhat satisfied	32 (3)	15 (3)	12 (4)	5 (2)	
Not satisfied	14 (1)	5 (1)	6 (2)	3 (1)	
Variables – combined ratings					
How much benefit did you derive from the joint/bursa injection?					0.002
No or very little benefit	283 (24)	117 (20)	83 (27)	83 (30)	
Some or a lot of benefit	905 (76)	481 (80)	228 (73)	196 (70)	
How would you rate your satisfaction in the procedure clinic?					0.12
Extremely satisfied or satisfied	1142 (96)	578 (97)	293 (94)	271 (97)	
Somewhat or not satisfied	46 (4)	20 (3)	18 (6)	8 (3)	

to three-fourths of patients with osteoarthritis of the knee, receiving a local injection therapy, experience a significant positive response.¹⁷ We found similar clinical results, with over 76% overall noting an improvement in pain. Another study reported a wide variation of success – from conservative management that included physical therapy, injections, and systemic pharmacotherapy – of 33% to 92%, in patients with shoulder pain secondary to rotator cuff tear.¹⁸ Clinical and patient predictors identifying those most likely to benefit are scarce,¹⁹ forming an area of much-needed research and insight.

The majority of patients, including those that reported no clinical benefit, reported being significantly satisfied by their experience with the administration of injection therapy. Thus, 96% were satisfied or extremely satisfied, despite only a 76% self-perceived benefit. This may reflect the patients' satisfaction with the attempt to improve the pain, even though the pain was not resolved. Many patients may perceive the benefits, which include improved pain levels, mobility, ability to participate in physical therapy, and QOL measures. However, a limiting factor of injection therapy is its short-term effectiveness.^{1,20} A Cochrane review reported corticosteroid-containing joint injections to be more effective than a placebo or hyaluronic acid, with respect to pain relief, for up to 3 weeks.¹ However, some reports suggest that hyaluronic acid may be better for longer-term pain relief beyond 4 weeks.²¹ The longer-term benefits of corticosteroids are questionable, and differences in outcomes between injection therapy and conservative management may not be significant.²²

This study had both strengths and weaknesses. First, it relied on a self-report option as opposed to anthropomorphic measurements, which may have led to recall bias or other biases. Additionally, there may be a placebo effect from the injection. In a systematic review of the Cochrane database, it was noted that for rotator cuff disease, subacromial steroid injection had a small benefit over placebo in some trials.²³ The QOL questionnaire and self-reported pain are standard measures, which should be reproducible in other offices and clinical settings. The providers who performed these procedures are primary care providers who devote a part of their clinical practice to procedures. Therefore, results should be generalizable to those family medicine physicians and internists who routinely perform procedures in the office; however, the results of physicians performing the procedures less frequently may differ. The population of Olmsted County, and of southeastern Minnesota in general, is largely Northern European, so the results may differ in other populations.

Conclusion

In this study of 1188 patients, we found pain relief and improved physical QOL in patients undergoing a joint injection in the knee, shoulder, or hip. This improvement did not change by age, gender, or injection site. Mental QOL improved overall, but did so specifically in women and those over 60 years of age. Knee injections similarly improved mental QOL. Localized corticosteroid injections can be successfully performed in an outpatient clinic by primary care physicians, and they can improve QOL.

Acknowledgments/disclosure

We would like to acknowledge the Primary Care Internal Medicine procedure nurses and support staff that have helped with surveys and follow-up phone calls: Christine Bray, Carol Fealey, Kristine McCoy, Jacqueline McGee, Jean Mensink, Lori Momany, and Julie Rindels. The authors report no conflicts of interest in this work.

References

1. Bellamy N, Campbell J, Robinson V, Gee T, Bourne R, Wells G. Intraarticular corticosteroid for treatment of osteoarthritis of the knee. *Cochrane Database Syst Rev*. 2006;2:CD005328.
2. Friedman DM, Moore ME. The efficacy of intraarticular steroids in osteoarthritis: a double-blind study. *J Rheumatol*. 1980;7(6):850–856.
3. Ravaud P, Moulinier L, Giraudeau B, et al. Effects of joint lavage and steroid injection in patients with osteoarthritis of the knee: results of a multicenter, randomized, controlled trial. *Arthritis Rheum*. 1999;42(3):475–482.
4. Skedros JG, Hunt KJ, Pitts TC. Variations in corticosteroid/anesthetic injections for painful shoulder conditions: comparisons among orthopaedic surgeons, rheumatologists, and physical medicine and primary-care physicians. *BMC Musculoskelet Disord*. 2007;8:63.
5. Burbank KM, Stevenson JH, Czarnecki GR, Dorfman J. Chronic shoulder pain: part II. Treatment. *Am Fam Physician*. 2008;77(4):493–497.
6. Ware J, Kosinski M, Dewey J, Gandek B. *How to Score and Interpret Single-item Health Status Measures: A Manual for Users of the SF-8 Health Survey*. 2nd Ed. Boston, MA: QualityMetric; 2001.
7. Quality Metric. *The SF-8™ Health Survey*. 2002 [webpage on the Internet]. Medical Outcomes Trust. Available from: <http://www.sf-36.org/tools/sf8.shtml>. Accessed October 27, 2008.
8. Smith MD. Can a rheumatologist be a procedural specialist? The experience of an Australian rheumatologist. *Int J Rheumatic Dis*. 2008;11(2):118–120.
9. El Miedany Y. Musculoskeletal US: examining the joints. *Br J Nurs*. 2012;21(6):340–344.
10. Goodyear MD, Krleza-Jeric K, Lemmens T. The Declaration of Helsinki. *BMJ*. 2007;335(7621):624–625.
11. Arroll B, Goodyear-Smith F. Corticosteroid injections for osteoarthritis of the knee: meta-analysis. *BMJ*. 2004;328(7444):869.
12. Godwin M, Dawes M. Intra-articular steroid injections for painful knees. Systematic review with meta-analysis. *Can Fam Physician*. 2004;50:241–248.
13. Leopold SS, Redd BB, Warne WJ, Wehrle PA, Pettis PD, Shott S. Corticosteroid compared with hyaluronic acid injections for the treatment of osteoarthritis of the knee. A prospective, randomized trial. *J Bone Joint Surg Am*. 2003;85-A(7):1197–1203.

14. Bliddal H. Placement of intra-articular injections verified by mini arthrography. *Ann Rheum Dis.* 1999;58(10):641–643.
15. Jones A, Regan M, Ledingham J, Patrick M, Manhire A, Doherty M. Importance of placement of intra-articular steroid injections. *BMJ.* 1993;307(6915):1329–1330.
16. Sambrook PN, Champion GD, Browne CD, et al. Corticosteroid injection for osteoarthritis of the knee: peripatellar compared to intra-articular route. *Clin Exp Rheumatol.* 1989;7(6):609–613.
17. Gaffney K, Ledingham J, Perry JD. Intra-articular triamcinolone hexacetonide in knee osteoarthritis: factors influencing the clinical response. *Ann Rheum Dis.* 1995;54(5):379–381.
18. Longo UG, Franceschi F, Berton A, Maffulli N, Droena V. Conservative treatment and rotator cuff tear progression. *Med Sport Sci.* 2012;57: 90–99.
19. Jones A, Doherty M. Intra-articular corticosteroids are effective in osteoarthritis but there are no clinical predictors of response. *Ann Rheum Dis.* 1996;55(11):829–832.
20. Hepper CT, Halvorson JJ, Duncan ST, Gregory AJ, Dunn WR, Spindler KP. The efficacy and duration of intra-articular corticosteroid injection for knee osteoarthritis: a systematic review of level I studies. *J Am Acad Orthop Surg.* 2009;17(10):638–646.
21. Bannuru RR, Natov NS, Obadan IE, Price LL, Schmid CH, McAlindon TE. Therapeutic trajectory of hyaluronic acid versus corticosteroids in the treatment of knee osteoarthritis: a systematic review and meta-analysis. *Arthritis Rheum.* 2009;61(12):1704–1711.
22. Brinks A, van Rijn RM, Willemsen SP, et al. Corticosteroid injections for greater trochanteric pain syndrome: a randomized controlled trial in primary care. *Ann Fam Med.* 2011;9(3):226–234.
23. Buchbinder R, Green S, Youd JM. Corticosteroid injections for shoulder pain. *Cochrane Database Syst Rev.* 2003;1:CD004016.

International Journal of General Medicine

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-general-medicine-journal>

Dovepress

A key focus is the elucidation of disease processes and management protocols resulting in improved outcomes for the patient. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.