Asymptomatic bacteriuria among elderly and middle-aged rural community-dwellers in South-Western Nigeria

OA Olowe¹
OB Makanjuola¹
KO Olabiyi¹
PO Akinwusi²
CO Alebiosu²
MA Isawumi³
MB Hassan³
EO Asekun-Olarinmoye⁴
WO Adebimpe⁴
TA Adewole⁵

¹Department of Medical Microbiology and Parasitology, ²Department of Medicine, ³Department of Ophthalmology, ⁴Department of Community Medicine, ⁵Department of Biochemistry, College of Health Sciences, Osun State University, Osogbo, Osun State, Nigeria

Abstract: Asymptomatic bacteriuria in elderly individuals has been well described in institutionalized settings, but to a lesser extent in the community. The purpose of this study was to determine the pathogens responsible for asymptomatic bacteriuria in elderly and middle-aged individuals in Alajue-Ede, South-Western Nigeria, and to identify any associated factors. Mid-stream urine samples were collected from apparently healthy elderly and middle-aged volunteers who were participating in community health screening. Samples were processed and bacterial isolates were identified following standard procedures. In total, 128 volunteers (48 men, 76 women) participated in the study. Twenty-eight (22.6%) urinary pathogens were isolated, comprising *Klebsiella* species in five (17.9%), *Pseudomonas aeruginosa* in one (3.6%), *Escherichia coli* in 19 (67.9%), and *Proteus* species in three (10.7%) cases. Women were identified as being at higher risk of asymptomatic bacteriuria, and the prevalence also increased with increasing age in men. The elderly in this community have a high prevalence of asymptomatic bacteriuria, and screening for comorbid medical conditions may be of benefit.

Keywords: asymptomatic bacteriuria, urinary pathogens, elderly, urinary tract infection

Introduction

Urinary tract infections (UTIs) are common and require treatment with antibiotics.¹ Asymptomatic bacteriuria is defined as significant bacteriuria without symptoms of UTI. Significant bacteriuria has been defined as the presence of 1×10^5 or more bacteria per milliliter of voided mid-stream urine.²

Asymptomatic bacteriuria is common, with varying prevalence according to age, gender, sexual activity, and the presence of genitourinary abnormalities. Several categories of people are reported to be at higher risk, including the elderly, pregnant women, and those with diabetes, particularly women.³ Among the elderly, those in institutionalized settings such as long-term care facilities and hospitals have a higher prevalence of asymptomatic bacteriuria than those living in the community.^{4,5} Therefore, interest in asymptomatic bacteriuria has shifted from the general population to these groups of individuals.⁶ The prevalence varies from 2% to 10% in pregnancy and from 15% to 50% in the elderly in long-term care facilities.^{5,7}

There has been controversy and conflicting reports as to the significance and benefits of treating individuals with asymptomatic bacteriuria.^{4,8} Significant asymptomatic bacteriuria has been described as an important cause of pyelonephritis and Gram-negative septicemia among predisposed individuals, such as diabetics.⁹ Asymptomatic bacteriuria is also believed to be harmful in pregnant women, men or women undergoing invasive genitourinary procedures, and renal transplant recipients. In these groups,

Correspondence: OB Makanjuola
Department of Medical Microbiology
and Parasitology, College of Health
Sciences, Osun State University,
PMB 4494, Osogbo, Osun State, Nigeria
Email funmifemi@yahoo.com

adverse outcomes can be prevented using antimicrobial drug therapy and studies have shown that these groups are also at increased risk of symptomatic UTI.⁶ However, there are doubts as to the benefit of treating asymptomatic bacteriuria in elderly individuals, even in institutionalized settings, despite the risk of symptomatic UTI.⁶

This study was undertaken to determine the prevalence of urinary tract pathogens in older people without symptoms in a community setting and to identify factors associated with asymptomatic bacteriuria in these people.

Materials and methods

This cross-sectional study in the village of Alajue-Ede, South-Western Nigeria, was part of a community health survey of antibiotic use and resistance on April 2011 during World Health Day. Participants in this study were community-dwellers who were at least 40 years of age. Those aged 40–69 years were regarded as middle-aged and those aged 70 years and older were regarded as elderly. The inclusion criteria was apparent good health and ability to give informed consent. Individuals who had been hospitalized during the previous month were excluded. The majority of participants were of low socioeconomic status. Most were peasant farmers and fishermen, while a small percentage were community traders. A free health care system funded by government helps to support community health in this village. This study was approved by the ethics committee of the College of Health Sciences, Osun State University, Osogbo, Nigeria. Informed consent was obtained from all participants after an adequate explanation of the study.

Procedure

A semistructured questionnaire was given to all volunteers after obtaining informed consent. Thereafter, mid-stream urine specimens were collected and sent to the microbiology laboratory for microscopy and culture. For isolation of organisms, each urine sample was mixed thoroughly and a standard urine sample was inoculated onto blood agar and MacConkey agar medium. All inoculated plates were incubated aerobically at 37°C for 24 hours. Bacteriuria $\geq 1 \times 10^5$ colony-forming units/mL of urine was regarded as significant bacteriuria. Standard microbiological methods of isolation and identification as outlined by Mackie and McCartney were carefully followed in processing the specimens.¹⁰

Statistical analysis

Patient demographic and microbiological data were entered using the standard format. All data were analyzed

using Statistical Package for the Social Sciences version 16 software (SPSS Inc, Chicago, IL, USA).

Results

Twenty-eight (22.6%) of 124 samples showed significant bacterial growth, comprising seven from 48 women and 21 from 76 men. The most frequent isolate was *Escherichia coli* (n = 19, 67.9%), followed by *Klebsiella* species (n = 5, 17.9%), *Proteus* species (n = 3, 10.7%), and *Pseudomonas aeruginosa* (one, 3.6%, Table 1). Table 2 shows the age distribution and prevalence of significant bacteriuria according to gender. The age range was 40 to more than 70 years, and the male to female ratio was 2:3. There was a higher prevalence of asymptomatic bacteriuria in elderly subjects than in their younger counterparts.

Discussion

To the authors' knowledge, this study is one of the few addressing the issue of significant bacteriuria in older individuals and the elderly who are not in institutional care. We found a prevalence of 22.6%, which is much higher than that reported by others for this group of individuals, ie, community-dwellers without overt clinical pathology and not in institutional care.^{1,7,11} However, our finding is quite comparable with what has been described in nursing homes and other long-term care facilities, where a high prevalence of asymptomatic bacteriuria has been attributed to impaired genitourinary status, instrumentation, and increased use of antibiotics.^{7,12} The prevalence of asymptomatic bacteriuria in our community-based study is also similar to findings for diabetics, as described by Alebiosu et al.¹³ However, ours was a group of apparently healthy individuals who were not taking antibiotics and had no overt evidence of UTI or pathology. This high prevalence could be a result of as yet undetected comorbid medical conditions in these individuals.

Table 1 Prevalence of bacteriuria and the range of uropathogens recovered from urine of asymptomatic volunteers

Characteristics	Male volunteers (n = 48)	Female volunteers (n = 76)	Total (n = 124)
Bacteriuria status			
Significant	7 (14.6%)	21 (27.6%)	28 (22.6%)
Non significant	10 (20.8%)	11 (14.5%)	21 (16.9%)
No growth	31 (64.6%)	44 (57.9%)	75 (60.5%)
Bacterial organisms			
<i>Escherichia coli</i>	5 (71.4%)	14 (66.7%)	19 (67.9%)
<i>Klebsiella</i> species	1 (14.3%)	4 (19.0%)	5 (17.9%)
<i>Proteus</i> species	1 (14.3%)	2 (9.5%)	3 (10.7%)
<i>Pseudomonas aeruginosa</i>	0 (0.0%)	1 (4.8%)	1 (3.6%)

Table 2 Age distribution and prevalence of significant bacteriuria in male and female volunteers

Age (years)	Sex	No of subjects	Significant bacteriuria no (%)	No (%) of isolates			
				<i>E. coli</i>	<i>Klebs spp</i>	<i>Proteus spp</i>	<i>P. aeruginosa</i>
40–49	M	8	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
	F	22	5 (22.7)	2 (40.0)	1 (20.0)	2 (40.0)	0 (0.0)
50–59	M	14	1 (7.1)	1 (100)	0 (0.0)	0 (0.0)	0 (0.0)
	F	21	7 (33.3)	6 (85.7)	1 (14.3)	0 (0.0)	0 (0.0)
60–69	M	15	3 (20)	2 (66.7)	1 (33.3)	0 (0.0)	0 (0.0)
	F	17	4 (23.5)	4 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
≥70	M	11	3 (27.3)	2 (66.7)	0 (0.0)	1 (33.3)	0 (0.0)
	F	16	5 (31.3)	2 (40.0)	2 (40.0)	0 (0.0)	1 (20.0)

It is important to screen such individuals for diabetes which is often found in association with asymptomatic bacteriuria.³ Our finding of an almost 23% prevalence of asymptomatic bacteriuria in older community-dwellers underscores the need for regular health screening, especially in high-risk groups.

E. coli was found to be the leading urinary pathogen in this study, and was documented in 66.7% and 71.4% of women and men, respectively. Becker et al, Drucker et al, and Olusanya and Olutiola have reported that the most common urinary pathogens causing asymptomatic bacteriuria are enteric organisms, ie, *E. coli* and other coliforms.^{14–16} In contrast with this observation, Dada-Adegbola and Muili reported that in their study *Klebsiella* was the leading urinary pathogen.¹⁷ Organisms such as *Pseudomonas* and *Proteus* were quite rare in this study, unlike in institutional settings where these pathogens are more common, likely because they are not usually acquired in the community.¹⁸

The results of our study also indicate that the prevalence of asymptomatic bacteriuria is higher in women than in men. This is in agreement with the findings of several earlier studies.^{7,11} This is related to the short course of the female urethra and its proximity to the anorectal region, making it easier for enteric organisms to colonize, ascend, and subsequently attach to the uroepithelium. The prevalence of asymptomatic bacteriuria also appears to increase with increasing age in men, as reflected in a prevalence of 7% in 50-year-olds, which increased to 20% among those in their 60's then to 31% in the elderly. It is uncertain why women did not show the same prevalence pattern, despite several reports that prevalence of asymptomatic bacteriuria increases with age, even among women.^{7,18} The picture in men is likely to be attributable to the prostatic enlargement that occurs with increasing age and causes urinary retention.^{5,7}

We also noted that the prevalence of asymptomatic bacteriuria in the middle-aged group was 20.6% whereas it

was 29.6% in the elderly group. Further investigation did not reveal any statistically significant difference in prevalence between these two age groups despite an initial indication of a significant difference. This might be because of the small size of our study, and further studies are needed to determine whether there is any difference in the prevalence of asymptomatic bacteriuria between middle-aged and elderly adults.

In conclusion, although there is controversy regarding whether to treat asymptomatic bacteriuria in an otherwise healthy adult, we recommend that further investigations be carried out in individuals living in the community setting to determine the cause of this high prevalence of asymptomatic bacteriuria. Underlying condition(s) could require treatment, which might prevent unnecessary morbidity.

Disclosure

The authors report no conflicts of interest in this work.

References

- Colgan R, Nicolle LE, McGlone A, Hooton TM. Asymptomatic bacteriuria in adults. *Am Fam Physician*. 2006;74:985–990.
- Rubin RH, Shapiro ED, Andriole VT, Davis RJ, Stamm WE. Evaluation of new anti-infective drugs for the treatment of urinary tract infection. *Clin Infect Dis*. 1992;15:216–227.
- Renko M, Tapanainen P, Tossavainen P, Pokka T, Uhari M. Meta-analysis of the significance of asymptomatic bacteriuria in diabetes. *Diabetes Care*. 2011;34:230–235.
- Nicolle LE, Bradley S, Colgan R, Rice KC, Schaeffer A, Hooton MT. Infectious Diseases Society of America guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis*. 2005;40:643–654.
- Nelius T, Filleur S, Nelson JS. *Asymptomatic Bacteriuria: Significance for Different Patient Population, Urinary Tract Infections*. Manhattan, NY: Intech; 2011. Available from: <http://www.intechopen.com/books/urinary-tract-infections/asymptomatic-bacteriuria-significance-for-different-patient-population>. Accessed May 16, 2013.
- Nicolle LE. Asymptomatic bacteriuria – important or not? *N Engl J Med*. 2000;343:1037–1039.
- Rodhe N, Molstad S, Englund L, Svardsudd K. Asymptomatic bacteriuria in a population of elderly residents living in a community setting: prevalence, characteristics and associated factors. *Fam Pract*. 2006;23:303–307.

8. Lee S, Kim K, Park Y, Kim Y. Association between asymptomatic urinary tract infection and postoperative spine infection in elderly women: a retrospective analysis study. *J Korean Neurosurg Soc.* 2010;47:265–270.
9. Kayima JK, Otieno LS, Twahir A, Njenga E. Asymptomatic bacteriuria among diabetics attending Kenyatta National Hospital. *East Afr Med J.* 1996;73:524–526.
10. Mackie, McCartney. *Practical Medical Microbiology*, 14th ed. London, UK: Churchill Livingstone; 1996.
11. Varli M, Guruzb H, Arasa S, Yalcina A, Atlia T, Turgaya M. Asymptomatic bacteriuria among the elderly living in the community: prevalence, risk factors and characteristics. *Eur Geriatr Med.* 2012;3:87–91.
12. Boscia JA, Kobasa WD, Knight RA, Abrutyn E, Levison ME, Kaye D. Epidemiology of bacteriuria in an elderly ambulatory population. *Am J Med.* 1986;80:208–214.
13. Alebiosu CO, Osinupebi OA, Olajubu FA. Significant asymptomatic bacteriuria among Nigerian type 2 diabetics. *J Natl Med Assoc.* 2003;95:344–349.
14. Watts ER, Hancock V, Ong CY, Vejborg RM, Mabbett AN, Totsika M, et al. *Escherichia coli* isolates causing asymptomatic bacteriuria in catheterized and noncatheterized individuals possess similar virulence properties. *J Clin Microbiol.* 2010;48(7):2449–2458.
15. Roos V, Ulett GC, Schembri MA, Klemm P. The asymptomatic bacteriuria *Escherichia coli* strain 83972 outcomes Uropathogenic *E. coli* strains in human urine. *Infect Immun.* January 2006;74(1):615–624.
16. Olusanya O, Olutiola PO. Studies on bacteriuria in patients and students in Ile-Ife, Nigeria. *West Afr J Med.* 1984;303:177–182.
17. Dada-Adegbola HO, Muili KA. Antibiotics susceptibility pattern of urinary tract pathogens in Ibadan, Nigeria. *Afr J Med Med Sci.* 2010;39:173–179.
18. Ariathianto Y. Asymptomatic bacteriuria – prevalence in the elderly population. *Aust Fam Physician.* 2011;40:805–809.

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

Submit your manuscript here: <http://www.dovepress.com/infection-and-drug-resistance-journal>

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress