Toxic effects of the interaction of titanium dioxide nanoparticles with chemicals or physical factors

Kui Liu
Xialu Lin
Jinshun Zhao

Public Health Department of
Medical School, Zhejiang Provincial
Key Laboratory of Pathological and
Physiological Technology, Ningbo
University, Ningbo, Zhejiang Province,
People's Republic of China

Abstract: Due to their chemical stability and nonallergic, nonirritant, and ultraviolet protective properties, titanium dioxide (TiO_2) nanoparticles (NPs) have been widely used in industries such as electronics, optics, and material sciences, as well as architecture, medicine, and pharmacology. However, increasing concerns have been raised in regards to its ecotoxicity and toxicity on the aquatic environment as well as to humans. Although insights have been gained into the effects of TiO_2 NPs on susceptible biological systems, there is still much ground to be covered, particularly in respect of our knowledge of the effects of the interaction of TiO_2 NPs with other chemicals or physical factors. Studies suggest that interactions of TiO_2 NPs with other chemicals or physical factors may result in an increase in toxicity or adverse effects. This review highlights recent progress in the study of the interactive effects of TiO_2 NPs with other chemicals or physical factors.

Keywords: titanium dioxide, TiO_2 , nanoparticles, interaction, chemicals, physical factors

Introduction

Nanoparticles (NPs) are raw materials used in nanotechnology, with a size range of 1–100 nm in no less than one of their three dimensions.^{1–3} Titanium dioxide (TiO_2) NPs consists of three polymorphs, including anatase, rutile, and brookite.⁴ TiO_2 NPs have been widely used in many products, such as toothpastes, sunscreens, cosmetics, food products, pharmaceuticals, and nanomedical reagents.⁵ TiO_2 particles have been considered as nontoxic mineral particles and traditionally used in the fields of cosmetics, food, and drugs. They were even used as “dust negative control” in many in vitro and in vivo toxicological investigations for many years.^{3,6} However, research evidence suggests that TiO_2 NPs may possess higher toxicity potential than their bulk materials.^{5,7,8} Zhao et al⁸ found that TiO_2 NPs caused higher cytotoxicity than fine particles in cell culture. Due to their very small size, NPs can penetrate basic biological structures, which may, in turn, disrupt their normal function.^{3,9} Recent research evidence shows that TiO_2 NPs may induce cellular toxicity effects in cardiac tissue.¹⁰ The toxicity effects of TiO_2 particles were also observed in cells of the circulatory system. Li et al⁶ found that the erythrocytes treated with TiO_2 NPs underwent abnormal sedimentation, hemagglutination, and hemolysis, which were totally different from those treated with TiO_2 fine particles. Lung tumors were also found in rats after lifetime exposure to high concentrations of TiO_2 particles.¹¹ Moreover, a recent study showed oxidative stress in mice brain as well as overproliferation of all glial cells.¹² These occurred in mice that were exposed to 2.5 mg/kg, 5 mg/kg, and 10 mg/kg body weight TiO_2 NPs through nasal administration for 90 days. The toxicokinetics (Figure 1) and toxic effects of

Correspondence: Jinshun Zhao
Public Health Department of Medical
School, Zhejiang Provincial Key
Laboratory of Pathological and
Physiological Technology, Ningbo
University, Ningbo 315211, Zhejiang
Province, People's Republic of China
Email zhaojinshun@nbu.edu.cn

TiO₂ NPs^{13–15} alone have been well documented in many in vivo and in vitro studies, but reviews of the effects (or toxicities) of the interaction of TiO₂ NPs with other chemicals or physical factors are currently unavailable. TiO₂ NPs may coexist with other chemicals or physical factors in the surrounding environment and occupational settings. In the field of nanomedicine, TiO₂ NPs are being used as drug carriers.³ Therefore, evaluating the interactive effects of TiO₂ NPs with other chemicals or physical factors is vital for the safe application of TiO₂ NPs. This review will mainly focus on the current knowledge concerning the effects of the interaction of TiO₂ NPs with other chemicals or physical factors and will identify areas where further improvement is needed.

Effects of TiO₂ NP interaction with metals and their compounds

With growing applications, TiO₂ NPs are rapidly entering the aquatic environment,¹⁶ and thus the aquatic environment is expected ultimately to be a sink for the sedimentation of these NPs. Consequently, these NPs will inevitably mix and

interact with other aquatic pollutants, including metals and their compounds.¹⁷ In addition, NPs have been found to be capable of absorbing and separating metals from aqueous or organic solutions.^{18,19}

TiO₂ NPs have been tested as a sorbent in the solid-phase extraction for preconcentrated lead (Pb) in river water and seawater.^{20–22} Zhang et al²⁰ investigated the potential acute toxicity of the interaction between TiO₂ NPs (50 nm and 120 nm) and lead acetate (PbAC) in mice. Suspensions of TiO₂ NPs (5 g/kg body weight) alone, PbAC (500 mg/kg) alone, and TiO₂ NPs (5 g/kg) plus PbAC (500 mg/kg) were administered to mice via oral gavage, respectively. No synergistic acute toxicity in mice was found after treatment with the combination of TiO₂ NPs and PbAC. However, Du et al²³ found that, compared with the control (1% dimethyl sulfoxide), mixtures of TiO₂ NPs of different doses (21 nm, 80% anatase, 20% rutile) plus PbAC (1 µg/mL) induced a significant increase in reactive oxygen species (ROS) generation (at 0.001 µg/mL, 0.01 µg/mL, 0.1 µg/mL, 1 µg/mL, and 10 µg/mL of TiO₂), intracellular superoxide dismutase (SOD) activity (at 0.1 µg/mL and

Figure 1 Toxicokinetics and accumulation sites of titanium dioxide nanoparticles.

Note: Reprinted from Shi et al,¹³ Copyright 2013, with permission from BioMed Central Publishing.

Abbreviation: GI, gastrointestinal.

0.01 µg/mL of TiO₂), glutathione (GSH) levels (at 0.01–1 µg/mL of TiO₂), 8-hydroxydeoxyguanosine levels (at 1 µg/mL and 10 µg/mL of TiO₂), 8-oxoguanine DNA glycosylase homologue 1 expression (at 0.001–1 µg/mL of TiO₂), and cytotoxicity (at 0.1 µg/mL, 1 µg/mL, and 10 µg/mL of TiO₂) in human embryo hepatocyte cells. These results suggest that interaction of TiO₂ NPs and PbAC may result in an increase of oxidative stress in culture cells.

Arsenic (As) is a metalloid. Chronic As exposure could cause cancer, neuropathies, and bronchopulmonary, cardiovascular, and metabolic diseases.²⁴ To reduce As pollution, TiO₂ NPs have been used as photocatalytic oxidants and/or absorption materials to remove As from water,²⁵ which implies an interaction between As and TiO₂ NPs. Due to their small diameter, large surface area, and the ability to uptake –OH ions from solution, TiO₂ NPs could adsorb metal ions through electrostatic interaction.^{26,27} Evidence shows that variables such as pH and temperature may affect the absorption and/or desorption of As (III) and As (V) by TiO₂ NPs in the aqueous solution. Pena et al²⁸ found that at 21°C–25°C and pH <8, TiO₂ NPs could be used to remove As (V) from solution through adsorption, but the maximum removal for As (III) occurred at about pH 7.5. Additionally, they demonstrated that the competing anions such as silicate, carbonate, and phosphate had a low effect on the adsorption capacities of TiO₂ NPs on As (III) and As (V) in a neutral pH range, which was in agreement with the results of Bang et al.²⁹ Niu et al found that the adsorption of As (V) was more favored in acid solution at 25°C, whereas the uptake of As (III) was preferred in alkaline solution by TiO₂ NPs at 25°C.³⁰ The maximum uptake of As (V) and As (III) was 208 mg/g (pH =3.0) and 60 mg/g (pH =7.0). Their experiments also suggested that more than 80% of As (III) and 95% of As (V) adsorbed on TiO₂ NPs could be desorbed with 1.0 M sodium hydroxide solution within 1 hour, as demonstrated by desorption tests, which was confirmed by Bang et al.²⁹ Jegadeesan et al³¹ indicated that the capacity for sorption to As by TiO₂ NP polymorphs might be affected by sorption site density, surface area (particle size), and crystalline structure. Wang et al³² stated that As toxicity on *Ceriodaphnia dubia* might increase when TiO₂ NPs (5–10 nm) interact with As in the ecosystem. They found that TiO₂ NPs less than 400 mg/L alone were nontoxic. The 24-hour median lethal concentration (LC₅₀) of As alone on *Ceriodaphnia dubia* was 3.68±0.22 mg/L. In addition, the presence of low concentrations of 50 mg/L TiO₂ NPs increased the toxicity of As significantly, and the LC₅₀ of As was also lowered to 1.43 mg/L. In summary, available studies show that pH,

temperature, composite method, and crystalline structure may all be important for the adsorption of As (III) and As (V) by TiO₂ NPs, and an alkaline environment is suitable for the desorption of As (III) and As (V). More studies are needed to investigate the toxicity effects after combination of As with TiO₂ NPs.

Copper (Cu) is an important element in human physiological processes. Exposure to low doses of Cu³³ is harmless because Cu is an essential trace element for the human body. Adverse immunotoxicological effects on human health could be caused only by an overexposure to Cu. Overexposure, especially a sublethal dose exposure of Cu, could induce immunotoxicity in mice, including cell-cycle arrest and cell death in the spleen and thymus.³⁴ Fan et al³⁵ found that TiO₂ NPs (at a concentration generally considered to be safe in the environment) remarkably enhanced the toxicity of Cu on *Daphnia magna* by increasing the bioaccumulation of Cu. In addition, they found that the Cu was adsorbed on to the TiO₂ NPs when ingested and was accumulated in the animals, thereby causing an increase in toxic effects.

Cadmium (Cd) is one of the most toxic elements to which human beings may be exposed.^{36–38} Cd compounds³⁹ are widely used in rechargeable nickel-Cd batteries. Cigarette smoke, polluted foods, and batteries are the major sources of Cd pollution. Xia et al⁴⁰ investigated the combined toxicity of cadmium chloride (CdCl₂) and TiO₂ NPs (25 nm) in human embryo kidney (HEK293T) cells. They found that cotreatment with 3.8 µm/L CdCl₂ and 7.5 µg/mL TiO₂ NPs exerted additive effects on the cellular oxidative damage by upregulation of heme oxygenase 1 gene expression, catalase activities, and malondialdehyde concentration. A combination of CdCl₂ (5.12 µm/L) and TiO₂ NPs (10.05 µg/mL) showed synergetic effects on activities of SOD and ROS concentrations. Zhang et al⁴¹ assessed the bioaccumulation of Cd in carp in the presence of TiO₂ NPs (21 nm) and found that the presence of TiO₂ NPs and the accumulation of Cd in carp was positively correlated.

Hu et al⁴² investigated the combined effects of TiO₂ NPs (21 nm) and humic acid (HA) on the bioaccumulation of Cd in zebrafish. They found that the presence of TiO₂ NPs at 5–20 mg/L in water containing HA could alter the exposure of Cd and other potential heavy metals to zebrafish. The presence of TiO₂ NPs or HA alone with Cd slightly increased the uptake rate constants of Cd in fish. TiO₂ NPs have a slightly higher uptake than HA, whereas mixtures of HA and TiO₂ NPs with Cd slightly reduced the uptake rate constants. The mechanism underlying these combined effects is unclear.

Yang et al⁴³ investigated Cd adsorption on polyacrylate-coated TiO₂ NPs, which could decrease the concentration of Cd ion in an aquatic environment and its effect on the bioavailability as well as toxicity of Cd to green algae *Chlamydomonas reinhardtii*. They found that Cd absorbed quickly on to TiO₂ NPs (anatase, 1–10 nm), reaching a steady state within 30 minutes. Interestingly, they found that the presence of TiO₂ NPs could alleviate the Cd toxicity to the green algae cells, which might be caused by TiO₂ NP adsorption on Cd²⁺, resulting in a decrease of free Cd ion in the medium and, further, its bioaccumulation in the algal cells. In addition, the electrostatic and potentially steric repulsions between TiO₂ NPs and algal cells might hinder their direct contact with each other and then prevent the internalization of TiO₂ NPs into the cells.

Taken together, toxicities of TiO₂ NPs alone have been well documented.^{10,44,45} However, knowledge of the

combined effects of TiO₂ NPs with other chemicals is limited. The existing evidence suggests that TiO₂ NPs can absorb metal ions, including Pb, As, Cu, and Cd, in the solution. Meanwhile, interaction of TiO₂ NPs and metal compounds (see Table 1) may also result in the increased toxicity demonstrated by increased oxidative stress to cells and decreased LC₅₀ to aquatic organisms.^{23,40} Therefore, more ecology investigations and biology experiments of their combined toxicity should be carried out.

Effects of TiO₂ NP interaction with organic and inorganic compounds

Bisphenol A (BPA) is an endocrine disruptor that can mimic estrogen and may lead to adverse health effects.^{46,47} Woodruff et al⁴⁸ analyzed data for 163 chemical analytes in 12 chemical classes in subsamples of 268 pregnant women from the National Health and Nutrition Examination Survey

Table 1 Studies in vitro on the interactive effects of TiO₂ NPs with chemicals or physical factors

Reference	Supplier	Characteristics of TiO ₂ NP			Dispersion method	Exposure concentration (µg/mL)
		Particle size (nm)	Crystal structure	Surface area (m ² /g)		
23	Degussa	21	80% anatase, 20% rutile	49.6	10 min ultrasonication and 30 s vortex mixing	0.001, 0.01, 0.1, 1, 10
40	Degussa	25	80% anatase, 20% rutile	50	Suspended fresh immediately before use	0, 0.25, 0.5, 0.75, 1, 1.25 TU (1 TU = 10.05 µg/mL)
49	Degussa	25–50	80% anatase, 20% rutile	50	Vortexed for 2 min, ultrasonicated for 10 min	0, 1, 5, 10
55	Degussa	25	80% anatase, 20% rutile	50	NA	0, 0.01, 0.1, 1
57	Degussa	21	80% anatase, 20% rutile	NA	Ultrasonicated for 40 min	0, 10
65	Degussa	20	rutile and anatase	50	Ultrasonicated for 15 min	50
66	Wanjin Material Corp; Degussa	4, 10, 21, 25, or 60	anatase, rutile; anatase/rutile (3:1)	NA	Freshly prepared and diluted	0, 10, 50, 100, 200
67	Degussa	20	70%–85% anatase and 30%–15% rutile	48.08	Sonicated for 30 min	0, 1, 5
68	Sigma Chemicals, Degussa	<25, <100; 31	anatase, rutile; 86% anatase, 14% rutile	NA	Sonicated for 30 min	50, 100
71	Degussa	21	25% rutile and 75% anatase	NA	Sonicated for 10–15 min	200
75	Sigma-Aldrich	NA	anatase and rutile	NA	Sonicated for 30 min	200
82	Degussa	NA	NA	NA	NA	61, 60

Abbreviations: ↑, combined effect showed a significant increase than TiO₂ NPs group and other factor group alone; ↓, combined effect showed a significant decrease than TiO₂ NPs group and other factor group alone; ↔, combined effect showed no significant difference than TiO₂ NPs group and other factor group alone; 8-OHdG, 8-hydroxydeoxyguanosine; GSH, glutathione; *HO-1* gene, *heme oxygenase 1* gene; MMP, mitochondrial membrane potential; NA, data not available; NPs, nanoparticles; OGG1, 8-oxoguanine DNA glycosylase homologue 1; ROS, reactive oxygen species; SOD, superoxide dismutase; TiO₂, titanium dioxide; UVA, ultraviolet A; DNA, deoxyribonucleic acid; MDA, malondialdehyde; CAT, catalase activities; NO, nitric oxide; PARP, poly (ADP-ribose) polymerase; MTS, 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium; NAC, N-acetyl cysteine; LDH, lactic acid dehydrogenase; mRNA, messenger ribonucleic acid; BAX, Bcl-2-associated X protein; DMEM, Dulbecco's Modified Eagle's Medium; TU, toxic unit; MEM, Minimum Essential Medium; ADP, adenosine diphosphate; min, minutes; PbAc, Plumbi Acetatis; BPA, bisphenol A; HBE, human bronchial epithelial cell; NaF, sodium fluoride; h, hours; HaCaT, a cell type belonging to an immortal human keratinocyte line; RPMI, Roswell Park Memorial Institute; BCL2, B-cell lymphoma 2 protein.

2003–04, a nationally representative sample of the US population. They found that BPA was detectable in 96% of pregnant women. Therefore, interactions between TiO₂ NPs and BPA may occur when TiO₂ NPs are used as drug carriers in the human body. Zheng et al⁴⁹ evaluated the interactive effects of TiO₂ NPs (25–50 nm) with BPA on their physiochemical properties and in vitro toxicity in human embryo hepatocytes (L-02 cells). They found that TiO₂ NPs alone (0 mg/L, 0.1 mg/L, 1 mg/L, and 10 mg/L) or BPA alone (0 µmol/L, 0.1 µmol/L, 1 µmol/L, and 10 µmol/L) did not exert significant DNA and chromosomal damage, whereas the mixture of TiO₂ NPs and BPA induced a significant increase in oxidative stress, DNA double-strand breaks, and micronuclei formation in a weak synergistic manner. An increase in intracellular levels of BPA bound by TiO₂ NPs was hypothesized to be the reason behind the synergistic toxicity. This could have been determined if the

investigators in this study used varying concentrations of TiO₂ NPs in the combination.

Dichlorodiphenyltrichloroethane (p,p'-DDT) was widely used as an effective insecticide, and had been proven to have genotoxicity, developmental toxicity, and endocrine-disruptive effects in human beings and a wide range of living organisms.^{50–52} Therefore, the degradation of DDT had attracted great attention.^{53,54} Recently, TiO₂ NPs have been tested to degrade the p,p'-DDT, which increases the risk of exposure to mixtures of TiO₂ NPs and p,p'-DDT. Shi et al⁵⁵ examined the interactive toxicities of p,p'-DDT and TiO₂ NPs (25 nm) at low concentrations in L-02 cells. The mixtures induced higher toxicity than TiO₂ NPs or p,p'-DDT alone. Combination of traces of TiO₂ NPs (0 µg/mL, 0.01 µg/mL, 0.1 µg/mL, and 1 µg/mL) and traces of p,p'-DDT (0 µmol/L, 0.001 µmol/L, 0.01 µmol/L, and 0.1 µmol/L) synergistically enhanced genotoxicity, as demonstrated by an increase in

Exposure time	Cell line	Culture medium	Combined factors/exposure condition	Combined effects
24 h	L-02	DMEM	PbAC/I µg/mL	Cell viability↓, ROS↑, GSH↑, SOD↓, 8-OHdG↑, OGGI expression↑
24 h	HEK293T	DMEM	CdCl ₂ /0, 0.25, 0.5, 0.75, 1, 1.25 TU (1 TU = 5.12 µmol/L)	<i>HO-1</i> gene express↑, OGGI expression↑, SOD↓, ROS↑, MDA↔, CAT↑
24 h	L-02	DMEM	BPA/0, 0.1, 1, 10 µmol/L	Cell viability↔, ROS↑, MDA↑, DNA double strand break↑, chromosomal damage↑
12, 24, 36 h	L-02	DMEM	p,p'-DDT/0, 0.001, 0.01, and 0.1 µmol/L	Cell viability↔, apoptosis test↔, ROS↑, MDA↑, 8-OHdG↑; DNA strand break↑; micronucleus frequency↑
72 h	I6-HBE	RPMI-1640	NaF/0, 10, 20, 30 mg/L	Cell viability↔, apoptosis test↑, SOD↓, MDA↑, NO↑
1 h	HaCaT	NA	Nitrite, UVA/nitrite: 0, 0.1, 0.5, 1, 2 mM; UVA: 365 nm, 0.6 mW/cm ² , 1 h	Cell viability↓, apoptosis test↑, protein tyrosine nitration↑
1 h	HaCaT	MEM	UVA/365 nm, 3.5 mW/cm ² , 1 h	Cell viability↓, SOD↓, ROS↑, MDA↑
0, 24, 48 h	Human peripheral blood lymphocytes	RPMI-1640	UVA/365 nm, 2.0 mW/cm ² , 0, 24, 48 h	Cell viability↓, sub-G1 phase↑, caspase-9↑, caspase-3↑, and PARP↑ MMP↓, ROS↑DNA damage↑micronucleus formation↑
4 h	HaCaT	DMEM	UVA/320–390 nm; 0, 2.5, 5.0, and 10 J/cm ²	MTS assay (A325, P25 and A25)↓, ROS↑
24 h	HaCaT	MEM	NAC, UVA/NAC: 5 mM, 2 h UVA: 365 nm, 3.5 W/cm ² , 1 h	Cell viability (UVA)↓ (UVA + NAC)↑, LDH (UVA+NAC)↓, apoptosis assay (UVA + NAC)↓, ROS (UVA + NAC)↓, MMP (UVA + NAC)↑, K6 mRNA (UVA + NAC)↑
12 h	U87-MG	DMEM	UVA/365 nm, 5 J/cm ² , 20 min	Cell viability↓, BCL2↓, BAX↑
10 min	Leukemia K562	RPMI-1640	Daunorubicin, UVA/daunorubicin: 0.14 mM, 0.2 mM UVA: 100 s	Drug accumulation↑

oxidative stress, oxidative DNA adducts, DNA breaks, and chromosomal damage in L-02 cells. The adsorption of p,p'-DDT by TiO₂ NPs was approximately 0.3 mmol/g. Synergistic genotoxicity induced by a combination of traces of p,p'-DDT and TiO₂ NPs may be a potential environmental risk factor. Sodium fluoride (NaF) and TiO₂ NPs are useful additives in household products such as toothpastes.^{3,56} Xie et al⁵⁷ examined the combined effects of NaF and TiO₂ NPs (21 nm) in human bronchial epithelial cells (16-HBE) and found that combined exposure of NaF and TiO₂ NPs could enhance the oxidative stress in 16-HBE cells. In another study, Xu et al⁵⁸ investigated the interactions of TiO₂ NPs with functional biomolecule lysozymes in culture cells. They found that lysozymes were adsorbed on to the surface of TiO₂ NPs (60 nm) via electrostatic attraction and hydrogen bonds. They therefore suggested that TiO₂ NPs might have some toxic impacts on biomolecules after interacting with biomolecule lysozymes. In summary, studies on the effects of TiO₂ NP interaction with organic or inorganic compounds are limited. The existing evidence indicates that interaction of TiO₂ NPs with BPA, p,p'-DDT, or NaF may result in enhancement of oxidative stress and further cyto or ecotoxicity.

Effects of TiO₂ NP interaction with the physical factor UVA

TiO₂ NPs possess excellent optical and electrical properties.^{59–61} Due to photocatalysis, TiO₂ NPs^{62,63} are used to degrade formaldehyde and thus improve air quality in the indoor environment. However, some studies suggest that TiO₂ NPs might be toxic under ultraviolet A (UVA), an electromagnetic radiation with wavelength range from 315 nm to 400 nm, ISO-21348. Lu et al⁶⁴ found that TiO₂ NPs (20 nm) could induce photocatalytic nitration of the protein tyrosine, which could lead to prevalent post-translational modification by TiO₂ NPs as a result of oxidative and nitrative stress. In another study, Tu et al⁶⁵ found that nitrative stress induced by TiO₂ NPs (20 nm) under UVA radiation triggered apoptotic cell death in human keratinocyte cells. This result suggests that skincare products with TiO₂ NP components may cause damage to human keratinocyte cells under UVA radiation. Xue et al⁶⁶ investigated the oxidative stress and cytotoxicity induced by different crystalline forms (anatase, rutile, and anatase/rutile) and sizes (4 nm, 10 nm, 21 nm, 25 nm, or 60 nm) of TiO₂ NPs in HaCaT cells under UVA irradiation. They found that TiO₂ NPs could induce ROS generation and toxicity in cells under UVA irradiation. Kang et al⁶⁷ also showed that TiO₂ NPs (20 nm) and UVA (0.6 mW/cm² for 1 hour) synergistically promoted

ROS generation and triggered cell apoptosis. Yin et al⁶⁸ examined the phototoxicity of TiO₂ NPs with different sizes and crystal forms (anatase and rutile) in human skin keratinocytes under UVA irradiation. They found that TiO₂ NPs are phototoxic to human skin keratinocytes, and that the phototoxicity is mediated by ROS generated during UVA radiation. Moreover, the phototoxicity of TiO₂ NPs was less with larger particle size and surface areas. Zhang et al⁶⁹ investigated the combined effects of TiO₂ NPs and UVA exposure on African clawed frogs (*Xenopus laevis*). They found that, regardless of UVA exposure, the rate of *X. laevis* survival decreased with increased concentrations of TiO₂ NPs. Exposure to 10 nm TiO₂ NPs and UVA significantly decreased the rate of *X. laevis* survival. However, exposure to 32 nm TiO₂ NPs and UVA had no statistical effect on the rate of *X. laevis* survival. This experiment suggests that toxicity is related to the size of TiO₂ NPs to some extent. Bar-Ilan et al⁷⁰ also found that TiO₂ NPs under UV produced ROS as the major phototoxic agent to the development of zebrafish. Xue et al⁷¹ examined the chemoprotective effects of N-acetylcysteine (NAC) (5 mM pretreated for 2 hours) on TiO₂ NP-induced (21 nm, 200 µg/mL for 24 hours) oxidative stress and apoptosis in human keratinocytes under UVA (3.5 mW/cm² for 1 hour). NAC,^{72–74} a nutritional supplement for cysteine donating, is widely used as an antioxidant. They found that NAC could prevent TiO₂ NP-induced oxidative stress and apoptosis in cells. The protective effects of NAC on TiO₂ NP-induced apoptosis were related to modulation of ROS and intracellular nitric oxide levels. It is worth noting that the combined effect of TiO₂ NPs and UVA may be a double-edged sword. Wang et al⁷⁵ investigated the antitumor effects of TiO₂ NPs excited with UVA irradiation both in vitro and in vivo. Their results revealed that TiO₂ NPs alone had no effect on glioma cell proliferation. However, when TiO₂ NPs were combined with UVA irradiation, the proliferation rate of cells was decreased significantly compared with controls (TiO₂ NPs alone or UVA alone). They further investigated the in vivo antitumor effects of combined TiO₂ NPs plus UVA on established glioma tumors. TiO₂ NPs plus UVA led to pronounced areas of necrosis, elevated indices of apoptosis, delayed tumor growth, and increased survival compared with the TiO₂ NPs alone or UVA alone. Moreover, the log-rank test for trend in survival analysis of tumors implanted in animals showed that the average survival duration was prolonged. Additionally, degradation, detoxification, and/or bactericidal effects of TiO₂ NPs under UVA to some pesticides or drugs in the ecosystem have been widely investigated.^{76,77} Under UV-irradiated conditions

($\lambda > 350$ nm), a TiO₂ electron excites an electron from the valance band to the conduction band and electron hole pairs are generated in the surface of TiO₂, which consists of the positive hole (h^+) and electron (e^-). The hole in the valence band has a positive redox potential and is capable of oxidizing organics, H₂O, and hydroxide ions on the surface of TiO₂ NPs, and eventually to generate hydroxyl radicals (OH). At the same time, the electron promoted from the valence band to the conduction band reduces oxygen into a superoxide radical (O₂⁻). OH, O₂⁻, and other perhydroxyl radicals have a strong oxidability to degradate pesticides, drugs, and organic substances (Figure 2). The photocatalytic effects of TiO₂ NPs have been indicated to degrade pesticides such as endosulfan and organochlorine.⁷⁸ Seitz et al⁷⁹ found that TiO₂ NPs (~100 nm; 0.2 mg/L) could reduce nearly 30% of the pirimicarb concentration under UV irradiation (40 W/m² for 15 minutes), which resulted in an almost complete removal of pirimicarb toxicity to *D. magna*. Zhao et al⁸⁰ indicated that TiO₂ NPs under UV irradiation could degrade oxytetracycline, which is widely used in both human and veterinary medicine. Higher degraded toxic byproducts were detected by a standardized bioluminescence assay of inhibition rate on *Vibrio qinghaiensis* sp.-Q67 (Q67), which indicates that a possible enhancement in ecological toxicity may occur after combination of TiO₂ NPs with oxytetracycline under UVA irradiation. These results indicate that the photocatalytic effect of TiO₂ NPs on pesticides and drugs under UVA irradiation may result in either increased or

decreased toxicity, depending on the characteristics of the byproducts.

Li et al⁸¹ demonstrated that Ag-TiO₂ NPs showed a greater synergistic bactericidal activity under UV than TiO₂ (inert) or pure Ag NPs alone to Gram-positive bacteria *Bacillus subtilis* and Gram-negative bacteria *Pseudomonas putida* at 25°C. Additionally, Song et al⁸² found that the synergistic effect of TiO₂ NPs under UV irradiation could enhance the drug accumulation dose in targeted leukemia K562 cells and inhibit multidrug resistance. Their findings suggest that combined effects of TiO₂ NPs and UVA irradiation may be beneficial for tumor treatment. TiO₂ NPs have been well investigated in recent years for enhancing the photocatalytic activity, coating, and doping of zinc oxide (ZnO) on to its surface.^{83,84} Liao et al⁸³ found that TiO₂/ZnO composite NPs had a higher photocatalytic activity in the degradation of methyl orange than both TiO₂ and shape-controlled NPs alone. Additionally, Jiang et al⁸⁵ demonstrated that a combination of the nanosized TiO₂ and ZnO powders displayed high photocatalytic activity toward the decolorization of C.I. Basic Blue 41 in water under solar radiation, and a Ti/Zn molar ratio of 1:1 showed highest photocatalytic activity.

In conclusion, TiO₂ NPs excited with UVA irradiation could induce ROS generation and thus trigger photocatalytic nitration of the protein tyrosine, oxidative stress, and eventually cell apoptosis. Increased oxidative stress damage may be the principal toxic mechanism. Combined toxicity of TiO₂ NPs under UVA may be through ROS-mediated upregulation

Figure 2 Interactive effects (degradation and absorption) of titanium dioxide (TiO₂) nanoparticles (NPs) with chemicals under ultraviolet A (UVA) radiation.

Abbreviations: OH, hydroxyl radicals; O₂, oxygen; O₂⁻, superoxide radical; H₂O, water; e⁻, electron; h⁺, the positive hole.

Table 2 Studies in vivo on the interactive effects of TiO₂ NPs with chemicals or physical factors

Reference	Supplier	Characteristics of TiO ₂ NP			Dispersion method	Living organism
		Particle size (nm)	Crystal structure	Surface area (m ² /g)		
20	Zhejiang Hongsheng Nanotechnology	50, 120	NA	NA	Sonicated for 20 min	Kun Ming mice
32	Skyspring	5–10	NA	NA	Mixed in a shaker for 24 h	<i>Ceriodaphnia dubia</i>
35	Nanomaterials Inc	13.5	Anatase	NA	Sonicated for at least 30 min	<i>Daphnia magna</i>
41	Technology Material Degussa	21	NA	50	NA	<i>Cyprinus carpio</i>
42	Evonik Degussa	21	NA	50 ± 15	Pre-equilibrated for at least 24 h	Zebrafish
43	Vivo Nano	1–10	Anatase	NA	Coated with hydrophilic sodium polyacrylate	<i>Chlamydomonas reinhardtii</i>
58	Degussa	21	NA	NA	Ultrasonicated for 10 min	<i>Micrococcus lysodeikticus</i>
69	Alfa Aesar	5, 10, 32	NA	210, 115, 45	NA	<i>Xenopus laevis</i> 0
75	Sigma-Aldrich	NA	Anatase and rutile	NA	Sonicated for 30 min	Female BALB/c nude mice
79	Degussa	21	80% anatase, 20% rutile	50 ± 15	Sonicated for 10 min	<i>D. magna</i>
80	Degussa	27	NA	50	Dispersed on 5 A or 13X surface	<i>Vibrio qinghaiensis</i> sp.-Q67

Note: ^aThe combined effects are effects with TiO₂ NPs and combined factors, comparing with effects of nTiO₂ or combined factor alone.

Abbreviations: ↓, inhibit, decrease, suppress, or delay; ↑, increase; ↔, no significant changes; GSH, glutathione; NA, data not available; NPs, nanoparticles; ROS, reactive oxygen species; SOD, superoxide dismutase; T, temperature; TiO₂, titanium dioxide; UVA, ultraviolet A; MDA, malondialdehyde; CD, cadmium; HA, humic acid; NaH₂AsO₄, sodiumarsenate; OTC, oxytetracycline; min, minutes; PbAC, Plumbi Acetatis; h, hours; d, days; pH, the acidity or basicity of an aqueous solution; nTiO₂, nano-TiO₂.

of the death receptor Fas, and activation of the preapoptotic protein Bax.⁸⁶ Meanwhile, ROS-induced p53-mediated DNA damage can lead to G₂/M cell-cycle arrest or delay and mitochondrial DNA damage.^{67,69,88} These procedures can ultimately lead to cell apoptosis. This combined toxicity could be alleviated by antioxidant NAC. The combined effects of TiO₂ NPs and UVA are a double-edged sword. Synergistic effects in ROS generation and cytotoxicity after a combination of TiO₂ NPs and UVA irradiation may be beneficial in the treatment of tumors. A combination of TiO₂ NPs and UVA irradiation may also be helpful in degradation of pesticide or drug content in the environment. The beneficial effects also include enhancement in bactericidal activity. However, degradation may also be a possible contributor. An enhancement in toxicity in the ecosystem occurs when higher toxics are degraded and byproducts are generated. Therefore, more studies are necessary to elucidate the effects of the interaction of TiO₂ NPs with different drugs or pesticides in ecosystems under UV radiation.

Summary

Research data on the effects of the interaction of TiO₂ NPs with chemicals or physical factors are emerging slowly. Experimental factors such as agglomeration/aggregation of NPs, cell type, dispersion method, or culture medium may all affect the experiment results,^{15,88–90} which may inevitably cause difficulties in interpreting these results. To give the whole picture of the experiments, we summarize in detail the combined toxicities of TiO₂ and chemicals or physical factors for all available studies in Tables 1 and 2.

The existing evidence suggests that TiO₂ NPs can absorb metals, including Pb, As, Cu, and Cd, or their ions in solution. A combination of TiO₂ NPs with metals, organic or inorganic compounds such as BPA, p,p-DDT, or NaF, and physical factors such as UVA light can result in an increase in oxidative stress and toxicity in culture cells or in aquatic animals. Oxidative stress may further induce tumor gene expression or lead to nuclear and mitochondrial DNA damage in mammalian cells. Further well-designed studies are necessary to

Exposure concentration	Exposure method	Exposure time	Exposure condition of combined factors and experiment	Combined effects ^a
5 g/kg	Oral gavage	7 d	PbAC: 500 mg/kg	Liver and kidney function↓; ROS: liver↑, kidney/cortex/hippocampus↔; MDA↔; liver/kidney: SOD↔, GSH-Px↔, cortex and hippocampus: SOD↓, GSH-Px↓
200 mg/L	In food	24 h	NaH ₂ AsO ₄ : 0, 0.45, 0.75, 1.5, 2.25, 2.5, 3, and 4.5 mg/L; pH = 7.8, T = 20°C	TiO ₂ NPs concentration: mortality at low dose↑; mortality at high dose↓
2 mg/L	In water	3 d	Copper nitrate: 10, 20, 30, 40, 50, 70 and 100 µg/L; pH = 7.6, T = 23°C	Cu ²⁺ LC50↓, metallothionein level↓
10.0 ± 1.3 mg/L	In water	0, 5, 10, 15, 20, 25 d	Cd: 3.4 ± 4.8 µg/L, 97.3 ± 6.9 µg/L; pH = 7.8, T = 23°C ± 2°C	Cd accumulation↑
5, 10, 20 mg/L	In water	0, 1, 2, 5, 8, 12, 16, 20 d	HA: 5, 10, 20 mg/L; Cd: 50 µg/L; pH = 7.0, T = 25°C ± 2°C	Uptake rate constants of Cd bioaccumulation: HA/TiO ₂ ↑, HA& TiO ₂ ↓
1, 3, 10, 30, and 100 mg/L	In water	0, 0.25, 0.5, 0.75, 1, 2, 6 h	Cd: 0, 0.1, 0.3, 0.5, 0.8, 1.0, and 3.0 mg/L; pH = 7.5 ± 0.1, T = 25°C	Free Cd ²⁺ concentration in media↓
0, 2, 4, 6, 8, 10, 15, 20, 25, 30, 40, 50, 60, 80, and 100 mg/L	In water	10 min	Lysozymes: 0, 0.42, 0.97, 1.39, 2.08 µM; pH = 7.4	Bacteriolysis activity of lysozyme↓
31, 1, 3.1, 10, 31, 100, 310, and 1000 mg/L	In water	14 d	UVA: 400 mW/m ² ; pH = 7.0–7.8	<i>X. laevis</i> survival↓; tadpole: body length↓, development stages↓
200 µg per tumor	Air pouch	12 h	UVA: 365 nm, 5 J/cm ² , 30 min	Necrosis↑, apoptosis↑, tumor growth↓, survival↑
0.02, 0.2, and 2.0 mg/L	In water	72 h	Pirimicarb: 20 µg/L; UVA: 300–400 nm, 40 ± 5 W/m ² , 15 min; T = 20°C ± 1°C	UVA and nTiO ₂ : pirimicarb concentration↓
1, 5, 10, and 15 wt%	In solution	30, 60, 270 min	OTC: 50 mg/L; UVA: 254 nm, 845 µW/cm ² , 2 h; pH = 7	(TiO ₂ /5A or TiO ₂ /13X) and UVA: inhibition rate↑

elucidate possible mechanisms of these combined effects. In addition, combined toxicity of TiO₂ NPs with ZnO/Fe₃O₄ is yet to be investigated by molecular biological and toxicological experiments. Meanwhile, studies on the combined toxicity of TiO₂ NPs with other nanosized particles in toxicology and nanomedicine are also urgently needed. Human epidemiological investigation on the combination of TiO₂ NPs with other chemicals is urgently encouraged because TiO₂ NPs are increasingly being used as drug carriers in nanomedicine. Studies on the combined effects of TiO₂ NPs with chemicals or physical factors on aquatic animals are also urgently needed. In addition, the combined effects of TiO₂ NPs with chemicals or physical factors may serve as a double-edged sword. Therefore, studies are also encouraged for TiO₂ NP application in heavy metal pollution prevention and tumor treatment.

Acknowledgments

The excellent assistance of Miss Ruth Magaye, Mrs Linda Bowman, and Mr Yaseen Habeeb in the preparation of this

article is greatly appreciated. This work was partly supported by the National Nature Science Foundation of China (Grant No 81273111), the Foundations of Innovative Research Team of Educational Commission of Zhejiang Province (T200907), the Nature Science Foundation of Ningbo City (Grant No 2012A610185), the Ningbo Scientific Project (Grant Nos 2012C5019 and SZX11073), the Scientific Innovation Team Project of Ningbo (No 2011B82014), Innovative Research Team of Ningbo (2009B21002), and KC Wong Magna Fund in Ningbo University.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Hagens WI, Oomen AG, de Jong WH, Cassee FR, Sips AJ. What do we (need to) know about the kinetic properties of nanoparticles in the body? *Regul Toxicol Pharmacol*. 2007;49:217–229.
2. Magaye R, Zhao J, Bowman L, Ding M. Genotoxicity and carcinogenicity of cobalt-, nickel- and copper-based nanoparticles. *Exp Ther Med*. 2012;4:551–561.

3. Zhao J, Castranova V. Toxicology of nanomaterials used in nanomedicine. *Toxicol Environ Health B Crit Rev*. 2011;14:593–632.
4. Cho WS, Kang BC, Lee JK, Jeong J, Che JH, et al. Comparative absorption, distribution, and excretion of titanium dioxide and zinc oxide nanoparticles after repeated oral administration. *Particle and fibre toxicology*. 2013;10:9.
5. Long TC, Tajuba J, Sama P, Saleh N, Swartz C, et al. Nanosize titanium dioxide stimulates reactive oxygen species in brain microglia and damages neurons in vitro. *Environ Health Perspect*. 2007;115:1631–1637.
6. Li SQ, Zhu RR, Zhu H, Xue M, Sun XY, et al. Nanotoxicity of TiO₂ nanoparticles to erythrocyte in vitro. *Food Chem Toxicol*. 2008;46:3626–3631.
7. Magaye R, Zhao J. Recent progress in studies of metallic nickel and nickel-based nanoparticles' genotoxicity and carcinogenicity. *Environ Toxicol Pharmacol*. 2012;34:644–650.
8. Zhao J, Bowman L, Zhang X, Vallyathan V, Young SH, et al. Titanium dioxide (TiO₂) nanoparticles induce JB6 cell apoptosis through activation of the caspase-8/Bid and mitochondrial pathways. *J Toxicol Environ Health A*. 2009;72:1141–1149.
9. Buzea C, Pacheco II, Robbie K. Nanomaterials and nanoparticles: sources and toxicity. *Biointerphases*. 2007;2:17–71.
10. Jawad H, Boccacini AR, Ali NN, Harding SE. Assessment of cellular toxicity of TiO₂ nanoparticles for cardiac tissue engineering applications. *Nanotoxicology*. 2011;5:372–380.
11. Lee KP, Trochimowicz HJ, Reinhardt CF. Pulmonary response of rats exposed to titanium dioxide (TiO₂) by inhalation for two years. *Toxicol Appl Pharmacol*. 1985;79:179–192.
12. Ze Y, Hu R, Wang X, Li B, Su J, et al. Neurotoxicity and gene-expressed profile in brain injured mice caused by exposure to titanium dioxide nanoparticles. *J Biomed Mater Res A*. March 27, 2013;00A:3–9 [Epub ahead of print.]
13. Iavicoli I, Leso V, Bergamaschi A. Toxicological effects of titanium dioxide nanoparticles: a review of in vivo studies. *J Nanomater*. 2012;2012:5.
14. Magdolenova Z, Collins AR, Kumar A, Dhawam A, Stone V, et al. Mechanisms of genotoxicity: review of recent in vitro and in vivo studies with engineered nanoparticles. *Nanotoxicology*. 2013;1–73.
15. Shi H, Magaye R, Castranova V, Zhao J. Titanium dioxide nanoparticles: a review of current toxicological data. *Part Fibre Toxicol*. 2013;10:15.
16. Sharma VK. Aggregation and toxicity of titanium dioxide nanoparticles in aquatic environment: a review. *J Environ Sci Health A Tox Hazard Subst Environ Eng*. 2009;44:1485–1495.
17. Hartmann NB, Legros S, Von der Kammer F, Hofmann T, Baun A. The potential of TiO₂ nanoparticles as carriers for cadmium uptake in *Lumbriculus variegatus* and *Daphnia magna*. *Aquat Toxicol*. 2012;118–119:1–8.
18. Tavallali H. Alumina-coated magnetite nanoparticles for solid phase extraction of Cd in water samples. *Chem Tech*. 2011;3:1647–1651.
19. Mashhadizadeh MH, Karami Z. Solid phase extraction of trace amounts of Ag, Cd, Cu, and Zn in environmental samples using magnetic nanoparticles coated by 3-(trimethoxysilyl)-1-propanthiol and modified with 2-amino-5-mercapto-1,3,4-thiadiazole and their determination by ICP-OES. *J Hazard Mater*. 2011;190:1023–1029.
20. Zhang R, Niu Y, Li Y, Zhao C, Song B, et al. Acute toxicity study of the interaction between titanium dioxide nanoparticles and lead acetate in mice. *Environ Toxicol Pharmacol*. 2010;30:52–60.
21. Bakircioglu Y, Bakircioglu D, Akman S. Biosorption of lead by filamentous fungal biomass-loaded TiO₂ nanoparticles. *J Hazard Mater*. 2010;178:1015–1020.
22. Kalfa OM, Yalcinkaya Ö, Türker AR. Synthesis of nano B₂O₃/TiO₂ composite material as a new solid phase extractor and its application to preconcentration and separation of cadmium. *J Hazard Mater*. 2009;166:455–461.
23. Du H, Zhu X, Fan C, Xu S, Wang Y, et al. Oxidative damage and OGG1 expression induced by a combined effect of titanium dioxide nanoparticles and lead acetate in human hepatocytes. *Environmental Toxicology*. 2011;27:590–597.
24. Barchowsky A, Cartwright IL, Reichard JF, Futscher BW, Lantz RC. Arsenic toxicology: translating between experimental models and human pathology. *Environ Health Perspect*. 2011;119:1356.
25. Guan X, Du J, Meng X, Sun Y, Sun B, et al. Application of titanium dioxide in arsenic removal from water: a review. *J Hazard Mater*. 2012;215–216:1–16.
26. Liu Y, Liang P, Guo L, Lu HB. Study on the adsorption behavior of heavy metal ions on nanometer TiO₂ supported on silica gel [in Chinese]. *Acta Chim Sin*. 2005;63:312–316.
27. Bleam WF, McBride MB. The chemistry of adsorbed Cu (II) and Mn (II) in aqueous titanium dioxide suspensions. *J Colloid Interface Sci*. 1986;110:335–346.
28. Pena ME, Korfiatis GP, Patel M, Lippincott L, Meng X. Adsorption of As (V) and As (III) by nanocrystalline titanium dioxide. *Water Research*. 2005;39:2327–2337.
29. Bang S, Patel M, Lippincott L, Meng X. Removal of arsenic from groundwater by granular titanium dioxide adsorbent. *Chemosphere*. 2005;60:389–397.
30. Niu HY, Wang JM, Shi YL, et al. Adsorption behavior of arsenic onto protonated titanate nanotubes prepared via hydrothermal method. *Micropor Mesopor Mat*. 2009;122:28–35.
31. Jegadeesan G, Al-Abed SR, Sundaram V, Choi H, Scheckel KG, et al. Arsenic sorption on TiO₂ nanoparticles: size and crystallinity effects. *Water Research*. 2010;44:965–973.
32. Wang D, Hu J, Irons DR, Wang J. Synergistic toxic effect of nano-TiO₂ and As (V) on *Ceriodaphnia dubia*. *STOTEN*. 2011;409:1351–1356.
33. Malkin R, Malmström BG. The state and function of copper in biological systems. *Adv Enzymol Relat Areas Mol Biol*. 1970:177–244.
34. Mitra S, Keswani T, Dey M, Bhattacharya S, Sarkar S, et al. Copper-induced immunotoxicity involves cell cycle arrest and cell death in the spleen and thymus. *Toxicology*. 2012;293:1–3.
35. Fan W, Cui M, Liu H, Wang C, Shi Z, et al. NanoTiO₂ enhances the toxicity of copper in natural water to *Daphnia magna*. *Environ Pollut*. 2011;159:729–734.
36. Kjellström T. Mechanism and epidemiology of bone effects of cadmium. *IARC Scientific Publications*. 1992:301.
37. Nordberg GF, Herber RFM, Alessio L. Cadmium in the human environment: toxicity and carcinogenicity. International Agency for Research on Cancer; 1992.
38. Flick D, Kraybill H, Dlmittroff J. Toxic effects of cadmium: a review. *Environmental Research*. 1971;4:71–85.
39. Järup L. Hazards of heavy metal contamination. *British Medical Bulletin*. 2003;68:167–182.
40. Xia B, Chen J, Zhou Y. Cellular oxidative damage of HEK293T cells induced by combination of CdCl₂ and Nano-TiO₂. *Journal of Huazhong University of Science and Technology Medical Sciences*. 2011;31:290–294.
41. Zhang X, Sun H, Zhang Z, Niu Q, Chen Y, et al. Enhanced bioaccumulation of cadmium in carp in the presence of titanium dioxide nanoparticles. *Chemosphere*. 2007;67:160–166.
42. Hu X, Chen Q, Jiang L, Yu Z, Jiang D, et al. Combined effects of titanium dioxide and humic acid on the bioaccumulation of cadmium in zebrafish. *Environ Pollut*. 2011;159:1151–1158.
43. Yang WW, Miao AJ, Yang LY. Cd²⁺ Toxicity to a green alga *Chlamydomonas reinhardtii* as influenced by its adsorption on TiO₂ engineered nanoparticles. *PLoS One*. 2012;7:e32300.
44. Shukla RK, Kumar A, Gurbani D, Pandey AK, Singh S, et al. TiO₂ nanoparticles induce oxidative DNA damage and apoptosis in human liver cells. *Nanotoxicology*. 2011:1–13.
45. Shukla RK, Sharma V, Pandey AK, Singh S, Sultana S, et al. ROS-mediated genotoxicity induced by titanium dioxide nanoparticles in human epidermal cells. *Toxicol In Vitro*. 2011;25:231–241.

46. Huang Y, Wong C, Zheng J, Bouwman H, Barra R, et al. Bisphenol A (BPA) in China: a review of sources, environmental levels, and potential human health impacts. *Environment International*. 2011;42:91–99.
47. Alexander HC, Dill DC, Smith LW, Guiney PD, Dorn P. Bisphenol A: acute aquatic toxicity. *Environ Toxicol Chem*. 1988;7:19–26.
48. Woodruff TJ, Zota AR, Schwartz JM. Environmental chemicals in pregnant women in the United States: NHANES 2003–2004. *Environ Health Perspect*. 2011;119:878–885.
49. Zheng D, Wang N, Wang X, Tang Y, Zhu L, et al. Effects of the interaction of TiO₂ nanoparticles with bisphenol A on their physicochemical properties and in vitro toxicity. *J Hazard Mater*. 2012;199–200:426–432.
50. Cohn BA, Wolff MS, Cirillo PM, Sholtz RI. DDT and breast cancer in young women: new data on the significance of age at exposure. *Environ Health Perspect*. 2007;115:1406.
51. Rogan WJ, Chen A. Health risks and benefits of bis (4-chlorophenyl)-1,1,1-trichloroethane (DDT). *Lancet*. 2005;366:763–773.
52. Rogan WJ, Ragan NB. Evidence of effects of environmental chemicals on the endocrine system in children. *Pediatrics*. 2003;112:247–252.
53. Zoro J, Hunter JM, Eglinton G, Ware G. Degradation of p,p'-DDT in reducing environments. *Nature*. 1974;247:235–237.
54. Garrison AW, Nzengung VA, Avants JK, Ellington JJ, Jones WJ, et al. Photodegradation of p,p'-DDT and the enantiomers of o,p'-DDT. *Environmental Science and Technology*. 2000;34:1663–1670.
55. Shi Y, Zhang JH, Jiang M, Zhu LH, Tan HQ, et al. Synergistic genotoxicity caused by low concentration of titanium dioxide nanoparticles and p, p'-DDT in human hepatocytes. *Environ Mol Mutagen*. 2010;51:192–204.
56. O'Mullane DM, Kavanagh D, Ellwood RP, Chesters RK, Schafer F, et al. A three-year clinical trial of a combination of trimetaphosphate and sodium fluoride in silica toothpastes. *J Dent Res*. 1997;76:1776–1781.
57. Xie C, Liang GY, Ye B, Pu YP. Combined effects of sodium fluoride and nano-TiO₂ on human bronchial epithelial cells. *Journal of Environmental and Occupational Medicine*. 2009;26:242–244.
58. Xu Z, Liu XW, Ma YS, Gao HW. Interaction of nano-TiO₂ with lysozyme: insights into the enzyme toxicity of nanosized particles. *Environ Sci Pollut Res Int*. 2010;17:798–806.
59. Hui YLS. Research progress and application of photocatalysis of TiO₂. *Materials Review*. 2000;12:23–25.
60. Kwon S, Fan M, Cooper AT, Yang H. Photocatalytic applications of micro-and nano-TiO₂ in environmental engineering. *Crit Rev Environ Sci Technol*. 2008;38:197–226.
61. Aarthi T, Madras G. Photocatalytic degradation of rhodamine dyes with nano-TiO₂. *Ind Eng Chem Res*. 2007;46:7–14.
62. Yu H, Zhang K, Rossi C. Experimental study of the photocatalytic degradation of formaldehyde in indoor air using a nano-particulate titanium dioxide photocatalyst. *Indoor and Built Environment*. 2007;16:529–537.
63. Wu YP, Zhang WM, Ma CF, Lu YW, Liu L. Photocatalytic degradation of formaldehyde by diffuser of solar light pipe coated with nanometer titanium dioxide thin films. *Science China Technological Sciences*. 2010;53:150–154.
64. Lu N, Zhu Z, Zhao X, Tao R, Yang X, et al. Nano titanium dioxide photocatalytic protein tyrosine nitration: a potential hazard of TiO₂ on skin. *Biochem Biophys Res Commun*. 2008;370:675–680.
65. Tu M, Huang Y, Li HL, Gao ZH. The stress caused by nitrite with titanium dioxide nanoparticles under UVA irradiation in human keratinocyte cell. *Toxicology*. 2012;299:60–68.
66. Xue C, Wu J, Lan F, Liu W, Yang X, et al. Nano titanium dioxide induces the generation of ROS and potential damage in HaCaT cells under UVA irradiation. *J Nanosci Nanotechnol*. 2010;10:8500–8507.
67. Kang SJ, Lee YJ, Kim BM, Choi YJ, Chung HW. Cytotoxicity and genotoxicity of titanium dioxide nanoparticles in UVA-irradiated normal peripheral blood lymphocytes. *Drug Chem Toxicol*. 2011;34:277–284.
68. Yin JJ, Liu J, Ehrenshaft M, Roberts JE, Fu PP, et al. Phototoxicity of nano titanium dioxides in HaCaT keratinocytes – generation of reactive oxygen species and cell damage. *Toxicol Appl Pharmacol*. 2012;263:81–88.
69. Zhang J, Wages M, Cox SB, Maul JD, Li Y, et al. Effect of titanium dioxide nanomaterials and ultraviolet light coexposure on African clawed frogs (*Xenopus laevis*). *Environ Toxicol Chem*. 2012;31:176–183.
70. Bar-Ilan O, Louis KM, Yang SP, Pedersen JA, Hamers RJ, et al. Titanium dioxide nanoparticles produce phototoxicity in the developing zebrafish. *Nanotoxicology*. 2012;6:670–679.
71. Xue C, Liu W, Wu J, Yang X, Xu H. Chemoprotective effect of N-acetylcysteine (NAC) on cellular oxidative damages and apoptosis induced by nano titanium dioxide under UVA irradiation. *Toxicology In Vitro*. 2011;25:110–116.
72. Smilkstein MJ, Knapp GL, Kulig KW, Rumack BH. Efficacy of oral N-acetylcysteine in the treatment of acetaminophen overdose. Analysis of the national multicenter study (1976 to 1985). *N Engl J Med*. 1988;319:1557.
73. Aruoma OI, Halliwell B, Hoey BM, Butler J. The antioxidant action of N-acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical, superoxide, and hypochlorous acid. *Free Radical Biology and Medicine*. 1989;6:593–597.
74. Pannu N, Manns B, Lee H, Tonelli M. Systematic review of the impact of N-acetylcysteine on contrast nephropathy. *Kidney Int*. 2004;65:1366–1374.
75. Wang C, Cao S, Tie X, Qiu B, Wu A, et al. Induction of cytotoxicity by photoexcitation of TiO₂ can prolong survival in glioma-bearing mice. *Mol Biol Rep*. 2011;38:523–530.
76. Moon J, Yun CY, Chung K-W, Kang M-S, Yi J. Photocatalytic activation of TiO₂ under visible light using Acid Red 44. *Catalysis Today*. 2003;87:77–86.
77. Srinivasan C, Somasundaram N. Bactericidal and detoxification effects of irradiated semiconductor catalyst, TiO₂. *Current Science-Bangalore*. 2003;85:1431–1438.
78. Thomas J, Kumar KP, Chitra K. Synthesis of Ag doped nano TiO₂ as efficient solar photocatalyst for the degradation of endosulfan. *Adv Sci Lett*. 2011;4:108–114.
79. Seitz F, Bundschuh M, Dabrunz A, Bandow N, Schaumann GE, et al. Titanium dioxide nanoparticles detoxify pirimicarb under UV irradiation at ambient intensities. *Environ Toxicol Chem*. 2012;31:518–523.
80. Zhao C, Deng H, Li Y, Liu Z. Photodegradation of oxytetracycline in aqueous by 5A and 13X loaded with TiO₂ under UV irradiation. *J Hazard Mater*. 2010;176:884–892.
81. Li M, Noriega-Trevino ME, Nino-Martinez N, Marambio-Jones C, Wang J, et al. Synergistic bactericidal activity of Ag-TiO₂ nanoparticles in both light and dark conditions. *Environ Sci Technol*. 2011;45:8989–8995.
82. Song M, Zhang R, Dai Y, Gao F, Chi H, et al. The in vitro inhibition of multidrug resistance by combined nanoparticulate titanium dioxide and UV irradiation. *Biomaterials*. 2006;27:4230–4238.
83. Liao D, Badour C, Liao B. Preparation of nanosized TiO₂/ZnO composite catalyst and its photocatalytic activity for degradation of methyl orange. *J Photochem Photobiol A Chem*. 2008;194:11–19.
84. Panigrahi S, Basak D. Core-shell TiO₂@ ZnO nanorods for efficient ultraviolet photodetection. *Nanoscale*. 2011;3:2336–2341.
85. Jiang Y, Sun Y, Liu H, Zhu F, Yin H. Solar photocatalytic decolorization of CI Basic Blue 41 in an aqueous suspension of TiO₂-ZnO. *Dyes Pigm*. 2008;78:77–83.
86. Yoo K-C, Yoon C-H, Kwon D, Hyun K-H, Woo SJ, et al. Titanium dioxide induces apoptotic cell death through reactive oxygen species-mediated Fas upregulation and Bax activation. *Int J Nanomedicine*. 2012;7:1203.
87. Sanders K, Degn LL, Mundy WR, Zucker RM, Dreher K, et al. in vitro phototoxicity and hazard identification of nano-scale titanium dioxide. *Toxicol Appl Pharmacol*. 2012;258:226–236.

88. Lankoff A, Sandberg WJ, Wegierek-Ciuk A, Lisowska H, Refsnes M, et al. The effect of agglomeration state of silver and titanium dioxide nanoparticles on cellular response of HepG2, A549 and THP-1 cells. *Toxicol Lett*. 2012;208:197–213.
89. Ekstrand-Hammarström B, Akfur CM, Andersson PO, Lejon C, Österlund L, et al. Human primary bronchial epithelial cells respond differently to titanium dioxide nanoparticles than the lung epithelial cell lines A549 and BEAS-2B. *Nanotoxicology*. 2012;6:623–634.
90. Magdolenova Z, Bilaničová D, Pojana G, Fjellsbø LM, Hudcová A, et al. Impact of agglomeration and different dispersions of titanium dioxide nanoparticles on the human related in vitro cytotoxicity and genotoxicity. *J Environ Monit*. 2012;14:455–464.

International Journal of Nanomedicine

Dovepress

Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-nanomedicine-journal>