Emerging targets in human lymphoma: targeting the *MYD88* mutation

James Q Wang*
Yogesh S Jeelall*
Keisuke Horikawa*

Department of Immunology, The John
Curtin School of Medical Research,
Australian National University,
Canberra, ACT, Australia

*All authors contributed equally
to this manuscript

Abstract: B cell neoplasms co-opt the molecular machinery of normal B cells for their survival. Technological advances in cancer genomics has significantly contributed to uncovering the root cause of aggressive lymphomas, revealing a previously unknown link between TLR signaling and B cell neoplasm. Recurrent oncogenic mutations in *MYD88* have been found in 39% of the activated B cell-like subtype of diffuse large B cell lymphoma (ABC DLBCL). Interestingly, 29% of ABC DLBCL have a single amino acid substitution of proline for the leucine at position 265 (L265P), and the exact same variant has also been identified in a number of lymphoid malignancies. The *MYD88* L265P variant was recently identified in 90% of Wadenstrom's macroglobulinemia patients. These recent developments warrant the need for novel diagnostic tools as well as targeted therapeutics. In this review, we discuss the physiological functions of *MYD88* and focus on its role in B cell lymphomas, evaluating the potential for targeting oncogenic *MYD88* in lymphoma.

Keywords: *MYD88*, L265P mutation, lymphoma, targeted therapy

Introduction

From one of the earliest detailed descriptions of lymphoma cases by Hodgkin in 1832, it was already evident that this group of cancer is very diverse.¹ Such heterogeneity poses significant challenges to the effective diagnosis, management and study of lymphomas. Following decades of progress in the understanding of the biology of white blood cells and 'the hallmarks of cancers', we now know lymphomas are characterized by neoplastic transformation of lymphocytes at various differentiation stages.^{2,3} Given the diversity in the subsets of lymphocytes and the numerous differentiation stages, from the common hematopoietic stem cell precursor to distinct differentiated states, the diagnosis, treatment and study of lymphoid neoplasms remain central clinical challenges.

The current classification of lymphomas resulted from a major collaborative effort by the World Health Organization (WHO) synthesizing information about the immunophenotype, genetic features, and clinical characteristics, along with the traditionally used cell/tissue morphology to define specific clinically relevant diseases.⁴ The WHO classification broadly segregates neoplasms based on myeloid and lymphoid lineages, followed by sub-categorization into functional or cell differentiation stages of the normal counterpart of each neoplasm. The 2008 WHO classification lists more than one hundred tumors of the hematopoietic and lymphoid tissues, for most of which the underlying causes are still unknown.

Correspondence: Yogesh S Jeelall
Immunogenomics Laboratory,
Department of Immunology, The John
Curtin School of Medical Research,
The Australian National University,
Building 131, Garran Road,
Acton, ACT 0200, Australia
Tel +612 6125 1393
Fax +612 6125 2595
Email yogesh.jeelall@anu.edu.au

Lymphoid neoplasms are the sixth most common cancer worldwide, with close to one million new cases expected to be diagnosed each year.⁵ The most common hematopoietic tumors diagnosed are non-Hodgkin lymphoma, leukemia, multiple myeloma and Hodgkin's lymphoma. B lymphocyte neoplasms account for about 90% of all newly diagnosed cases, among which diffuse large B cell lymphoma (DLBCL) and follicular lymphoma are the most prevalent.⁶

B cell lymphomas are thought to co-opt the molecular features of normal B cells for their survival, such that the phenotype of malignant B cells mirror the state of differentiation from which they originate. During B cell development, B cells express a number of DNA-modifying enzymes such as recombinase-activating gene (RAG1 and RAG2), which primarily serves to increase diversity of antibodies in the repertoire.⁷ A side consequence of such enzymes is the generation of chromosomal translocations, which may contribute to malignancy.⁷ Another stage of differentiation at which B cells are very susceptible to genomic alterations is the transient germinal centre (GC) stage.⁸ During the GC stage, B cells express activation-induced cytidine deaminase (AID), a DNA-modifying enzyme that is required for somatic hypermutation and class switching of antibodies.⁹ The off-target effect of AID may also contribute to the oncogenic load in B cells.^{10–12} Thus, GC B cells may give rise to several types of lymphoma, including the diffuse-large B cell lymphoma, follicular lymphoma and Burkitt's lymphoma.¹³

Diffuse large B cell lymphoma (DLBCL) is one of the most common forms of lymphoma, accounting for 30%–40% of all newly diagnosed cases.⁶ DLBCL is also currently one of the least curable lymphoma, with about 50% success using a combination of chemotherapy and rituximab.¹⁴ With the advent of genome-wide gene expression profiling, DLBCL has been subdivided into three molecular subtypes.¹⁵ The activated B cell (ABC), germinal-center B cell (GCB) and the primary mediastinal B cell lymphoma (PMBL) subtypes are histologically indistinguishable, but differ in the expression of hundreds of signature genes.¹⁵ The subdivision of DLBCL holds promise for better diagnosis and improved treatment regimes, even though the use of gene expression profiling is yet to be translated into clinical practice.

Among the three subtypes of DLBCL, the ABC subtype has been associated with the lowest success rates following standard treatment regimes.¹⁴ Interestingly, gene expression profiling and drug inhibition studies revealed that the ABC subtype has a striking dependence on signaling pathways activating the transcription factor NFκB.^{15,16} The constitutive NFκB activation in ABC DLBCL could contribute to

the poor response following chemotherapy as the targets of this family of transcription factors prevent apoptosis.¹⁷ These findings emphasized the need for the development of therapeutics targeting NFκB signaling for the treatment of aggressive lymphomas.

A recent wave of progress in cancer genomics triggered by next-generation sequencing technologies have significantly contributed to uncovering the root cause of the high NFκB activity in ABC DLBCL (Figure 1). The survival of this aggressive lymphoma subtype relies on signaling from the antigen receptor to the NFκB transcription factors, with CARD11, BCL10 and MALT1 being essential components of the signaling apparatus.¹⁶ In approximately 10% of patients, gain-of-function mutations in the *CARD11* oncogene have been found to activate NFκB and prolong cell survival.¹⁸ In addition, about 20% of ABC lymphomas have mutations in *CD79A* or *CD79B*, which are rare or absent in GCB and other lymphoma subtypes.¹⁹ Loss of function mutations resulting in the inactivation of A20, a negative regulator of NFκB signaling, has been found to occur in 25% of ABC lymphomas.^{20–22} Crippling the activity of A20 increases the activity of NFκB signaling in malignant B cells.²³

More recently, high-throughput RNA resequencing of DLBCL has identified recurrent oncogenic mutations in *MYD88* in 39% of ABC DLBCL tumors.^{24,25} These findings established a previously unknown link between TLR signaling and B cell lymphoma. Interestingly, 29% of ABC DLBCL have a single amino acid substitution of proline for the leucine at position 265 (L265P) in the TIR domain.^{24,25} The *MYD88* L265P variant has also been identified in a number of lymphoid malignancies (Table 1). The *MYD88* L265P variant was recently identified in about 90% of Wadenstrom's macroglobulinemia patients, revealing a central pathogenic feature of this tumor.²⁶ These recent developments warrant the need for novel diagnostic tools as well as targeted therapeutics. In this review, we discuss the physiological functions of *MYD88* and focus on its role in B cell lymphomas, evaluating the potential for targeting oncogenic *MYD88* in lymphoma.

Physiological function of MYD88

MYD88 was originally identified as a myeloid differentiation primary response gene in hematopoiesis.²⁷ Following treatment of myeloid cell precursors with interleukin 6 (IL6), the levels of *MYD88* transcript was found to increase as cells terminally differentiated.²⁷ *MYD88* is essential for the mammalian innate immune response. Individuals with *MYD88* deficiency suffer life threatening recurrent pyogenic

Figure 1 Oncogenic mutations targeting the NFκB pathway. Oncogenic mutations frequently target the MYD88, CD79, and CARD11 (part of the CMB complex) in aggressive lymphomas. Consequences of these mutations include disruption to normal cellular signal transduction events such as protein phosphorylation, ubiquitylation or deubiquitylation, which converge onto aberrant NFκB activity, a hallmark of lymphomas with L265P MYD88. Specific inhibitors targeting (1) MYD88, (2) IRAK4, (3) Toll-like receptor (TLR), (4) NFκB, and BTK are currently in clinical trials.

Note: Targeting these molecular pathways may provide effective treatment to patients.

Abbreviations: AP-1, activator protein 1; BCR, B cell receptor; BTK, Bruton tyrosine kinase; CBM, CARD11-BCL10-MALT1 complex; CD79, cluster of differentiation 79; Fyn, Src family protein tyrosine kinase Fyn; Lyn, Src family protein tyrosine kinase Lyn; IκBα, inhibitor of NFκB alpha; IKK, inhibitor of κB alpha kinase; IRAK1, interleukin 1 receptor associated kinase 1; IRAK4, interleukin 1 receptor associated kinase 4; MAPK, mitogen activated protein kinase; MYD88, myeloid differentiation factor 88; NFκB, nuclear factor kappa-light-chain-enhancer of activated B cells; Syk, spleen tyrosine kinase; TAB1, TGF beta activated kinase 1 binding protein 1; TAB2, TGF beta activated kinase 2; TAK1, TGF beta activated kinase 1; TGF, transforming growth factor; TLR, Toll like receptor; TNF, tumor necrosis factor; TRAF6, TNF receptor associated factor 6.

bacterial infection by *Streptococcus pneumonia*, *Staphylococcus aureus* and *Pseudomonas aeruginosa*, suggesting that MYD88 plays a crucial role in the innate immune response in the Toll/IL1 receptor pathways.^{28,29} Furthermore, mice lacking MYD88 also show impaired Toll receptor and IL1/IL18 responses in addition to defects in T cell proliferation and Th1 response.³⁰ These defects result from the inability of signals to be transmitted from ligand activated receptors to the NFκB and JNK pathways.³⁰ Thus, MYD88 plays a central role as an adaptor molecule to transducing signals from receptors such as Toll-like receptor/IL-1 receptor to transcription factors.

A homolog of MYD88 in *Drosophila*, known as Tube, was initially described as an adaptor protein for Toll-Dorsal signaling and is essential for establishing dorsal-ventral

polarity during embryogenesis.³¹ Moreover, MYD88 deficient *Drosophila* flies have crippled defense against fungal and microbial infections, and mammalian models of MYD88 deficiency have drastically poor defense against a plethora of pathogens.^{32,33} Given the crucial role of MYD88 in the immune system of a wide range of organisms, it is not surprising that this adaptor molecule has been evolutionarily conserved.

Toll like receptors (TLR) play essential roles as the danger sensing molecular detector in an innate immune response.³⁴ TLRs are type I transmembrane protein that share homology with the interleukin-1 receptor. In total, ten different members of TLRs are differentially expressed amongst different immune cell subtypes, each responding to a different type of stimulus.³⁵

Table I Frequency of somatic *MYD88* mutations in B cell neoplasm

Disease	<i>MYD88</i> mutation	Frequency (case/sample)	Reference
ABC DLBCL	L265P	29% [#]	24
ABC DLBCL	Others	10% [#]	24
BL	Other	5% [#]	24
CBCL	L265P	69% (11/16)	70
CLL	L265P	2.9% (9/310)	71
CLL	M232T	2.2% (2/91)	72
CLL	P258L	1.1% (1/91)	72
CLL	L265P	6.6% (6/91)	72
GCB DLBCL	Other	6% [#]	24
IgM MGUS	L265P	10% (2/21)	73
LPL	L265P	91% (49/54)	73
MALT	L265P	9% [#]	24
MALT	L265P	3.8% (2/53)	74
MALT	27 bp deletion*	1.9% (1/53)	74
MZL	L265P	6.5% (3/46)	75
PCNSL	L265P	36% (5/14)	76
PCNSL	L103L	7% (1/14)	76
PCNSL	Q143E	7% (1/14)	76
PCNSL	L265P	38% (11/29)	77
SMZL	L265P	13% (6/46)	78
SMZL	L265P	5.1% (6/117)	79

Notes: [#]Percentage in published text inconsistent with calculated percentage from biopsy number; *deletion occurred between gene sequence 1039–1065, resulting in amino acid deletion between V286–T294.

Abbreviations: ABC-DLBCL, activated B-cell like diffuse large B-cell lymphoma; BL, Burkitt's lymphoma; CBCL, cutaneous diffuse large B-cell lymphoma (leg type); CLL, chronic lymphocytic leukemia; GCB-DLBCL, germinal center B-cell like diffuse large B cell lymphoma; IgM MGUS, IgM monoclonal gammopathy of undetermined significance; LPL, lymphoplasmacytic lymphoma (Waldenström's Macroglobulinemia or non-IgM LPL); MALT, gastric mucosa-associated lymphoid tissue lymphoma; MZL, marginal zone lymphoma; PCNSL, primary central nervous system lymphoma; SMZL, splenic marginal zone lymphoma.

Different combinations of adaptor molecules create distinct signaling platforms, which recruit additional signal transduction molecules giving rise to a range of responses governed by differential gene expression.³⁴ Signaling by all TLRs, with the exception of TLR3, requires MYD88 as an adaptor molecule.³⁴

During bacterial infections, macrophages form an important first line of defense as part of the innate immune response. Macrophages can be potently activated by lipopolysaccharide (LPS), a major component of the Gram-negative bacteria outer membrane, through the stimulation of Toll-like receptor 4.³⁴ Stimulated macrophages produce various cytokines such as tumor necrosis factor alpha (TNF α), IL1, IL6/10 and inflammatory effector chemokines.³⁶ By using MYD88 deficient mice, Akira and colleagues elucidated the role of MYD88 in macrophage activation in response to endotoxin.³⁷ MYD88 deficient mice fail to produce proinflammatory cytokines such as TNF α , IL1 and IL6 after LPS challenge.³⁷ Cultured macrophages from MYD88 deficient

mice show no increase in mRNA level in proinflammatory cytokines upon LPS treatment, highlighting the importance of MYD88 in response to bacterial infection.³⁷

MYD88 signaling in B lymphocytes

B lymphocytes have also been shown to respond to LPS through their Toll-like receptors, resulting in proliferation and production of cytokines such as IL1, IL6, IL8 and TNF α .³⁸ LPS activated B cells also enhance their antigen presentation capacity through increased MHC II expression as well as secretion of large amounts of LPS-neutralizing antibodies, such that in response to LPS MYD88 deficient B cells have impaired MHC II upregulation as well as poor proliferation and antibody secretion.^{37,38} The response following recognition of bacterial DNA by TLR9 through a specific CpG motif,^{39,40} requires MYD88 as B cells without the adaptor molecule fail to proliferate in response to CpG DNA.⁴¹ MYD88 plays central role in the response of B cells following various stimuli through the Toll-like receptors.

MYD88 acts as a key adaptor protein linking danger signals from Toll-like receptors to transcription factors, which regulate cellular gene expression. Molecular studies revealed that MYD88 forms a protein complex with interleukin 1 receptor (IL1R) and interleukin 1 receptor-associated kinase (IRAK), a serine threonine protein kinase, in the presence of IL1.^{42,43} MYD88 is first recruited to the cytoplasmic tail of IL1R or TLRs following their engagement via homophilic TIR interactions and the formation of a homodimer.⁴⁴ IRAK4 is then recruited to the site of activation through the interactions between the death domains, resulting in the activation of IRAK4 and the phosphorylation of the downstream protein kinase IRAK1.⁴² Subsequently, phosphorylated IRAK1 associates with TRAF6 in the cytoplasm and leads to the activation of the NF κ B and MAPK pathways (Figure 1).³⁴ Through a poorly defined mechanism, following IRAK1 mediated phosphorylation, TRAF6 disassociates from the receptor complex and forms a cytoplasmic complex that consists of TRAF6/TAB2/TAK1.³⁴ Activated TAK1 phosphorylates both the β subunit of IKK and MAPK kinase 6.³⁴ Activated IKK β in turn phosphorylate I κ B α , leading to its ubiquitylation and proteasomal degradation, allowing the NF κ B dimer to translocate to the nucleus and activate gene transcription that regulates cell activation, proliferation and immune responses.³⁴

Oncogenic MYD88 in lymphoma

MYD88 forms an important link in the activation and proliferation of B cells under a number of different extra-

cellular stimuli. It is thus not surprising that defects in this critical signal relay molecule may result in pathology in B cell activation and proliferation that result from aberrant NF κ B and MAPK activity.³⁴ ABC DLBCL has characteristically constitutive NF κ B activity that enhances the proliferation and survival of the affected B cell populations. Sequencing studies of DLBCL samples and RNA interference screens using human lymphoma cell lines revealed *MYD88* mutations are present in 39% of samples.²⁴ MYD88 was found to be required for the survival of the cell lines through constitutive activation of NF κ B signaling.²⁴

The MYD88 L265P variant was recently identified in about 90% of Wadenstrom's macroglobulinemia patients, constituting a significant clinical feature for this disease.²⁶ Albeit at lower frequencies, L265P MYD88 was also found in cases of chronic lymphocytic leukemia, splenic marginal zone lymphoma, primary central nervous system lymphoma and gastric mucosa-associated lymphoid tissue lymphoma (Table 1). The *L265P* mutation affects the MYD88 TIR domain, which is responsible for recruiting the protein to the cytoplasmic tail of TLRs to form an active complex, which subsequently activates the kinases IRAK1 and IRAK4 to signal downstream.⁴⁵ A hyperphosphorylated slow migrating isoform of IRAK1 associates strongly with the *L265P* mutant MYD88 but not wild type MYD88 suggesting a gain-of-function activity in the *L265P* mutant.²⁴ The *MYD88 L265P* mutation was also found to be a potent driver of high NF κ B activity, which is characteristic of ABC DLBCL.²⁴

Interestingly, in addition to enhancing NF κ B signaling, MYD88 L265P seems to increase JAK-STAT signaling and interferon production, indicating the potential involvement of a niche microenvironment important for tumor survival.²⁴ STAT3 signaling induced by cytokines such as IL6 could provide additional survival signals sustaining lymphoma survival, given that transgenic mice expressing supra-physiological amounts of IL6 develop a range of lymphoid malignancies, including DLBCL.⁴⁶ Thus, IL6 and IL10 production potentially form an important autocrine feedback loop that activates JAK-STAT signaling to enhance the survival of ABC DLBCL.^{24,47} Interestingly, overexpression of the L265P MYD88 variant has recently been associated with reduced disease free survival and increased disease recurrence in DLBCL patients.⁴⁸ These recent developments highlight MYD88 as a specific target for therapeutic intervention, and warrants the development of inhibitors targeting the MYD88-NF κ B signaling axis.

Targeting oncogenic MYD88

Aberrant NF κ B activation has been associated with poor clinical outcomes in many lymphomas and leukemia. Thus, effective therapeutic agents targeting the NF κ B pathway may allow the achievement of desirable outcome for a subset of patients. Given the large proportion of lymphoid neoplasms with aberrant TLR signaling, targeting the MYD88 pathway is becoming an attractive option for clinicians and researchers.

Direct inhibition of MYD88

A critical event in MYD88 mediated signaling is the homodimerization of MYD88 through its TIR and DD domains.⁴⁹ The dimerization of MYD88 promotes its recruitment to the plasma membrane and docking with the TIR domain of the cytoplasmic tails of TLRs or IL1R, as well as the recruitment of IRAK4 and IRAK1 through the interaction between their DD domains.^{43,50} Given signal transduction through MYD88 requires its homodimerization and lymphoma associated *MYD88* mutations occur exclusively in the TIR domain, one appealing option would be to 'switch off' this signaling relay event so that MYD88 homodimerization and downstream signaling is inhibited.

TIR-TIR interaction in MYD88 is achieved by distinct conserved residues in a structure known as the BB-loop that lies between the second β -strand and the second α -helix.⁵¹ The interference of this interaction was successfully achieved by the use of heptapeptides mimicking BB-loop by Sette and colleagues.⁵¹ When this dimerization is inhibited, significant reduction in NF κ B activity is achieved in cells stimulated with IL1 or TLR agonists but not poly(I:C), a TLR3 agonist, suggesting this component selectively inhibits MYD88 dependent signaling.⁵¹ The same group also identified a novel synthetic compound, ST2825, which mimics the heptapeptide in the BB-loop of MYD88 and this compound is currently undergoing preclinical evaluations.⁵²

Alternative options to specifically target MYD88-MYD88 and MYD88-receptor interactions through the TIR domain would be to use small molecule inhibitors such as Hydrocinnamoyl-L-valyl pyrrolidine (compound 4a).⁵³ This particular low molecular weight compound is cell membrane permeable and specifically disrupts MYD88-receptor interactions by inhibiting TIR domain interactions. Another peptide Pepinh-MYD developed by InvivoGen, which carries a 26 amino acid MYD88 homodimerization motif, could also potentially be used to treat lymphoma patients with L265P MYD88. However, these potential MYD88-specific therapeutic options are yet to be trialed in large clinical cohorts.

Targeting IRAK4, downstream of receptor-MYD88 signaling

Interleukin receptor-associated kinases (IRAKs) are a key component of the signal transduction pathways downstream of IL1 receptor or TLRs, and are required for the activation of NF κ B and MAPKs in response to the activation of these receptors.³⁴ In particular, IRAK4 serves as the “master IRAK” by having the ability to phosphorylate and activate other IRAKs. The *MYD88 L265P* mutation result in a constitutively activated signaling complex, which includes IRAK4 and phosphorylated IRAK1.²⁴ Since ABC DLBCL lines depend on kinase activity of IRAK4, but not IRAK1, inhibiting IRAK4 kinase activity could be a potent way of ‘tuning down’ aberrant NF κ B and MAPK pathways activated by *MYD88* mutations.

The activation of IRAK4 is regulated by the autophosphorylation of three serine and threonine sites in its activation loop.⁵⁴ The autophosphorylation of these key activating residues confer a conformation change to allow the ATP-binding site to become activated.⁵⁴ The kinase activity of IRAK4 is thus often targeted by small molecular inhibitors that bind to the ATP-binding site. In ABC DLBCL cell lines carrying the *MYD88 L265P* mutation, disrupting IRAK4 signaling by a small molecule inhibitor led to the decreased phosphorylation of IRAK1, I κ B α , NF κ B p65 and STAT3 (unpublished data). Two small molecule IRAK4 inhibitors (ND-2110 and ND-2158) developed by Nimbus Discovery are highly selective against more than 300 kinases and seem promising, although these drugs are still in the preclinical stage.

Targeting the two ends-inhibiting TLRs and the NF κ B signaling

It has long been speculated that cognate antigen stimulation might contribute to lymphomagenesis.⁵⁵ Since *MYD88* physiologically serves as an adaptor molecule for TLR sensing pathogens, it is an attractive idea that lymphomas with *MYD88* mutations need external or internal signals from TLRs for their survival. However, it remains unclear whether oncogenic mutant *MYD88* proteins require upstream signaling from ligand-activated receptors for enhancing its activity. But, in the potential scenario where ligand-TLR engagement would be required for oncogenic *MYD88* activation, inhibiting TLR signaling upstream of *MYD88* can be a potential therapeutic target for treating B cell malignancies carrying *MYD88* mutations. TLR antagonists which are currently under preclinical and clinical evaluation could be potentially used to inhibit receptor signaling.⁵⁶

An alternative point at which signaling from oncogenic *MYD88* could be interrupted would be through direct inhibition of NF κ B activity. In recent years, more than 800 drugs that inhibit NF κ B signaling have been developed, and their mechanisms of action have been characterized.^{57,58} For instance drugs such as emetine, bithionol, narasin and lestaurtinib inhibit NF κ B signaling via inhibition of I κ B α phosphorylation, a critical step required for activation of NF κ B, while drugs such as bortezomib and Carfilzomib inhibit NF κ B activity by dampening proteosomal degradation of I κ B α .⁵⁹ Even though a number of NF κ B inhibitors are FDA-approved for use in particular cancer types, inhibition of this particular family of transcription factors would be accompanied by a number of undesired side-effects. NF κ B is known to have a critical role across many cellular processes including cell proliferation, apoptosis, immune responses to infection, and inflammation, such that the beneficial effects and potential collateral damage must be carefully examined.⁶⁰

Combination therapy

Lenalidomide (Revlimid®, Celgene Corporation, Summit, NJ, USA), a derivative of thalidomide, is an immunomodulatory drug and currently used as a treatment of multiple myeloma and myelodysplastic syndromes.^{61,62} Although the mechanism of action still remains unclear, clinical trials showed lenalidomide is effective against most lymphomas. ABC DLBCL had a significantly higher response rate to the thalidomide analog compared to GC DLBCL, indicating that one of its actions is a suppressive effect on the NF κ B pathway.⁶³

A recent study from Staudt et al found synergistic effects between lenalidomide and a BTK inhibitor, ibrutinib.⁶⁴ They found that lenalidomide was partially toxic to ABC DLBCL lines by inhibiting NF κ B, JAK and *MYD88* signaling as well as augmenting IFN beta signature.⁶⁴ IFN beta production and upregulation of IFN beta-responsive genes is characteristic of ABC DLBCL harboring *MY88* mutations, although its pathological roles are still unclear since IFN beta is known to induce cell cycle arrest and apoptosis paradoxically.⁶⁵ IFN beta production is induced by IRF7 in a positive-feedback manner and IRF4/SPIB transcription factor network represses IFN beta production by downregulating IRF7.⁶⁶ Ibrutinib cooperates with lenalidomide to kill ABC DLBCL lines, presumably due to inducing IFN beta pathway by changing the balance between IRF7 and IRF4/SPIB.⁶⁴

Another potential strategy of combination therapy would be to block IRAK4 signaling and BTK signaling in synergy. This strategy is consistent with the observation that survival of ABC DLBCL cell lines require a signal through both the

TLR and the BCR as knocking down CD79 molecule of the BCR signaling compartment synergistically kills ABL-DLBCL cell lines with MYD88 knockdown.²⁴ Therefore synergistic killing of lymphoid neoplasms with L265P MYD88 using a combination of IRAK inhibitor and BTK inhibitor strongly indicate that most potent therapy should target both the TLR and BCR pathways simultaneously.

Concluding remarks

Recent discoveries have uncovered the involvement of L265P MYD88 in a number of diseases, but the effects of this oncogenic variant remain unclear. Future research would help elucidate the effects of deregulated MYD88 on cellular signaling pathways, the activity of transcription factors, and gene expression changes. However, from the initial biochemical characterization of oncogenic MYD88, it is apparent that several key signaling pathways are disrupted such that effective therapeutic strategies would involve multi-pronged approaches with combination of specific agents targeting biochemically distinct pathways simultaneously. Such therapeutic regimes remain to be evaluated in large cohorts of patients, and a factor central to evaluation of the effectiveness of such treatment regimes would be the appropriate selection of patients. Recently, a number of diagnostic tools have been developed to specifically detect the presence of the L265P MYD88 variant in clinical samples.^{67–69} With the refinement of these diagnostic methods and the development of targeted therapeutics, clinicians would, in the near future, be in a better position to provide treatment options to previously incurable diseases.

Disclosure

The authors have no conflict of interest to declare.

References

- Hodgkin T. On some Morbid Appearances of the Absorbent Glands and Spleen. *Med Chir Trans*. 1832;17:68–114.
- Jaffe ES, Pittaluga S. Aggressive B-cell lymphomas: a review of new and old entities in the WHO classification. *Hematology Am Soc Hematol Educ Program*. 2011;2011:506–514.
- Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646–674.
- Swerdlow S, Campo E, Harris N, et al. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*, 4 ed. Lyon, France: IARC Press; 2008.
- Bray F, Ren JS, Masuyer E, Ferlay J. Global estimates of cancer prevalence for 27 sites in the adult population in 2008. *Int J Cancer*. 2013;132(5):1133–1145.
- Siegel R, Ward E, Brawley O, Jemal A. Cancer statistics, 2011: the impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin*. 2011;61(4):212–236.
- Jung D, Giallourakis C, Mostoslavsky R, Alt FW. Mechanism and control of V(D)J recombination at the immunoglobulin heavy chain locus. *Annu Rev Immunol*. 2006;24:541–570.
- Victoria GD, Nussenzweig MC. Germinal centers. *Annu Rev Immunol*. 2012;30:429–457.
- Muramatsu M, Kinoshita K, Fagarasan S, Yamada S, Shinkai Y, Honjo T. Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell*. 2000;102(5):553–563.
- Liu M, Duke JL, Richter DJ, et al. Two levels of protection for the B cell genome during somatic hypermutation. *Nature*. 2008;451(7180):841–845.
- Chiarle R, Zhang Y, Frock RL, et al. Genome-wide translocation sequencing reveals mechanisms of chromosome breaks and rearrangements in B cells. *Cell*. 2011;147(1):107–119.
- Klein IA, Resch W, Jankovic M, et al. Translocation-capture sequencing reveals the extent and nature of chromosomal rearrangements in B lymphocytes. *Cell*. 2011;147(1):95–106.
- Shaffer AL 3rd, Young RM, Staudt LM. Pathogenesis of human B cell lymphomas. *Annu Rev Immunol*. 2012;30:565–610.
- Rosenwald A, Wright G, Chan WC, et al. The use of molecular profiling to predict survival after chemotherapy for diffuse large-B-cell lymphoma. *N Engl J Med*. 2002;346(25):1937–1947.
- Alizadeh AA, Eisen MB, Davis RE, et al. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature*. 2000;403(6769):503–511.
- Ngo VN, Davis RE, Lamy L, et al. A loss-of-function RNA interference screen for molecular targets in cancer. *Nature*. 2006;441(7089):106–110.
- Baldwin AS. Control of oncogenesis and cancer therapy resistance by the transcription factor NF-kappaB. *J Clin Invest*. 2001;107(3):241–246.
- Lenz G, Davis RE, Ngo VN, et al. Oncogenic CARD11 mutations in human diffuse large B cell lymphoma. *Science*. 2008;319(5870):1676–1679.
- Davis RE, Ngo VN, Lenz G, et al. Chronic active B-cell-receptor signalling in diffuse large B-cell lymphoma. *Nature*. 2010;463(7277):88–92.
- Compagno M, Lim WK, Grunn A, et al. Mutations of multiple genes cause deregulation of NF-kappaB in diffuse large B-cell lymphoma. *Nature*. 2009;459(7247):717–721.
- Kato M, Sanada M, Kato I, et al. Frequent inactivation of A20 in B-cell lymphomas. *Nature*. 2009;459(7247):712–716.
- Honma K, Tsuzuki S, Nakagawa M, et al. TNFAIP3/A20 functions as a novel tumor suppressor gene in several subtypes of non-Hodgkin lymphomas. *Blood*. 2009;114(12):2467–2475.
- Wertz IE, O'Rourke KM, Zhou H, et al. De-ubiquitination and ubiquitin ligase domains of A20 downregulate NF-kappaB signalling. *Nature*. 2004;430(7000):694–699.
- Ngo VN, Young RM, Schmitz R, et al. Oncogenically active MYD88 mutations in human lymphoma. *Nature*. 2011;470(7332):115–119.
- Jeelall YS, Horikawa K. Oncogenic MYD88 mutation drives Toll pathway to lymphoma. *Immunology and Cell Biology*. 2011;89(6):659–660.
- Treon SP, Xu L, Yang G, et al. MYD88 L265P somatic mutation in Waldenstrom's macroglobulinemia. *N Engl J Med*. 2012;367(9):826–833.
- Lord KA, Hoffman-Liebermann B, Liebermann DA. Nucleotide sequence and expression of a cDNA encoding MyD88, a novel myeloid differentiation primary response gene induced by IL6. *Oncogene*. 1990;5(7):1095–1097.
- von Bernuth H, Picard C, Jin Z, et al. Pyogenic bacterial infections in humans with MyD88 deficiency. *Science*. 2008;321(5889):691–696.
- Picard C, Casanova JL, Puel A. Infectious diseases in patients with IRAK-4, MyD88, NEMO, or IkappaBalpha deficiency. *Clin Microbiol Rev*. 2011;24(3):490–497.
- Adachi O, Kawai T, Takeda K, et al. Targeted disruption of the MyD88 gene results in loss of IL-1- and IL-18-mediated function. *Immunity*. 1998;9(1):143–150.
- Hashimoto C, Hudson KL, Anderson KV. The Toll gene of *Drosophila*, required for dorsal-ventral embryonic polarity, appears to encode a transmembrane protein. *Cell*. 1988;52(2):269–279.

32. Belvin MP, Anderson KV. A conserved signaling pathway: the Drosophila toll-dorsal pathway. *Annu Rev Cell Dev Biol.* 1996;12:393–416.
33. Kopp EB, Medzhitov R. The Toll-receptor family and control of innate immunity. *Curr Opin Immunol.* 1999;11(1):13–18.
34. Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol.* 2004;4(7):499–511.
35. Muzio M, Bosio D, Polentarutti N, et al. Differential expression and regulation of toll-like receptors (TLR) in human leukocytes: selective expression of TLR3 in dendritic cells. *J Immunol.* 2000;164(11):5998–6004.
36. Ulevitch RJ, Tobias PS. Receptor-dependent mechanisms of cell stimulation by bacterial endotoxin. *Annu Rev Immunol.* 1995;13:437–457.
37. Kawai T, Adachi O, Ogawa T, Takeda K, Akira S. Unresponsiveness of MyD88-deficient mice to endotoxin. *Immunity.* 1999;11(1):115–122.
38. Morrison DC, Ryan JL. Endotoxins and disease mechanisms. *Annu Rev Med.* 1987;38:417–432.
39. Krieg AM. The role of CpG motifs in innate immunity. *Curr Opin Immunol.* 2000;12(1):35–43.
40. Yamamoto S, Yamamoto T, Tokunaga T. The discovery of immunostimulatory DNA sequence. *Springer Semin Immunopathol.* 2000;22(1–2):11–19.
41. Schnare M, Holt AC, Takeda K, Akira S, Medzhitov R. Recognition of CpG DNA is mediated by signaling pathways dependent on the adaptor protein MyD88. *Curr Biol.* 2000;10(18):1139–1142.
42. Wesche H, Henzel WJ, Shillinglaw W, Li S, Cao Z. MyD88: an adapter that recruits IRAK to the IL-1 receptor complex. *Immunity.* 1997;7(6):837–847.
43. Akira S. Toll-like receptor signaling. *J Biol Chem.* 2003;278(40):38105–38108.
44. Dunne A, Ejdeback M, Ludidi PL, O'Neill LA, Gay NJ. Structural complementarity of Toll/interleukin-1 receptor domains in Toll-like receptors and the adaptors Mal and MyD88. *J Biol Chem.* 2003;278(42):41443–41451.
45. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol.* 2010;11(5):373–384.
46. Kishimoto T. IL-6: from its discovery to clinical applications. *Int Immunol.* 2010;22(5):347–352.
47. Ding BB, Yu JJ, Yu RY, et al. Constitutively activated STAT3 promotes cell proliferation and survival in the activated B-cell subtype of diffuse large B-cell lymphomas. *Blood.* 2008;111(3):1515–1523.
48. Choi JW, Kim Y, Lee JH, Kim YS. MYD88 expression and L265P mutation in diffuse large B-cell lymphoma. *Hum Pathol.* In press 2013.
49. Burns K, Martinon F, Esslinger C, et al. MyD88, an adapter protein involved in interleukin-1 signaling. *J Biol Chem.* 1998;273(20):12203–12209.
50. Janssens S, Beyaert R. Functional diversity and regulation of different interleukin-1 receptor-associated kinase (IRAK) family members. *Mol Cell.* 2003;11(2):293–302.
51. Loiarro M, Sette C, Gallo G, et al. Peptide-mediated interference of TIR domain dimerization in MyD88 inhibits interleukin-1-dependent activation of NF- κ B. *J Biol Chem.* 2005;280(16):15809–15814.
52. Loiarro M, Capolunghi F, Fanto N, et al. Pivotal Advance: Inhibition of MyD88 dimerization and recruitment of IRAK1 and IRAK4 by a novel peptidomimetic compound. *J Leukoc Biol.* 2007;82(4):801–810.
53. Bartfai T, Behrens MM, Gaidarova S, Pemberton J, Shivanyuk A, Rebek J Jr. A low molecular weight mimic of the Toll/IL-1 receptor/resistance domain inhibits IL-1 receptor-mediated responses. *Proc Natl Acad Sci U S A.* 2003;100(13):7971–7976.
54. Cheng H, Addona T, Keshishian H, et al. Regulation of IRAK-4 kinase activity via autophosphorylation within its activation loop. *Biochem Biophys Res Commun.* 2007;352(3):609–616.
55. Kuppers R. Mechanisms of B-cell lymphoma pathogenesis. *Nat Rev Cancer.* 2005;5(4):251–262.
56. Hennessy EJ, Parker AE, O'Neill LA. Targeting Toll-like receptors: emerging therapeutics? *Nat Rev Drug Discov.* 2010;9(4):293–307.
57. Gilmore TD, Herscovitch M. Inhibitors of NF-kappaB signaling: 785 and counting. *Oncogene.* 2006;25(51):6887–6899.
58. Miller SC, Huang R, Sakamuru S, et al. Identification of known drugs that act as inhibitors of NF-kappaB signaling and their mechanism of action. *Biochem Pharmacol.* 2010;79(9):1272–1280.
59. Huang R, Southall N, Wang Y, et al. The NCGC pharmaceutical collection: a comprehensive resource of clinically approved drugs enabling repurposing and chemical genomics. *Sci Transl Med.* 2011;3(80):80ps16.
60. Vallabhapurapu S, Karin M. Regulation and function of NF-kappaB transcription factors in the immune system. *Annu Rev Immunol.* 2009;27:693–733.
61. Reeder CB, Ansell SM. Novel therapeutic agents for B-cell lymphoma: developing rational combinations. *Blood.* 2011;117(5):1453–1462.
62. Younes A. Beyond chemotherapy: new agents for targeted treatment of lymphoma. *Nat Rev Clin Oncol.* 2011;8(2):85–96.
63. Hernandez-Ilizaliturri FJ, Deeb G, Zinzani PL, et al. Higher response to lenalidomide in relapsed/refractory diffuse large B-cell lymphoma in nongerminal center B-cell-like than in germinal center B-cell-like phenotype. *Cancer.* 2011;117(22):5058–5066.
64. Yang Y, Shaffer AL 3rd, Emre NC, et al. Exploiting synthetic lethality for the therapy of ABC diffuse large B cell lymphoma. *Cancer Cell.* 2012;21(6):723–737.
65. Stark GR, Kerr IM, Williams BR, Silverman RH, Schreiber RD. How cells respond to interferons. *Annu Rev Biochem.* 1998;67:227–264.
66. Honda K, Yanai H, Negishi H, et al. IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature.* 2005;434(7034):772–777.
67. Xu L, Hunter ZR, Yang G, et al. MYD88 L265P in Waldenstrom's Macroglobulinemia, IgM Monoclonal Gammopathy, and other B-cell Lymphoproliferative Disorders using Conventional and Quantitative Allele-Specific PCR. *Blood.* 2013;121(11):2051–2058.
68. Varettoni M, Arcaini L, Zibellini S, et al. Prevalence and clinical significance of the MYD88 (L265P) somatic mutation in Waldenstrom's macroglobulinemia and related lymphoid neoplasms. *Blood.* 2013;121(13):2522–2528.
69. Gachard N, Parrens M, Soubeyran I, et al. IGHV gene features and MYD88 L265P mutation separate the three marginal zone lymphoma entities and Waldenstrom macroglobulinemia/lymphoplasmacytic lymphomas. *Leukemia.* 2013;27(1):183–189.
70. Pham-Ledard A, Cappellen D, Martinez F, Vergier B, Beylot-Barry M, Merlio JP. MYD88 somatic mutation is a genetic feature of primary cutaneous diffuse large B-cell lymphoma, leg type. *J Invest Dermatol.* 2012;132(8):2118–2120.
71. Puente XS, Pinyol M, Quesada V, et al. Whole-genome sequencing identifies recurrent mutations in chronic lymphocytic leukaemia. *Nature.* 2011;475(7354):101–105.
72. Wang L, Lawrence MS, Wan Y, et al. SF3B1 and other novel cancer genes in chronic lymphocytic leukemia. *N Engl J Med.* 2011;365(26):2497–2506.
73. Treon SP, Xu L, Yang G, et al. MYD88 L265P somatic mutation in Waldenstrom's macroglobulinemia. *N Engl J Med.* 2012;367(9):826–833.
74. Li ZM, Rinaldi A, Cavalli A, et al. MYD88 somatic mutations in MALT lymphomas. *Br J Haematol.* 2012;158(5):662–664.
75. Gachard N, Parrens M, Soubeyran I, et al. IGHV gene features and MYD88 L265P mutation separate the three marginal zone lymphoma entities and Waldenstrom macroglobulinemia/lymphoplasmacytic lymphomas. *Leukemia.* Jan 2013;27(1):183–189.
76. Montesinos-Rongen M, Godlewska E, Brunn A, Wiestler OD, Siebert R, Deckert M. Activating L265P mutations of the MYD88 gene are common in primary central nervous system lymphoma. *Acta Neuropathol.* Dec 2011;122(6):791–792.
77. Gonzalez-Aguilar A, Idhah A, Boisselier B, et al. Recurrent mutations of MYD88 and TBL1XR1 in primary central nervous system lymphomas. *Clin Cancer Res.* Oct 1, 2012;18(19):5203–5211.

78. Yan Q, Huang Y, Watkins AJ, et al. BCR and TLR signaling pathways are recurrently targeted by genetic changes in splenic marginal zone lymphomas. *Haematologica*. Apr 2012;97(4): 595–598.
79. Rossi D, Trifonov V, Fangazio M, et al. The coding genome of splenic marginal zone lymphoma: activation of NOTCH2 and other pathways regulating marginal zone development. *J Exp Med*. Aug 27, 2012;209(9):1537–1551.

Blood and Lymphatic Cancer: Targets and Therapy

Dovepress

Publish your work in this journal

Blood and Lymphatic Cancer: Targets and Therapy is an international, peer-reviewed, open access journal focusing on blood and lymphatic cancer research, identification of therapeutic targets and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the

cancer patient. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/blood-and-lymphatic-cancer-targets-and-therapy-journal>