Monitoring human papillomavirus prevalence in urine samples: a review [Corrigendum]

Espen Enerly, Cecilia Olofsson, Mari Nygård. *Clin Epidemiol.* 2013;5:67–79.

On page 72, the following sentence was open to misinterpretation: “The most common urine sampling procedure in both female and male populations was first-void urine in contrast to mid-stream urine, which Vorsters et al suggested would be preferable if the goal were to analyse a maximum number of exfoliated cells”. The correct sentence is “The most common urine sampling procedure in both female and male populations was first-void urine, which Vorsters et al suggested would be preferable to mid-stream urine if the goal were to analyse a maximum number of exfoliated cells”.

Clinical Epidemiology

Dovepress

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification, systematic

reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Submit your manuscript here: <http://www.dovepress.com/clinical-epidemiology-journal>