Inhibition of choriocarcinoma by Fe_3O_4 -dextran-anti- β -human chorionic gonadotropin nanoparticles containing antisense oligodeoxynucleotide of heparanase

Liu Huining¹
Zhang Yi¹
Tang Dihong²
Pan Yifeng³
Xia Man²
Yang Ting²
Cai Jingting^{1,2}

¹Department of Obstetrics and Gynecology, Xiangya Hospital,

²Department of Gynecological Oncology, Hunan Provincial Tumor Hospital, ³National Hepatobiliary and Enteric Surgery Research Center, Central South University, Changsha, Hunan, People's Republic of China

Objective: To observe the influence of Fe_3O_4 -dextran-anti- β -human chorionic gonadotropin (HCG) carrying heparanase (Hpa) antisense oligodeoxynucleotide (ASODN), via the invasion, proliferation, and Hpa expression of JEG-3 cell lines and inhibitory effect of transplanted choriocarcinoma tumor growth.

Methods: The different abilities of invasion and proliferation between transfected JEG-3 and untransfected JEG-3 were measured by Matrigel invasion assay and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay in vitro. The effect of Hpa ASODN transfection on the expression of Hpa mRNA and protein was measured by reverse-transcription polymerase chain reaction and Western blot. The transplanted choriocarcinoma tumors were taken out to calculate the inhibitory effect on tumor growth of Hpa ASODN.

Results: In this study, we found that: (1) the invasive ability of JEG-3 cells was inhibited sufficiently ($P < 0.05$) after JEG-3 cells were transfected by Fe_3O_4 -dextran-anti- β HCG carrying Hpa ASODN; (2) after JEG-3 cells were transfected by Fe_3O_4 -dextran-anti- β HCG carrying Hpa ASODN at 48 and 72 hours, the proliferative ability of JEG-3 cells was inhibited sufficiently ($P < 0.05$); (3) the expression of Hpa mRNA and protein in JEG-3 cells was inhibited efficiently after JEG-3 cells were transfected by Fe_3O_4 -dextran-anti- β HCG carrying Hpa ASODN ($P < 0.05$); and (4) Fe_3O_4 -dextran-anti- β HCG carrying Hpa ASODN had an inhibitory effect on the transplanted choriocarcinoma tumor growth ($P < 0.05$) and was harmless on nude mice.

Conclusion: Fe_3O_4 -dextran-anti- β HCG carrying Hpa ASODN weakened the invasive and proliferative ability of choriocarcinoma, with a significant inhibitory effect on the transplanted choriocarcinoma tumor. Therefore, Fe_3O_4 -dextran-anti- β HCG carrying Hpa ASODN is an effective gene therapy, and Fe_3O_4 -dextran-anti- β HCG nanoparticles are a harmless and effective gene vector.

Keywords: heparanase, antisense oligodeoxynucleotide, Fe_3O_4 -dextran-anti- β HCG nanoparticles, choriocarcinoma, invasive ability

Introduction

Heparanase (Hpa) is a mammalian endo- β -D-glucuronidase responsible for heparan sulfate degradation.¹ Hpa activity may therefore play an important role in fundamental biological processes associated with extracellular matrix remodeling and cell invasion.² In our study, we have found out that the expression of Hpa plays an important role in the metastatic capability of human choriocarcinoma.³ Therefore, we deem that Hpa is the therapeutic target of choriocarcinoma and try to prove this hypothesis.

Correspondence: Cai Jingting
Department of Gynecological Oncology,
Hunan Provincial Tumor Hospital,
Changsha, Hunan 410013, People's
Republic of China
Tel +86 731 8976 2081
Fax +86 731 8887 6554
Email cjingting114@yahoo.com.cn

Gene therapy is the insertion of genes into an individual's cells and tissues to treat a disease, and hereditary diseases in particular. Gene therapy typically aims to supplement a defective mutant allele with a functional one. Although the technology is still in its infancy, it has been used with some success. In antisense therapy, an antisense copy of the messenger RNA (mRNA) of the target gene is used to inhibit translation and production of the protein. In gene therapy, a carrier (vector) must be used to deliver the therapeutic gene to the patient's target cells. Ideally, a gene-delivery system should be stable, biocompatible, nontoxic, cost-effective, and capable of transferring exogenous highly anionic genetic materials (ie, DNA, antisense oligonucleotides [ASODNs], short interfering RNA) into tissue-specific sites. A major goal of gene therapy is to obtain targeted vectors that transfer genes efficiently to specific cell types. The appearance of nanoparticles provides a new idea for gene vector. As a gene vector, nanoparticles possess several advantages⁴: (1) nanoparticles are not biologic materials, so they will not induce immune responses to human beings; (2) unlike viral carriers, nanoparticles have no genotoxicity or cytotoxicity; (3) nanoparticles are an effective gene vector; (4) nanoparticles can insert foreign genes in DNA in chromosomes of host cells, and as a result, foreign genes are expressed in host cells stably; (5) nanoparticles can defend foreign genes from complements, enzymes, etc; and (6) nanoparticles can counteract or kill some viruses. Having so many advantages, nanoparticles are potentially the most promising gene vector.

We have found out previously that Fe_3O_4 -dextran-anti- β -human chorionic gonadotropin (HCG) nanoparticles represent a versatile vector for gene therapy with high stability towards degradation.⁵ They combine a high introduction potential with good biocompatibility and low cytotoxicity. With magnetic and immune targeting, they offer the possibility of additional targeting strategies. In this study, we synthesized magnetic nanoparticles, and using those as a gene vector, we transfected Hpa ASODNs into choriocarcinoma JEG-3 cells. Afterwards, we estimated the curative effect of Hpa ASODNs on choriocarcinoma. We propose that the antitumor effect of Fe_3O_4 -dextran-anti- β HCG nanoparticles carrying Hpa ASODN has been demonstrated as proof of principle.

Material and methods

Cells and animals

Cell-culture studies

JEG-3 and JAR are human choriocarcinoma cell lines, and were bought from the American Type Culture Collection

(ATCC, Manassas, VA, USA). Both cell lines were cultured in Roswell Park Memorial Institute (RPMI) medium (Life Technologies, Carlsbad, CA, USA) supplemented with 10% fetal calf serum (FCS) in a humidified atmosphere of 5% CO_2 and 37°C. Culture medium was replaced every other day.

Animals

Female BALB/c nude mice, 4 weeks old, were supplied and fed by the animal division of Central South University, Hunan, People's Republic of China.

Preparation of magnetic nanoparticles of Fe_3O_4 -dextran-anti- β HCG

Materials

Dextran with a molecular weight of 40,000 (T-40) and bovine serum albumin (BSA) were bought from Sigma-Aldrich, St Louis, MO, USA. Mouse anti-human β HCG monoclonal antibody was purchased from Wolvo Biotechnology, Shanghai, People's Republic of China. $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$, and other chemicals were from local suppliers and were of analytical grade. Hpa ASODNs and non-sense oligonucleotides (NSODNs) were synthesized by Shanghai Sangon Biological Engineering Technology and Services (Shanghai, People's Republic of China). Hpa ASODNs' sequence was 5'-GGCTTCGAGCGCAGCAG CAT-3', and Hpa NSODNs' sequence was 5'-TCAGCTAGCGAGGCTGCGC A-3'. After being diluted to 100 $\mu\text{mol/L}$, all samples were preserved in -20°C . Mixtures of Fe_3O_4 -dextran-anti- β HCG-Hpa ASODN or NSODN Fe_3O_4 -dextran-anti- β HCG nanoparticles were prepared in a manner similar to Jingting et al.⁴

Preparation of mixtures of Fe_3O_4 -dextran-anti- β HCG-Hpa ASODNs or NSODNs

$\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (0.552 g), $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ (0.45 g), and dextran (5 g) were dispersed in 10 mL H_2O . While under magnetism stirring with a velocity of 800 rpm, the mixture was titrated to alkaline by the dropwise addition of 10 mL 5 mol/L ammonia water and quickly heated to 60°C. After 30 minutes of reaction, the resulting black suspension was neutralized with acetic acid. Aggregates were removed by centrifugation for 10 minutes at 8,000 rpm. The dextran-coated iron oxide nanoparticles were separated from unbound dextran by magnetic separator and eluted with 10 mL sodium acetate (pH 6.5). The dextran-coated iron oxide suspension was oxidized with sodium periodate (NaIO_4) under magnetism stirring in the dark at 4°C for 1 hour to obtain aldehyde termination, and the solution was dialyzed by magnetic separator again.

Afterwards, 100 µg mouse anti-human βHCG monoclonal antibody and 200 µL 100 µmol/L Hpa ASODNs or NSODNs were put into 1 mL aldehyde dextran-coated iron oxide nanoparticles and mixed together fully. After being reacted in the dark at 4°C for 24 hours, the mixture was blocked by 50 µL 10 wt% BSA. For improved stability, 1 mol/L NaBH₄ was added to the compound at 4°C for 3 hours and stirred. Lastly, the suspension was centrifuged for 20 minutes at 13,000 rpm three times. The production was dispersed again by 100 µL NaBO₃ (pH 8.5).

Introduction of Hpa ASODN or NSODN to JEG-3 cells mediated by Fe₃O₄-dextran-anti-βHCG nanoparticles

When JEG-3 cells had grown to 70%–80%, the cells were washed with free Dulbecco's Modified Eagle's Medium (DMEM) (Sigma-Aldrich), and then 4.5 mL DMEM that contained 10% FCS and 500 µL Fe₃O₄-dextran-anti-βHCG-Hpa ASODN or NSODN compounds, which contained 20 µmol/L Hpa ASODN or NSODN in every culture bottle (37.5 µg/mL Fe³⁺ and 100 µg/mL mouse anti-human βHCG monoclonal antibody in introduction reagent). After the introduction (24 hours), the cells were used for later experimentation.

MTT assay

Transfected and untransfected JEG-3 cells were seeded in 96-well plates (5,000 cells/well) in DMEM containing 10% FCS. After 24, 48, and 72 hours, MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide; Sigma-Aldrich) was added to the culture medium at a concentration of 5 mg/mL for 4 hours at 37°C. Subsequently, the cells were washed with phosphate-buffered saline, and the purple formazan was solubilized in 100% dimethyl sulfoxide. Absorbance was read at 570 nm. Five separate sets of experiments were performed and averaged.

Matrigel invasion assay

A Matrigel invasion assay was prepared as described in Staff et al.⁶ Briefly, diluted 1:2 Matrigel (1.75 µg/µL; BD Biosciences, Beit Ha'Emek, Israel) in serum-free cell-culture media was added to the upper chamber of a 24-well transwell plate and incubated at 37°C for 3–4 hours for gelling. Transfected and untransfected JEG-3 cells were harvested from tissue-culture flasks by trypsin/ethylene-diaminetetraacetic acid, washed and resuspended in 0.1% FCS in DMEM, and added to upper wells at a density of 10⁵ cells/well in 100 µL medium, while 600 µL medium

was added to the lower wells. Plates were incubated at 37°C for 24 hours, and then the cells remaining on the upper surface of the membrane were removed with a cotton swab and the filters fixed by 95% ethanol for 30 minutes. Cells that had invaded the lower surface of the filter were counted under an inverted microscope; ten fields per well were counted. All experiments were performed five times, and the results from the five separate sets of experiments were averaged.

Reverse-transcription polymerase chain reaction

RNA extraction

The commercially available Trizol reagent (Life Technologies, Carlsbad, CA, USA) was used to extract total RNA from cells. In brief, cells were milled and lysed in 1 mL of Trizol and left at room temperature for 5–10 minutes. Subsequently, 0.2 mL of chloroform was added, followed by vigorous agitation for 15 seconds. The sample mixture was kept at room temperature for 2–3 minutes, and then centrifuged at 12,000 g for 15 minutes at 4°C. The aqueous phase was incubated with 0.5 mL of isopropanol for 10 minutes at room temperature, and then centrifuged at 12,000 g for 10 minutes at 4°C. The precipitated RNA was washed with 75% ethanol and dissolved in 0.1% diethylpyrocarbonate-treated water. RNA concentrations were measured spectrophotometrically (1 OD₂₆₀ 40 mg/mL of RNA).

Reverse-transcription polymerase chain reaction

The expression levels of Hpa and glyceraldehyde-3-phosphodehydrogenase (*GAPDH*) genes were determined by reverse-transcription polymerase chain reaction (RT-PCR). Total RNA was reverse-transcribed in 30 mL of a solution containing 1 µg of total RNA random hexamer primers, 5× reaction buffer, RiboLock RNase inhibitor, 10 mM deoxy-ribonucleotide mix, Revert Aid Moloney murine leukemia virus reverse transcriptase, and diethylpyrocarbonate-treated water (Thermo Fisher Scientific, Waltham, MA, USA). PCR was performed to detect the expression of Hpa mRNA in transfected and untransfected JEG-3 cells. Based on published DNA sequences of human *HPA* and *GAPDH* genes, primers for Hpa¹ (sense, HPU-355 5'-TTCGATC-CCAAG AAGGAATCAAC-3', antisense, HPL-229 5'-GTAGTGATGCCATGTAAGTGAATC-3') and *GAPDH* (sense, 5'-GCTGGCGCTGAGTACGTCGT-3', antisense, 5'-TGGGTGTCGCTGTTGAAGTC-3') were obtained. PCR was performed using an MJ Research (Bio-Rad, Waltham, MA, USA) PCR system. The following conditions

were applied to Hpa PCR amplifications: 94°C for 5 minutes, 38 cycles (denaturation at 94°C for 45 seconds, annealing at 55°C for 45 seconds, extension at 72°C for 1 minute), and 72°C for 10 minutes. The conditions for GAPDH PCR amplifications were 94°C for 5 minutes, 35 cycles (denaturation at 94°C for 45 seconds, annealing at 55°C for 50 seconds, extension at 72°C for 1 minute), and 72°C for 10 minutes. Both reactions were in 25 mL mixtures containing 5 µL of the reverse-transcription mix, 10× Taq buffer with KCl, 25 mM MgCl₂, 10 mM deoxy-ribonucleotide mix, Taq DNA polymerase, 10 pmol/L of each sense and antisense primer, and ddH₂O (Takara Bio, Otsu, Japan). Aliquots (10 µL) of the amplification products were resolved by 1.5% agarose gel (Promega, Fitchburg, WI, USA) electrophoresis and visualized by ethidium bromide staining. The fragment size and signal intensity were analyzed by GeneScan Analysis and GeneScan Genotyper software (Life Technologies).

Western blotting

Extracting proteins from tissues

Proteins were extracted in a homogenate suspension buffer consisting of 10 mM Tris-HCl, pH 7.6, 100 mM NaCl, and a protease-inhibitor cocktail (Complete; Roche Diagnostics, Rotkreuz, Switzerland). Protein concentration was measured with a BCA protein assay (Thermo Fisher Scientific).

Western blotting

About 50 µg of protein extract was subjected to sodium dodecyl sulfate–polyacrylamide gel electrophoresis and transferred to nitrocellulose filters by electroblotting. Nonspecific binding was blocked by incubating the membranes in 5% nonfat milk in TBS-T (20 mM Tris-HCl, 150 mM NaCl, 0.1% Tween-20). Membranes were incubated with the rabbit anti-Hpa polyclonal antibody (1:100 in a dilution) (Boster Biotechnology, Wuhan, People's Republic of China) overnight at 4°C. Membranes were then washed with TBS-T and incubated with a horseradish peroxidase-conjugated rabbit anti-goat immunoglobulin G (1:2000 dilution) (Boster Biotechnology) for 1 hour at room temperature. Signals were detected by enhanced chemiluminescence (GE Healthcare, Piscataway, NJ, USA) according to the instructions of the manufacturer, and the data were analyzed using the Un-Scan-It program (Silk Scientific, Orem, UT, USA). The same membrane was reprobated with β-actin-specific antibody to ensure equal control.

Inhibitory effect of Fe₃O₄-dextran-anti-βHCG-Hpa ASODN on hypodermal transplant tumors in nude mice

Cultured JEG-3 cells were harvested and resuspended in unsupplemented RPMI medium at 4–6 × 10⁶ cells/mL, then 100 µL was immediately injected into the flanks of 20 nude mice. The mice were divided into four groups, five in each, and further growth of induced tumors was monitored for 10 days. Tumor diameters measured 8–10 mm in all groups. Four different types of solutions were injected hypodermally in each nude mouse: group 1, 0.2 mL Fe₃O₄-dextran-anti-βHCG-Hpa ASODN solutions, which contained 20 µmol/L Hpa ASODN; group 2, 0.2 mL Fe₃O₄-dextran-anti-βHCG-Hpa NSODN solutions, which contained 20 µmol/L Hpa NSODN; group 3, 0.2 mL Fe₃O₄-dextran-anti-βHCG solutions; and group 4, 0.2 mL saline control. These solutions were injected every 2 days into all nude mice for 20 days (ten times) aggregately. We weighed the transplanted tumor of every nude mouse. Inhibition rate (%) = [(the weight of transplant tumor of control group – the weight of transplant tumor of curing group)/the weight of transplant tumor of control group] × 100%.

Statistical analysis

Results were shown as means ± standard error of mean. The statistical significance of differences in mean values was assessed by Student's *t*-test or one-way analysis of variance followed by the Dunnett's multiple-comparison test using SAS software (SAS Institute, Cary, NC, USA). Differences among means were considered significant at *P* < 0.05.

Results

Effect of Fe₃O₄-dextran-anti-βHCG-Hpa ASODN compounds on proliferative ability of JEG-3

As shown in Table 1, after 24 hours there were not any significant differences among proliferative abilities of untransfected Hpa NSODN and Hpa ASODN JEG-3 cells (*P* > 0.05).

Table 1 Difference in proliferative ability of untransfected and transfected JEG-3 cells

Group	24 hours	48 hours	72 hours
Untransfected	0.317 ± 0.02	0.646 ± 0.04	0.871 ± 0.05
Hpa NSODNs	0.311 ± 0.03	0.613 ± 0.05	0.860 ± 0.04
Hpa ASODNs	0.308 ± 0.02	0.402 ± 0.02	0.471 ± 0.03

Abbreviations: Hpa, heparanase; NSODN, nonspecific oligodeoxynucleotide; ASODN, antisense oligodeoxynucleotide.

At the same time, we found that after 48 and 72 hours, the proliferative ability of Hpa ASODN JEG-3 cells was obviously inhibited ($P < 0.05$), whereas there were not any significant differences between the untransfected group and the Hpa NSODN group ($P > 0.05$).

Effect of Fe₃O₄-dextran-anti-βHCG-Hpa ASODN compounds on invasive ability of JEG-3 cells

As shown in Figure 1, invasive ability was decreased in JEG-3 cells treated by Fe₃O₄-dextran-anti-βHCG-Hpa ASODN (the average cell number on the lower chamber was 109 ± 10) compared to Hpa NSODN (213 ± 18) and untransfected control JEG-3 cells (221 ± 25) ($P < 0.01$). There were not any significant differences between Fe₃O₄-dextran-anti-βHCG-Hpa NSODN-treated JEG-3 cells and untransfected control JEG-3 cells ($P > 0.05$).

Effect of Fe₃O₄-dextran-anti-βHCG-Hpa ASODN compounds on Hpa mRNA expression in JEG-3 cells

As shown in Figure 2, there was an almost 2.7-fold decrease in Hpa mRNA in JEG-3 cells when these cells were treated with Hpa ASODN (0.481 ± 0.04 vs 1.288 ± 0.12 , $P < 0.01$). This decrease in Hpa mRNA is sequence-specific because the NSODN-treated JEG-3 cells produced equivalent amounts

of Hpa mRNA transcripts (1.117 ± 0.10 vs 1.288 ± 0.12 , $P > 0.05$).

Effect of Fe₃O₄-dextran-anti-βHCG-Hpa ASODN compounds on Hpa protein expression in JEG-3 cells

Confirmation of decreased translation of Hpa protein was demonstrated by Western blot analysis. As shown in Figure 3, Hpa protein decreased in JEG-3 cells treated with magnetic nanoparticle Hpa ASODNs (16.1 ± 1.4) compared to Fe₃O₄-dextran-anti-βHCG-Hpa NSODNs (35.3 ± 3.2) and untransfected control JEG-3 cells (39.7 ± 2.9) ($P < 0.05$). But there was no significant difference between Hpa protein in JEG-3 cells treated with Fe₃O₄-dextran-anti-βHCG-Hpa NSODNs and that in untransfected control JEG-3 cells ($P > 0.05$).

Effect of Fe₃O₄-dextran-anti-βHCG-Hpa ASODN on hypodermal transplant tumors of nude mice

As shown in Figure 4, after being treated by Fe₃O₄-dextran-anti-βHCG-Hpa ASODN, the growth of hypodermal transplant tumors was obviously inhibited ($P < 0.01$; the average weight was 0.615 ± 0.061 g). The inhibitory rate was 58.6%, and compared with the control group (the average weight was 1.487 ± 0.159 g), there was a significant difference ($P < 0.01$). The average weight of hypodermal transplant tumors in the Fe₃O₄-dextran-anti-βHCG-Hpa NSODN group and the Fe₃O₄-dextran-anti-βHCG nanoparticle group was 1.417 ± 0.136 g and 1.428 ± 0.145 g, respectively, and their inhibitory rates were 4.7% and 4.0%, respectively. Compared with the control group, the growth of hypodermal transplant tumors of the Fe₃O₄-dextran-anti-βHCG-Hpa NSODN group and the Fe₃O₄-dextran-anti-βHCG nanoparticle group was not obviously inhibited ($P > 0.05$).

Figure 1 (A–C) Effects of Fe₃O₄-dextran-anti-β-human chorionic gonadotropin (HCG) carrying heparanase (Hpa) antisense oligodeoxynucleotide (ASODN) on JEG-3 cell invasion. **(A)** Untransfected cells. Cells on the lower surface of the insert chamber were counted under microscope with 80× magnification. **(B)** Fe₃O₄-dextran-anti-βHCG-Hpa ASODN-treated cells. **(C)** Fe₃O₄-dextran-anti-βHCG-Hpa NSODN-treated cells.

Abbreviation: NSODN, non-sense oligonucleotides.

Discussion

The key role played by Hpa in malignancy was confirmed by antisense complementary DNA introduction studies that showed a significant reduction in the invasive and metastatic properties of tumor cells.⁷ Expression of an anti-Hpa ribozyme construct in human MDA-MB-435 breast carcinoma cells significantly reduced Hpa enzyme activity as well as invasion through a Matrigel basement membrane.⁸ A large number of publications now clearly link Hpa expression to the process of tumorigenesis in a wide range

Figure 2 (A–C) Effect of Fe_3O_4 -dextran-anti- β -human chorionic gonadotropin (HCG) carrying heparanase (Hpa) antisense oligodeoxynucleotide (ASODN) compounds on Hpa messenger RNA (mRNA) expression in JEG-3 cells. **(A)** Hpa mRNA expression in JEG-3 cells. **(B)** Hpa mRNA expression in JEG-3 cells after been treated with Fe_3O_4 -dextran-anti- β HCG-Hpa ASODNs. **(C)** Hpa mRNA expression in JEG-3 cells after been treated with Fe_3O_4 -dextran-anti- β HCG-Hpa NSODNs.

Abbreviations: GAPDH, glyceraldehyde-3-phosphodehydrogenase; NSODN, non-sense oligonucleotides.

of cancers, including bladder,⁹ colon,¹⁰ gastric,¹¹ breast,¹² oral,¹³ esophageal,¹⁴ pancreatic,¹⁵ brain,¹⁶ thyroid,¹⁷ prostate,¹⁸ and acute myeloid leukemia.¹⁹ Collectively, this evidence suggests that Hpa plays a fundamental role in sustaining the pathology of malignant diseases, and therefore that it may provide a potential target for anticancer therapy.

Human gene therapy is one of those new therapeutic approaches emerging from this molecular biology and biotechnology revolution.²⁰ ASODN is one gene-therapy approach; it offers the ability to regulate genes involved in cancer progression, especially those that are not amenable to small-molecule or antibody inhibition.²¹ ASODNs are designed to be complementary to a selected gene's mRNA and thereby specifically inhibit expression of that gene. It is estimated that any sequence of at least 13 bases in RNA and 17 bases in DNA is represented only once within the human genome. Thus, the specificity implicit in the design of ASODNs theoretically leads to decreased off-target toxicity. ASODN technology enables rapid movement from preclinical models to testing in the clinic. Challenges remain

to optimize tissue exposure and cellular uptake, and to demonstrate mechanisms and antitumor activity. Therefore, it is very important for us to find a targeted vector.

Magnetic Fe_3O_4 nanoparticles are considered ideal gene vectors, because they are simple to manufacture, have diameters of 10 nm, are able to absorb significant amounts of DNA, demonstrate limited cellular toxicity, and can be guided to specific sites via magnets. In addition, to improve biocompatibility, Fe_3O_4 nanoparticles can be embedded in various macromolecules, such as polysaccharides and proteins. However, human organs, such as liver, spleen, and lymph (among others), are infused with a reticuloendothelial system and have the ability to absorb nanoparticles passively through phagocytosis. Because β HCG is expressed only in choriocarcinomas, anti- β HCG monoclonal antibodies were attached to magnetic particles to allow the nanoparticles to target the tissues of interest.

We have confirmed that magnetic nanoparticles represent a versatile vector for gene therapy, with high stability towards degradation. They combine a high introduction potential with good biocompatibility and low cytotoxicity. With magnetic and immune targeting, they offer the pos-

Figure 3 (A–C) Effect of Fe_3O_4 -dextran-anti- β -human chorionic gonadotropin (HCG) carrying heparanase (Hpa) antisense oligodeoxynucleotide (ASODN) compounds on Hpa protein expression in JEG-3 cells. **(A)** Hpa protein expression in JEG-3 cells. **(B)** Hpa protein expression in Fe_3O_4 -dextran-anti- β HCG-Hpa ASODN-treated cells. **(C)** Hpa protein expression in Fe_3O_4 -dextran-anti- β HCG-Hpa non-sense oligonucleotides-treated cells.

Figure 4 (A–D) Effect of Fe_3O_4 -dextran-anti- β -human chorionic gonadotropin (HCG) carrying heparanase (Hpa) antisense oligodeoxynucleotide (ASODN) on hypodermal transplant tumors in nude mice. **(A)** Saline solution-treated hypodermal transplant tumor. **(B)** Fe_3O_4 -dextran-anti- β HCG nanoparticle-treated hypodermal transplant tumor. **(C)** Fe_3O_4 -dextran-anti- β HCG-Hpa nonspecific oligodeoxynucleotide-treated hypodermal transplant tumor. **(D)** Fe_3O_4 -dextran-anti- β HCG-Hpa ASODN-treated hypodermal transplant tumor.

sibility of additional targeting strategies.⁵ In this study, we used MTT and Matrigel invasion assay to detect changes in the proliferative and invasive abilities of choriocarcinoma JEG-3 cells after being treated by magnetic nanoparticle Hpa ASODN compounds. The results revealed that the invasive ability of JEG-3 cells was inhibited, and that after 48 and 72 hours, the proliferative ability was also obviously inhibited. Subsequently, we found that magnetic nanoparticle Hpa ASODN compounds can weaken the expression of Hpa mRNA and protein. Finally, we tested the effect of magnetic nanoparticle Hpa ASODN compounds on choriocarcinoma hypodermal tumors of nude mice. The results indicated that magnetic nanoparticle Hpa ASODN compounds can inhibit the growth of choriocarcinoma tumors. All of these results were identical with expectations. However, an unexpected outcome was the finding that proliferative ability after 24 hours was not affected. It was difficult for us to explain this result. Our initial view was that magnetic nanoparticle Hpa ASODN compounds did not produce any effect on JEG-3 cell proliferation after 24 hours.

Transfected by Fe₃O₄-dextran-anti-βHCG magnetic nanoparticles, Hpa ASODNs inhibited the expression of JEG-3 cells' Hpa directly, which hints that Fe₃O₄-dextran-anti-βHCG magnetic nanoparticles are a kind of effective gene vector. Subsequently, we found that the invasive and proliferative abilities of JEG-3 cells and the growth of choriocarcinoma hypodermal tumors of nude mice had been obviously controlled, and this result should be connected with the inhibition of Hpa expression. The reason is that the growth of tumors is not only relevant to the proliferation of tumor cells but is also associated with the invasion of cells. If tumor cells want to invade surrounding organs, it is necessary for them to dissolve extracellular matrix. Thus when the expression of Hpa is inhibited, the solvency of tumor cells is weakened, and so the growth of tumor cells is inhibited also. On the other hand, we found that magnetic nanoparticle Hpa ASODN compounds do not have any toxicity to nude mice, so it is reasonable for us to consider this as a safe and effective tumor remedy.

Acknowledgments

This project was supported by the National Natural Science Foundation of China (grant 81101995) and the Natural Science Foundation of Hunan Province (grant 13JJ4120).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Vlodavsky I, Friedmann Y, Elkin M, et al. Mammalian heparanase: gene cloning, expression and function in tumor progression and metastasis. *Nat Med*. 1999;5:793–802.
2. Vlodavsky I, Friedmann Y. Molecular properties and involvement of heparanase in cancer metastasis and angiogenesis. *J Clin Invest*. 2001;108:341–347.
3. Jingting C, Yangde Z, Yi Z, Huining L, Rong Y, Yu Z. Heparanase expression correlates with metastatic capability in human choriocarcinoma. *Gynecol Oncol*. 2007;107:22–29.
4. Zhang YD. *Nanobiotechnology*. Beijing: Scientific Publishing; 2005.
5. Jingting C, Huining L, Yi Z. Preparation and characterization of magnetic nanoparticles containing Fe₃O₄-dextran-anti-β-human chorionic gonadotropin, a new generation choriocarcinoma-specific gene vector. *Int J Nanomedicine*. 2011;6:285–294.
6. Staff AC, Ranheim T, Henriksen T, Halvorsen B. 8-Iso-prostaglandin F_{2α} reduces trophoblast invasion and matrix metalloproteinase activity. *Hypertension*. 2000;35:1307–1313.
7. Uno F, Fujiwara T, Takata Y, et al. Antisense-mediated suppression of human heparanase gene expression inhibits pleural dissemination of human cancer cells. *Cancer Res*. 2001;61:7855–7860.
8. Edovitsky E, Elkin M, Zcharia E, Peretz T, Vlodavsky I. Heparanase gene silencing, tumor invasiveness, angiogenesis, and metastasis. *J Natl Cancer Inst*. 2004;96:1219–1230.
9. Gohji K, Okamoto M, Kitazawa S, et al. Heparanase protein and gene expression in bladder cancer. *J Urol*. 2001;166:1286–1290.
10. Sato T, Yamaguchi A, Goi T, et al. Heparanase expression in human colorectal cancer and its relationship to tumor angiogenesis, hematogenous metastasis, and prognosis. *J Surg Oncol*. 2004;87:174–181.
11. Takaoka M, Naomoto Y, Ohkawa T, et al. Heparanase expression correlates with invasion and poor prognosis in gastric cancers. *Lab Invest*. 2003;83:613–622.
12. Zhang YL, Fu ZR, Chen XY, Yang JH, Wang YH, Shen Q. The inhibitory effect of heparanase antisense oligodeoxynucleotides on the invasiveness of human mammary carcinoma cell line. *Zhonghua Shi Yan Wai Ke Za Zhi*. 2004;3:367–369.
13. Ikuta M, Podyma KA, Maruyama K, Enomoto S, Yanagishita M. Expression of heparanase in oral cancer cell lines and oral cancer tissues. *Oral Oncol*. 2001;37:177–184.
14. Mikami S, Ohashi K, Usui Y, et al. Loss of syndecan-1 and increased expression of heparanase in invasive esophageal carcinomas. *Jpn J Cancer Res*. 2001;92:1062–1073.
15. Kim AW, Xu X, Hollinger EF, Gattuso P, Godellas CV, Prinz RA. Human heparanase-1 gene expression in pancreatic adenocarcinoma. *J Gastrointest Surg*. 2002;6:167–172.
16. Marchetti D, Nicolson GL. Human heparanase: a molecular determinant of brain metastasis. *Adv Enzyme Regul*. 2001;41:343–359.
17. Xu X, Quiros RM, Maxhimer JB, et al. Inverse correlation between heparan sulfate composition and heparanase-1 gene expression in thyroid papillary carcinomas: a potential role in tumor metastasis. *Clin Cancer Res*. 2003;9:5968–5979.
18. Ogishima T, Shiina H, Breault JE, et al. Increased heparanase expression is caused by promoter hypomethylation and up-regulation of transcriptional factor early growth response-1 in human prostate cancer. *Clin Cancer Res*. 2005;11:1028–1036.
19. Vlodavsky I, Goldshmidt O, Zcharia E, et al. Mammalian heparanase: involvement in cancer metastasis, normal development and angiogenesis. *Semin Cancer Biol*. 2002;12:121–129.
20. Anderson WF. Human gene therapy. *Nature*. 1998;392:25–30.
21. Gleave ME, Monia BP. Antisense therapy for cancer. *Nat Rev Cancer*. 2005;5:468–479.

International Journal of Nanomedicine**Dovepress****Publish your work in this journal**

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-nanomedicine-journal>