Impact of AT₂-receptor stimulation on vascular biology, kidney function, and blood pressure

Leon A Danyel¹
 Patrick Schmerler¹
 Ludovit Paulis^{1–3}
 Thomas Unger⁴
 U Muscha Steckelings^{1,5}

¹Center for Cardiovascular Research, Institute of Pharmacology, Charité Medical Faculty, Berlin, Germany;

²Institute of Pathophysiology, Faculty of Medicine, Comenius University, Bratislava, Slovak Republic; ³Institute of Normal and Pathological Physiology, Slovak Academy of Sciences, Bratislava, Slovak Republic;

⁴CARIM, Maastricht University, Maastricht, the Netherlands; ⁵Institute of Molecular Medicine, Department of Cardiovascular and Renal Physiology, University of Southern Denmark, Odense, Denmark

Abstract: The angiotensin type 2 receptor (AT₂R) and the receptor MAS are receptors within the renin–angiotensin system, which mediate tissue-protective actions such as anti-inflammation, antifibrosis, and antiapoptosis. In recent years, several programs have been launched in order to develop drugs that act as agonists on the AT₂R or MAS to take therapeutic advantage of the protective and regenerative properties of these receptors. This review article will focus on recent data obtained in preclinical animal and in vitro models with new AT₂R-agonistic molecules (Compound 21 and β -amino acid substituted angiotensin II) and with relevance for blood pressure (BP) regulation or hypertensive end-organ damage. These data will include studies on vasodilation/vasoconstriction in isolated resistance arteries ex vivo, studies on kidney function, studies on vascular remodeling, and studies that measured the net effect of AT₂R stimulation on BP in vivo. Current data indicate that although AT₂R stimulation causes vasodilation ex vivo and promotes natriuresis, it does not alter BP levels in vivo acutely – at least as long as there is no additional low-dose blockade of AT₁R. However, AT₂R stimulation alone is able to attenuate hypertension-induced vascular remodeling and reduce arterial stiffening, which in more chronic settings and together with the natriuretic effect may result in modest lowering of BP. We conclude from these preclinical data that AT₂R agonists are not suitable for antihypertensive monotherapy, but that this new future drug class may be beneficial in combination with established antihypertensives for the treatment of hypertension with improved protection from end-organ damage.

Keywords: renin–angiotensin system, AT₂-receptor, vasodilation, blood pressure, kidney function, vascular remodeling

Introduction

The renin–angiotensin system (RAS) is essentially involved in the control of blood pressure (BP) and body volume.¹ Angiotensin II (Ang II), acting via the angiotensin type 1 receptor (AT₁R), causes vasoconstriction as well as sodium and water retention.² Furthermore, it is involved in the pathogenesis of hypertensive and diabetic end-organ damage by promoting inflammation and fibrosis. Pharmacological interference with the RAS by direct renin inhibitors, angiotensin-converting enzyme (ACE) inhibitors, or AT₁R blockers (ARB) is a common therapeutic approach for the treatment of hypertension.³ In addition, ACE inhibitors and ARBs are standard treatment for heart failure and diabetic nephropathy.

In the recent past, research on potential new drug targets within the RAS and the development of respective novel drugs have gained significant momentum.^{4–6} These current efforts mainly aim at making therapeutic use of the so-called protective RAS,

Correspondence: U Muscha Steckelings
 Institute of Molecular Medicine,
 Department of Cardiovascular and
 Renal Physiology, University of Southern
 Denmark, JB Winslows Vej 21–3,
 5000 Odense C, Denmark
 Tel +45 6550 3760
 Fax +45 6613 3479
 Email usteckelings@health.sdu.dk

which comprises the AT2R and the ACE2-angiotensin-(1-7) (Ang-[1-7])-MAS axis. Both the AT2R and the receptor MAS mediate a broad array of tissue-protective effects, including anti-inflammation, antifibrosis, antiapoptosis, neuroprotection, favorable metabolic effects, and vasodilation. In particular, their ability to counteract vasoconstriction, inflammation, and fibrosis makes the AT2R and MAS potential drug targets for the treatment of hypertension and related end-organ damage. Interestingly, there seems to be a positive feedback loop within the protective RAS because, as has been published very recently, the expression of ACE2, Ang-(1-7), and MAS, as well as ACE2 activity, were increased in the kidneys of obese Zucker rats treated for 2 weeks with the AT2R agonist CGP 42112A.⁷

Regarding the AT2R, there are currently three types of new agonistic molecules with the potential for drug development: 1) the nonpeptide small molecule agonist Compound 21 (C21; Vicore Pharma, Gothenburg, Sweden, www.vicorepharma.com),⁸ 2) the cyclic Ang II derivative (one amino acid exchanged for another, unknown [unpublished] amino acid) LP2-3 (Lanthio Pharma, Groningen, the Netherlands, www.lanthiopep.nl),⁹ and 3) a group of Ang II derivatives in which individual amino acids in the sequence of native Ang II are substituted by the respective β -amino acid.¹⁰ C21 is currently in the final stage of preclinical development and is expected to enter clinical testing in 2014, the status of LP2-3 is unknown, and the β -amino acid substituted molecules are currently used only for academic purposes.

This review article will discuss physiology and potential therapeutic use of the AT2R with a focus on its role in BP regulation and hypertensive end-organ damage.

Vasodilation and blood pressure

Stimulation of the AT2R has been shown to act in a vasodilatory way in various species and multiple vascular beds such as mesenteric,^{11–15} renal,^{16–18} coronary,¹⁹ cerebral,²⁰ cutaneous,²¹ and uterine arteries.^{22,23} In addition, AT2R knockout mice exhibit higher basal BP levels than wild-type mice, and they react with a stronger increase in BP to infusion of Ang II.^{24,25} Conversely, in mice overexpressing AT2Rs in the vasculature, the pressor response to Ang II is markedly impaired.²⁶

Vasodilation was also shown for more recently developed AT2R agonists such as the first nonpeptide agonist C21⁸ or for the new peptide agonists generated by substituting individual amino acids in the sequence of native Ang II by the respective β -amino acid (Table 1).⁹ Vasodilation in response to these new ligands at concentrations between 10^{-11} and 10^{-6} M was

observed in aorta from normotensive mice or hypertensive rats and in mouse mesenteric arteries in an AT2R-dependent manner, because these effects could be blocked by the AT2R antagonist PD 123319.^{10,27} Vasodilation in response to C21 was also reported by Verdonk et al²⁸ in coronary, iliac, and mesenteric arteries of rats and mice; however, only at concentrations between 10^{-6} and 10^{-3} M. This vasodilatory response of C21 was unrelated to the AT2R because it could not be blocked by the AT2R antagonist PD 123319, which with 1 μ mol/L, however, was underdosed for most tested concentrations of C21, and because it was absent in AT2R-deficient mice. The mechanism of AT2R-unrelated vasodilation in response to C21 is still unknown but may involve blockade of calcium transport into the cell. Vasodilation caused by very high concentrations of C21 was preceded by a short vasoconstriction, which was obviously due to AT1R stimulation, because it could be blocked by AT1R antagonists.²⁸ This observation was not surprising, as Bosnyak et al²⁷ had previously described that C21 stimulates the AT1R at very high concentrations, resulting in a rise in BP. In general, most small molecule drugs lose specificity at concentrations >1 μ M or even lower. Well-known examples are β_2 -receptor mimetics used for the treatment of asthma, or β_1 -receptor blockers used for the treatment of hypertension, both of which bind to the respective other receptor subtype at high concentrations. For example, the K_i of the β_2 -receptor mimetic salmeterol, which possesses the highest selectivity of this drug class, is 24.6 nM for the β_2 -receptor and 1,600 nM for the β_1 -receptor; the selective β_1 -receptor blocker metoprolol has a K_i of 47 nM for the β_1 -receptor and of 2.960 nM for the β_2 -receptor.^{29,30} With a K_i for the AT2R of 0.4 nM (2 nM for the human receptor) and $>10,000$ nM for the AT1R, the selectivity of C21 is not any worse than selectivity of so-called selective β -blockers or β_2 -receptor mimetics.

It seems logical that a hormone or drug that is able to cause vasodilation in isolated blood vessels would also be able to lower BP in animals and/or humans, as the vascular tone in resistance arteries is a major determinant of systolic BP. However, for the AT2R, this seems not to be the case (Table 2). Several studies using peptide or nonpeptide AT2R agonists, including the new molecules discussed previously, demonstrated that an AT2R-mediated decrease in BP occurred only when there was a concomitant low-dose blockade of AT1Rs. For example, this was shown by the Carey et al³¹ in conscious, normotensive Sprague Dawley rats and by Barber et al³² in conscious, normotensive Wistar Kyoto and spontaneously hypertensive rats (SHR) in a PD 123319-reversible manner.^{10,27} The only exceptions were BP measurements in anesthetized rats, which were part of the first

Table 1 Effects of new AT₂R agonists on vascular tone *ex vivo*

AT ₂ agonist	Type of vessel	Species	Effective concentration	Result	Reference
β-substituted Ang peptides	Aorta	Mouse	10 ⁻¹² to 10 ⁻⁶ M	Vasodilation in presence of AT ₁ R blocker	8
β-substituted Ang peptides	Aorta	Mouse	10 ⁻¹² to 10 ⁻⁶ M	Vasodilation in absence of AT ₁ R blocker	8
C21	Aorta	Mouse	10 ⁻¹⁰ to 10 ⁻⁶ M	Vasodilation in presence of AT ₁ R blocker	14
C21	Aorta	SHR	10 ⁻¹¹ to 10 ⁻⁶ M	Vasodilation in presence of AT ₁ R blocker	14
C21	Mesenteric artery	Mouse	10 ⁻¹⁰ to 10 ⁻⁶ M	Vasodilation in presence of AT ₁ R blocker	14
C21	Aorta	Mouse	10 ⁻⁰⁹ to 10 ⁻⁶ M	Vasodilation in absence of AT ₁ R blocker	14
C21	Coronary microartery	Human	10 ⁻⁰⁶ to 10 ⁻⁴ M	Vasodilation in absence of AT ₁ R blocker	15
C21	Iliac artery	Wistar rat	10 ⁻⁰⁷ to 10 ⁻⁴ M	Vasodilation in absence of AT ₁ R blocker	15
C21	Mesenteric artery	Wistar rat	10 ⁻⁰⁷ to 10 ⁻⁴ M	Vasodilation in absence of AT ₁ R blocker	15
C21	Mesenteric artery	SHR	10 ⁻⁰⁸ to 10 ⁻⁴ M	Vasodilation in absence of AT ₁ R blocker	15
C21	Iliac artery	Wistar rat	10 ⁻¹⁰ to 10 ⁻⁴ M	No effect	15
C21	Iliac artery	SHR	10 ⁻⁰⁵ to 10 ⁻⁴ M	Vasoconstriction	15
C21	Iliac artery	Mouse	10 ⁻⁰⁷ to 10 ⁻⁴ M	Vasodilation in absence of AT ₁ R blocker	15
C21	Iliac artery	AT ₂ -KO	10 ⁻⁰⁷ to 10 ⁻⁴ M	Vasodilation in absence of AT ₁ R blocker	15
C21	Coronary arteries	Wistar rat	10 ⁻⁰⁴ to 10 ⁻³ M	Initial inhibition of coronary flow	15
C21	Coronary arteries	Wistar rat	10 ⁻³ M	Increase in coronary flow	15
C21	Coronary arteries	SHR	10 ⁻⁰⁵ to 10 ⁻³ M	Initial inhibition of coronary flow	15
C21	Coronary arteries	SHR	10 ⁻⁰⁴ to 10 ⁻³ M	Increase in coronary flow	15
C21	Coronary arteries	Mouse (wild-type and AT ₂ -KO)	10 ⁻¹⁰ to 10 ⁻³ M	No effect	15
C21	Mesenteric arteries	SHR	10 ⁻¹⁰ to 10 ⁻⁵ M	Vasodilation in presence of AT ₁ R blocker	32

Abbreviations: AT₂R, angiotensin type 2 receptor; KO, knockout; SHR, spontaneously hypertensive rats; AT₁R, angiotensin type 1 receptor.

description of design and synthesis of C21.⁸ In this study, C21 caused a fall in BP of up to 25 mmHg. However, it has to be noted that anesthesia probably caused unphysiological effects of this pharmacological intervention with the RAS, which is activated by anesthesia.³³

The observation that AT₂R stimulation lowers BP only in the presence of low-level AT₁R blockade indicates that *in vivo* a continuous angiotensinergic tone mediated via the AT₁R seems dominant over any vasodilatory effect of AT₂Rs. Consequently, AT₂R agonists will most likely not become antihypertensive drugs suitable for monotherapy. However, due to their tissue-protective effects discussed as follows, the combination of established antihypertensives with AT₂R agonists may result in better long-term prevention of hypertensive end-organ damage. Moreover, long-term therapy with AT₂R agonists may have a modest BP-lowering effect due to structural changes of

the vessel walls (see paragraph about “Vascular remodeling” below) and due to a recently described diuretic effect (see paragraph about “AT₂ receptor activation in renal physiology and disease” below).^{34–36} AT₂R stimulation may further act antihypertensive by CNS related mechanisms that become apparent only if C21, which crosses the blood–brain barrier only very poorly, is applied intracerebroventricularly.³⁷

Normal pregnancy and preeclampsia are conditions during which expression of components of the RAS is altered in a way that the peripheral and tissue RAS are activated.³⁸ In preeclampsia in patients or in respective animal models, the ratio of expression of AT₁R and AT₂R changes in favor of the AT₁R in placental and uterine tissue and arteries.^{39–41} The AT₂R has been described to act in a vasodilatory manner and to counteract the enhanced vasoconstrictive effect of Ang II via the AT₁R during pregnancy.²² Thus, it can

Table 2 Effects of new AT2R agonists on blood pressure

AT2 agonist	Species/strain	Dosage	Result	Reference
β-substituted Ang peptides	SHR	15 pmol/kg/min IV	MAP ↓ only in presence of a low-dose AT1R blocker	8
C21	SHR	100/300 ng/kg/min IV	MAP ↓ only in presence of a low-dose AT1R blocker; AT2-dependent	14
C21	SHR	1,000 ng/kg/min IV	MAP ↑; AT1R-dependent	14
C21	Wistar–Kyoto rats	50–300 ng/kg/min IV	No effect	14
C21	Sprague Dawley rats	100–300 ng/kg/min IV	No effect	22
C21	Obese Zucker rats	1 μg/kg/min IV	No effect	23
C21	Sprague Dawley rats	0.5 μg/μl/h ICV	MAP ↓ by central mechanisms	24
C21	L-NAME-induced hypertension	0.3 mg/kg BW IP	No effect	31
C21	SHR	1 mg/kg BW PO	Increase after 1 week; no effect in weeks 2–5	32
C21	2K1C hypertension	0.3 mg/kg BW IP	No effect	39
C21	SHR-SP	10 mg/kg/day extended release from 0.5% Na-carboxymethylcellulose	No effect	40
C21	Wistar rats	0.03/0.3 mg/kg BW IP	No effect	59
C21	C57Bl-6	1/3/10 μg/kg BW IP	No effect	60
C21	Obese Zucker rats	300 μg/kg/day IP	No effect	61
C21	KK-Ay mice	10 μg/kg BW IP	No effect	62

Abbreviations: 2K1C, two-kidney, one-clip rat model; AT2R, angiotensin type 2 receptor; BW, body weight; ICV, intracerebroventricularly; IP, intraperitoneally; IV, intravenously; L-NAME, N ω -nitro-L-arginine methyl ester hydrochloride; PO, orally; SHR, spontaneously hypertensive rats; SP, stroke-prone; AT1R, angiotensin type 1 receptor; MAP, mean arterial pressure.

be speculated that the relative deficiency in AT2R during preeclampsia may contribute to the development of hypertension. In a recent study, Hladunewich et al⁴² could, in fact, demonstrate that there is a strong correlation between AT1R/AT2R ratio and the change in BP (the higher the ratio, the stronger the increase in BP; $r=0.54$) in women with previous severe preeclampsia who received a graded infusion of Ang II (1–3 ng/kg/min). The importance of the role of the AT2R for the development of preeclampsia still needs confirmation, but if it should turn out to really be of importance, AT2R agonists may be a treatment option of interest, although under “normal” conditions they do not lower BP.

Vascular remodeling

Chronically elevated BP causes an inflammatory response followed by excess synthesis and accumulation of extracellular matrix mainly in the left cardiac ventricle, the kidneys, and the vascular wall. This fibrotic response to hypertension leads to deteriorated organ function and manifests as heart

failure, renal disease, or vascular stiffening, the latter two of which reinforce the development of hypertension, thus establishing a vicious circle.⁴³

Early studies on the role of AT2R stimulation on vascular remodeling could show that the beneficial effects of ARBs were, at least in part, due to indirect AT2R stimulation (use of ARBs leads to an increase in renin release and thus a rise in angiotensin II levels, which, in turn, stimulates the unopposed AT2Rs; ARB-induced Ang II levels are, however, much lower than pharmacological AT2 agonist levels), as these effects could be reversed by blockade of AT2Rs.^{44,45} Further indirect evidence for a favorable role of AT2Rs in vascular remodeling is coming from studies in AT2R-deficient mice, which responded with augmented vascular hypertrophy of coronary, aortic, and femoral arteries to chronically elevated BP.^{46–48}

The impact of direct AT2R stimulation on hypertension-induced vascular remodeling was studied recently in two studies in which C21 was applied orally to rats (Table 3).^{49,50}

Table 3 Effects of new AT2R agonists on vascular remodeling

AT2 agonist	Species/model	Dosage	Result	Reference
C21	L-NAME-induced hypertension	0.3 mg/kg BW IP	Reduced aortic wall thickness and collagen content; lowered pulse wave velocity	31
C21	SHR	1 mg/kg BW PO	Reduced mesenteric artery stiffness; reduced aortic collagen and fibronectin content; lowered oxidative stress	32

Abbreviations: AT2R, angiotensin type 2 receptor; BW, body weight; IP, intraperitoneally; L-NAME, N ω -nitro-L-arginine methyl ester hydrochloride; PO, orally; SHR, spontaneously hypertensive rats.

The authors of these studies used different models of hypertension: ie, stroke-prone spontaneously hypertensive rats (SHR-SP) in one study,⁵⁰ and inhibition of endothelial nitric oxide synthase by application of *N* ω -nitro-L-arginine methyl ester hydrochloride (L-NAME) in the other study.⁴⁹ In both studies, the effect of AT2R stimulation was compared with the effect of an ARB alone or the combination of both. The main finding of both studies was that treatment with an AT2R agonist widely prevented the development of vascular hypertrophy and fibrosis. Remarkably, this reversal of hypertension-induced pathology was achieved without any significant effect on BP: ie, although BP stayed elevated in C21-treated rats, vascular remodeling was almost completely prevented.^{49,50} The preventive effect of AT2R stimulation on vascular remodeling was shown in both studies to be related to a decrease in collagen deposition. Moreover, in our study in L-NAME-induced hypertension in rats, we measured pulse wave velocity, a marker for arterial stiffness and independent predictor of cardiovascular risk in patients, and we could again show that this indicator of vascular remodeling, which was increased in vehicle-treated hypertensive animals, was significantly attenuated in C21-treated, (still) hypertensive animals.⁴⁹

BP in our study was not significantly changed by treatment with C21, but there was a trend toward a reduction in BP, which may have been a result of reduced arterial stiffness.⁴⁹

Kidney function

AT2-receptor deficiency in renal disease and injury

Considering the ubiquitous expression of the AT2R in fetal kidney, it may be surprising that AT2R knockout mice show no renal abnormalities in histology.^{24,51} However, obvious differences in disease progression in AT2R knockout mice when compared with wild-type mice were revealed in the renal ablation model of renal injury.⁵¹ Aggravated glomerular damage and impairment of renal function were shown in AT2R-deficient mice compared with wild-type mice, subsequently culminating in higher overall mortality. Moreover, albuminuria in knockout mice was pronounced and renal macrophage infiltration of glomerulus and interstitium increased compared with wild-type mice. Benndorf et al⁵¹ ruled out systolic BP differences, podocyte or basal membrane damage, or upregulation of AT1Rs as possible causes for enhanced renal injury. Nonetheless, they elucidated one of the possible underlying mechanisms of disease progression in AT2R-deficient mice by showing a significant upregulation

of asymmetric dimethylarginine, an endogenous inhibitor of nitric oxide synthase, in the knockout group, suggesting that the impairment in synthesis of nitric oxide may account for the aggravation of glomerular damage observed.

In accordance with these results, aggravated renal injury was observed in unilateral ureteral obstruction of AT2R-deficient mice, comprising severe interstitial fibrosis and greater abundance of fibroblasts and myofibroblasts in ipsilateral kidneys.⁵²

Moreover, in a model of type 1 diabetes in mice, AT2R knockout animals exhibited accelerated development of diabetic nephropathy.⁵³ Extracellular matrix (ECM) protein accumulation was measured by quantification of periodic acid–Schiff and Masson trichrome staining, as well as real-time quantitative polymerase chain reaction analysis of renal collagen IV messenger ribonucleic acid expression. Terminal deoxynucleotidyl transferase dUTP nick end labeling assay was performed to semiquantify tubular apoptosis between groups. ECM accumulation and renal expression of collagen IV were significantly enhanced in nondiabetic AT2R knockout mice compared with nondiabetic controls. Additionally, tubular apoptosis was significantly increased in nondiabetic AT2R knockout mice. Accordingly, similar observations of increased ECM production and tubular apoptosis in AT2R-deficient animals were made in diabetic knockout animals when compared with wild-type mice, although values reached significance only for Masson trichrome staining arbitrary units. Chang et al⁵³ proposed that an increase in Heme oxygenase 1 expression, observed in proximal tubule cells of AT2R knockout mice, enhances oxidative stress. Moreover, an elevated ACE/ACE2 ratio in knockout animals could account for aggravation of renal injury during diabetic nephropathy.

It is important to note that deterioration of renal injury in AT2R-deficient mice may result from a persistent but yet unnoticed impairment in renal function that evolved during fetal development, due to the absence of the AT2R, which is usually expressed in high density during fetal life, and may not represent missing counter-regulatory actions of the receptor itself in the adult organism.

AT2 receptor activation in renal physiology and disease

So far, studies investigating the role of the AT2R in renal disease were conducted under 1) AT1R blockade with subsequent Ang II administration, 2) using the peptide agonist CGP 42112, or 3) under AT2R blockade achieved by administration of PD 123319. Findings in this regard have been recently

summarized and will not be further discussed in this review.^{54,55} Interpretation of results, however, is confounded by possible antagonistic effects of CGP 42112 and by agonistic effects of PD 123319 on the AT1R or even the AT2R, depending on timing and dosage.^{28,56} Since C21 became available for experimental research, data about the role of the AT2R in kidney function have become more consistent and unequivocally strengthen the idea of a protective role of the AT2R in renal disease and of a natriuretic effect of AT2R stimulation (Table 4).

In Sprague Dawley rats, graded infusion of C21 (peak dosage: 300 ng/kg/min) significantly enhanced renal blood flow by simultaneously reducing renal vascular resistance in both female and male animals.³⁵ In addition, urine flow, urinary sodium excretion, and fractional sodium excretion showed a significant increase when compared with vehicle-treated groups. All effects observed were abolished by concomitant administration of PD 123319. Interestingly, glomerular filtration rate remained stable in the C21 group, albeit the presence of renal vasodilation suggesting that the C21-induced increase in natriuresis is altered due to an effect on tubular function, but not due to hemodynamic effects. All effects were BP independent, as there was no statistically significant difference in BP between vehicle- and C21-treated animals. However, there was a small but significant difference between the C21- and the C21 plus PD 123319-treated animals (BP in PD 123319-treated animals being lower), but from these data it cannot be decided whether this difference was due to an effect of C21 or of PD 123319 or of both.³⁵

An AT2R-dependent but BP-independent natriuretic effect of C21 was further described recently in obese Zucker rats.³⁶ In analogy to the study by Hillard et al,³⁵ glomerular filtration rate remained unchanged in these animals, speaking again for a direct, tubular effect of C21.

In the two-kidney, one-clip rat model of hypertension (2K1C), Matavelli et al⁵⁷ evaluated the effects of AT2R stimulation in early renal inflammation. In 2K1C rats, kidney perfusion is reduced by unilateral clipping of the renal artery. Inflammatory markers of the subsequently

developing ischemia were determined by in vivo recovery levels of renal interstitial fluid. Animals received vehicle, C21 (0.3 mg/kg/day, intraperitoneally), PD 123319 (10 mg/kg/day, osmotic minipump), or C21 plus PD 123319 over the 4-day course of the study. Unilateral stricture of the renal artery led to a significant increase in AT2R protein expression, which was even further enhanced in the C21 group but abolished by administration of PD 123319. Expression of tumor necrosis factor (TNF)- α , transforming growth factor (TGF)- β 1, and interleukin (IL)-6 messenger ribonucleic acid and their renal interstitial fluid (RIF) recovery rates were significantly elevated in 2K1C compared with sham-operated animals. In contrast, nitric oxide and cyclic guanosine monophosphate RIF recovery rates were significantly reduced. Treatment with C21 reduced TNF α , TGF β 1, and IL-6 expression and, moreover, increased RIF recovery rates for nitric oxide and cyclic guanosine monophosphate. Extensive inflammatory cell infiltration observed by histological (hematoxylin/eosin) staining of 2K1C renal cortex and medulla was significantly reduced with C21 treatment when compared with the vehicle group. These effects were only partially inhibited by the AT2R antagonist PD 123319.⁵⁷

SHR-SPs represent an animal model of hypertension, characterized by the progressive development of renal damage and brain abnormality against the background of elevated Ang II blood levels. Oral treatment with C21 (peak dosage group: 10 mg/kg, suspended release from 0.5% sodium carboxymethylcellulose) significantly delayed the development of proteinuria and prevented the accumulation of high-molecular-weight proteins, which present markers of renal inflammation, in 24-hour urine electrophoresis.⁵⁸ Although plasma renin activity increased significantly in the vehicle group, plasma renin activity remained at basal levels in animals treated with C21. Finally, analysis of kidney histopathology revealed reduced renal lesions (attenuation of vascular lesions, tubular damage, luminal cast formation, glomerular sclerosis, inflammatory infiltrates) and attenuated renal macrophage infiltration in rats treated with C21.

Table 4 Effects of new AT2R agonists on kidney function and pathology

AT2 agonist	Species/strain	Dosage	Result	Reference
C21	Sprague Dawley rats	100–300 ng/kg/min IV	Natriuresis, GFR unchanged	22
C21	Obese Zucker rats	5 μ g/kg/min IV	Natriuresis, GFR unchanged	23
C21	2K1C hypertension	0.3 mg/kg BW IP	Attenuated renal inflammation	39
C21	SHR-SP	10 mg/kg/day in 0.5% carboxymethyl-cellulose	Attenuation of albuminuria	40
C21	SHR-SP	10 mg/kg/day in 0.5% carboxymethyl-cellulose	Attenuated renal fibrosis and inflammation	40
C21	Obese Zucker rats	300 μ g/kg/day IP	Attenuated renal inflammation	61

Abbreviations: 2K1C, two-kidney, one-clip rat model; AT2R, angiotensin type 2 receptor; BW, body weight; GFR, glomerular filtration rate; IP, intraperitoneally; IV, intravenously; SHR, spontaneously hypertensive rats; SP, stroke-prone.

Furthermore neoexpression of vimentin, a marker of tubulointestinal injury, was completely prevented by AT2R stimulation with C21.⁵⁸

Summary and conclusion

AT2R stimulation elicits effects on several physiological mechanisms, which contribute to the regulation of BP. Specifically, AT2R agonists induce vasodilation in isolated vessels ex vivo and they enhance natriuresis in vivo. Furthermore, in hypertensive rats, AT2R stimulation attenuates arterial stiffening. Nevertheless, in vivo BP is not altered acutely and only modestly chronically, the latter usually not reaching statistical significance. The lack of acute effect is probably due to some kind of counter-regulatory mechanism(s), which has not been characterized yet, but may involve activation of the “classic” RAS (increased levels of Ang II acting on the AT1R) or of the sympathetic nervous system. The trend toward a reduction in BP in more chronic settings seems likely due to secondary mechanisms such as natriuresis leading to volume reduction or attenuated vascular remodeling, resulting in lower peripheral resistance.

According to these reviewed data from preclinical animal or in vitro/ex vivo studies, it is likely that AT2R agonists will be unsuitable as antihypertensive monotherapeutics in clinical use. However, they may enhance the effectiveness of other, established antihypertensives, and they may provide additive benefit in respect of protection from hypertensive end-organ damage. However, these assumptions need to be verified in future clinical studies.

Disclosure

U Muscha Steckelings received modest research support from Vicore Pharma. The other authors report no conflicts of interest in this work.

References

1. Steckelings UM, Unger T. The renin-angiotensin-aldosterone system. In: Mancina G, Grassi G, Kjeldsen SE, editors. *Manual of Hypertension of the European Society of Hypertension*. 1st ed. London, UK: Informa Healthcare; 2008:110–116.
2. De Gasparo M, Catt KJ, Inagami T, Wright JW, Unger T. International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacol Rev*. 2000;52(3):415–472.
3. Mentz RJ, Bakris GL, Waeber B, et al. The past, present and future of renin-angiotensin aldosterone system inhibition. *Int J Cardiol*. 2013;167(5):1677–1687.
4. Bader M, Santos RA, Unger T, Steckelings UM. New therapeutic pathways in the RAS. *J Renin Angiotensin Aldosterone Syst*. 2012;13(4):505–508.
5. Ferreira AJ, Bader M, Santos RAS. Therapeutic targeting of the angiotensin-converting enzyme 2/angiotensin-(1-7)/Mas cascade in the renin-angiotensin system: a patent review. *Expert Opin Ther Pat*. 2012;22(5):567–574.

6. Steckelings UM, Paulis L, Unger T, Bader M. Emerging drugs which target the renin-angiotensin-aldosterone system. *Expert Opin Emerg Drugs*. 2011;16(4):619–630.
7. Ali Q, Wu Y, Hussain T. Chronic AT2 receptor activation increases renal ACE2 activity, attenuates AT1 receptor function and blood pressure in obese Zucker rats. *Kidney Int*. Epub July 3, 2013.
8. Wan Y, Wallinder C, Plouffe B, et al. Design, synthesis, and biological evaluation of the first selective nonpeptide AT2 receptor agonist. *J Med Chem*. 2004;47(24):5995–6008.
9. Wagenaar GT, Laghmani el H, Fidler M, et al. Agonists of MAS oncogene and angiotensin II type 2 receptors attenuate cardiopulmonary disease in rats with neonatal hyperoxia-induced lung injury. *Am J Physiol Lung Cell Mol Physiol*. 2013;305(5):L341–L351.
10. Jones ES, Del Borgo MP, Kirsch JF, et al. A single beta-amino acid substitution to angiotensin II confers AT2 receptor selectivity and vascular function. *Hypertension*. 2011;57(3):570–576.
11. Dimitropoulou C, White RE, Fuchs L, Zhang H, Catravas JD, Carrier GO. Angiotensin II relaxes microvessels via the AT(2) receptor and Ca(2+)-activated K(+) (BK(Ca)) channels. *Hypertension*. 2001;37(2):301–307.
12. Katada J, Majima M. AT(2) receptor-dependent vasodilation is mediated by activation of vascular kinin generation under flow conditions. *Br J Pharmacol*. 2002;136(4):484–491.
13. Matrougui K, Lévy BI, Henrion D. Tissue angiotensin II and endothelin-1 modulate differently the response to flow in mesenteric resistance arteries of normotensive and spontaneously hypertensive rats. *Br J Pharmacol*. 2000;130(3):521–526.
14. Matrougui K, Loufrani L, Heymes C, Lévy BI, Henrion D. Activation of AT(2) receptors by endogenous angiotensin II is involved in flow-induced dilation in rat resistance arteries. *Hypertension*. 1999;34(4 Pt 1):659–665.
15. Widdop RE, Matrougui K, Levy BI, Henrion D. AT2 receptor-mediated relaxation is preserved after long-term AT1 receptor blockade. *Hypertension*. 2002;40(4):516–520.
16. Arima S, Endo Y, Yaoita H, et al. Possible role of P-450 metabolite of arachidonic acid in vasodilator mechanism of angiotensin II type 2 receptor in the isolated microperfused rabbit afferent arteriole. *J Clin Invest*. 1997;100(11):2816–2823.
17. Endo Y, Arima S, Yaoita H, et al. Function of angiotensin II type 2 receptor in the postglomerular efferent arteriole. *Kidney Int Suppl*. 1997;63:S205–S207.
18. Endo Y, Arima S, Yaoita H, Tsunoda K, Omata K, Ito S. Vasodilation mediated by angiotensin II type 2 receptor is impaired in afferent arterioles of young spontaneously hypertensive rats. *J Vasc Res*. 1998;35(6):421–427.
19. Batenburg WW, Garrelts IM, Bernasconi CC, et al. Angiotensin II type 2 receptor-mediated vasodilation in human coronary microarteries. *Circulation*. 2004;109(19):2296–2301.
20. Haberl RL. Role of angiotensin receptor subtypes in the response of rabbit brain arterioles to angiotensin. *Stroke*. 1994;25(7):1476–1479; discussion 1480.
21. Savoia C, Touyz RM, Volpe M, Schiffrin EL. Angiotensin type 2 receptor in resistance arteries of type 2 diabetic hypertensive patients. *Hypertension*. 2007;49(2):341–346.
22. Pulgar VM, Yamashiro H, Rose JC, Moore LG. Role of the AT2 receptor in modulating the angiotensin-II contractile response of the uterine artery at mid gestation. *J Renin Angiotensin Aldosterone Syst*. 2011;12(3):176–183.
23. Widdop RE, Jones ES, Hannan RE, Gaspari TA. Angiotensin AT2 receptors: cardiovascular hope or hype? *Br J Pharmacol*. 2003;140(5):809–824.
24. Hein L, Barsh GS, Pratt RE, Dzau VJ, Kobilka BK. Behavioural and cardiovascular effects of disrupting the angiotensin II type-2 receptor in mice. *Nature*. 1995;377(6551):744–747.
25. Ichiki T, Labosky PA, Shiota C, et al. Effects on blood pressure and exploratory behaviour of mice lacking angiotensin II type-2 receptor. *Nature*. 1995;377(6551):748–750.

26. Tsutsumi Y, Matsubara H, Masaki H, et al. Angiotensin II type 2 receptor overexpression activates the vascular kinin system and causes vasodilation. *J Clin Invest*. 1999;104(7):925–935.
27. Bosnyak S, Welungoda IK, Hallberg A, Alterman M, Widdop RE, Jones ES. Stimulation of angiotensin AT₂ receptors by the non-peptide agonist, Compound 21, evokes vasodepressor effects in conscious spontaneously hypertensive rats. *Br J Pharmacol*. 2010;159(3):709–716.
28. Verdonk K, Durik M, Abd-Alla N, et al. Compound 21 induces vasorelaxation via an endothelium- and angiotensin II type 2 receptor-independent mechanism. *Hypertension*. 2012;60(3):722–729.
29. Baker JG. The selectivity of beta-adrenoceptor agonists at human beta₁-, beta₂- and beta₃-adrenoceptors. *Br J Pharmacol*. 2010;160(5):1048–1061.
30. Griffin PP, Schubert-Zsilavecz M, Stark H. [Inhibitors of beta-adrenergic receptors: Similarities and differences] Hemmstoffe von beta-Adrenozeptoren: Gemeinsamkeiten und Unterschiede. *Pharmazie in Unserer Zeit*. 2004;33(6):442–449. German.
31. Carey RM, Howell NL, Jin XH, Siragy HM. Angiotensin type 2 receptor-mediated hypotension in angiotensin type-1 receptor-blocked rats. *Hypertension*. 2001;38(6):1272–1277.
32. Barber MN, Sampey DB, Widdop RE. AT₂ receptor stimulation enhances antihypertensive effect of AT₁ receptor antagonist in hypertensive rats. *Hypertension*. 1999;34(5):1112–1116.
33. Efrati S, Berman S, Abu Hamad R, et al. Hyperglycaemia, inflammation, RAS activation: three culprits to blame for acute kidney injury emerging in healthy rats during general anaesthesia. *Nephrology (Carlton)*. 2012;17(7):591–602.
34. Carey RM, Padia SH. Role of angiotensin at₂ receptors in natriuresis: intrarenal mechanisms and therapeutic potential. *Clin Exp Pharmacol Physiol*. 2013;40(8):527–534.
35. Hilliard LM, Jones ES, Steckelings UM, Unger T, Widdop RE, Denton KM. Sex-specific influence of angiotensin type 2 receptor stimulation on renal function: a novel therapeutic target for hypertension. *Hypertension*. 2012;59(2):409–414.
36. Ali Q, Hussain T. AT₂ receptor non-peptide agonist C21 promotes natriuresis in obese Zucker rats. *Hypertens Res*. 2012;35(6):654–660.
37. Gao J, Zhang H, Le KD, Chao J, Gao L. Activation of central angiotensin type 2 receptors suppresses norepinephrine excretion and blood pressure in conscious rats. *Am J Hypertens*. 2011;24(6):724–730.
38. August P, Lenz T, Ales KL, et al. Longitudinal study of the renin-angiotensin-aldosterone system in hypertensive pregnant women: deviations related to the development of superimposed preeclampsia. *Am J Obstet Gynecol*. 1990;163(5 Pt 1):1612–1621.
39. Anton L, Merrill DC, Neves LA, et al. The uterine placental bed renin-angiotensin system in normal and preeclamptic pregnancy. *Endocrinology*. 2009;150(9):4316–4325.
40. Anton L, Merrill DC, Neves LA, et al. Activation of local chorionic villi angiotensin II levels but not angiotensin (1-7) in preeclampsia. *Hypertension*. 2008;51(4):1066–1072.
41. McMullen JR, Gibson KJ, Lumbers ER, Burrell JH. Selective down-regulation of AT₂ receptors in uterine arteries from pregnant ewes given 24-h intravenous infusions of angiotensin II. *Regul Pept*. 2001;99(2–3):119–129.
42. Hladunewich MA, Kingdom J, Odutayo A, et al. Postpartum assessment of the renin angiotensin system in women with previous severe, early-onset preeclampsia. *J Clin Endocrinol Metab*. 2011;96(11): 3517–3524.
43. Schmieder RE. End organ damage in hypertension. *Dtsch Arztebl Int*. 2010;107(49):866–873.
44. Jones ES, Black MJ, Widdop RE. Angiotensin AT₂ receptor contributes to cardiovascular remodelling of aged rats during chronic AT₁ receptor blockade. *J Mol Cell Cardiol*. 2004;37(5):1023–1030.
45. Tea BS, Der Sarkissian S, Touyz RM, Hamet P, deBlois D. Proapoptotic and growth-inhibitory role of angiotensin II type 2 receptor in vascular smooth muscle cells of spontaneously hypertensive rats in vivo. *Hypertension*. 2000;35(5):1069–1073.
46. Akishita M, Iwai M, Wu L, et al. Inhibitory effect of angiotensin II type 2 receptor on coronary arterial remodeling after aortic banding in mice. *Circulation*. 2000;102(14):1684–1689.
47. Brede M, Hadamek K, Meinel L, et al. Vascular hypertrophy and increased P70S6 kinase in mice lacking the angiotensin II AT₂ receptor. *Circulation*. 2001;104(21):2602–2607.
48. Wu L, Iwai M, Nakagami H, et al. Effect of angiotensin II type 1 receptor blockade on cardiac remodeling in angiotensin II type 2 receptor null mice. *Arterioscler Thromb Vasc Biol*. 2002;22(1):49–54.
49. Paulis L, Becker ST, Lucht K, et al. Direct angiotensin II type 2 receptor stimulation in N^ω-nitro-L-arginine-methyl ester-induced hypertension: the effect on pulse wave velocity and aortic remodeling. *Hypertension*. 2012;59(2):485–492.
50. Rehman A, Leibowitz A, Yamamoto N, Rautureau Y, Paradis P, Schiffrin EL. Angiotensin type 2 receptor agonist compound 21 reduces vascular injury and myocardial fibrosis in stroke-prone spontaneously hypertensive rats. *Hypertension*. 2012;59(2):291–299.
51. Benndorf RA, Krebs C, Hirsch-Hoffmann B, et al. Angiotensin II type 2 receptor deficiency aggravates renal injury and reduces survival in chronic kidney disease in mice. *Kidney Int*. 2009;75(10):1039–1049.
52. Ma J, Nishimura H, Fogo A, Kon V, Inagami T, Ichikawa I. Accelerated fibrosis and collagen deposition develop in the renal interstitium of angiotensin type 2 receptor null mutant mice during ureteral obstruction. *Kidney Int*. 1998;53(4):937–944.
53. Chang S-Y, Chen Y-W, Chenier I, Tran SLM, Zhang S-L. Angiotensin II type II receptor deficiency accelerates the development of nephropathy in type I diabetes via oxidative stress and ACE2. *Exp Diabetes Res*. 2011;2011:521076.
54. Jöhren O, Dendorfer A, Dominiak P. Cardiovascular and renal function of angiotensin II type-2 receptors. *Cardiovasc Res*. 2004;62(3): 460–467.
55. Wenzel UO, Krebs C, Benndorf R. The angiotensin II type 2 receptor in renal disease. *J Renin Angiotensin Aldosterone Syst*. 2010;11(1): 37–41.
56. Stoll M, Steckelings UM, Paul M, Bottari SP, Metzger R, Unger T. The angiotensin AT₂-receptor mediates inhibition of cell proliferation in coronary endothelial cells. *J Clin Invest*. 1995;95(2):651–657.
57. Matavelli LC, Huang J, Siragy HM. Angiotensin AT₂ receptor stimulation inhibits early renal inflammation in renovascular hypertension. *Hypertension*. 2011;57(2):308–313.
58. Gelosa P, Pignieri A, Fändriks L, et al. Stimulation of AT₂ receptor exerts beneficial effects in stroke-prone rats: focus on renal damage. *J Hypertens*. 2009;27(12):2444–2451.
59. Kaschina E, Grzesiak A, Li J, et al. Angiotensin II type 2 receptor stimulation: a novel option of therapeutic interference with the renin-angiotensin system in myocardial infarction? *Circulation*. 2008;118(24): 2523–2532.
60. Jing F, Mogi M, Sakata A, et al. Direct stimulation of angiotensin II type 2 receptor enhances spatial memory. *J Cereb Blood Flow Metab*. 2012;32(2):248–255.
61. Dhande I, Ali Q, Hussain T. Proximal tubule angiotensin AT₂ receptors mediate an anti-inflammatory response via interleukin-10: role in renoprotection in obese rats. *Hypertension*. 2013;61(6):1218–1226.
62. Ohshima K, Mogi M, Jing F, et al. Direct angiotensin II type 2 receptor stimulation ameliorates insulin resistance in type 2 diabetes mice with PPAR γ activation. *PLoS One*. 2012;7(11):e48387.

Integrated Blood Pressure Control

Dovepress

Publish your work in this journal

Integrated Blood Pressure Control is an international, peer-reviewed open-access journal focusing on the integrated approach to managing hypertension and risk reduction. Treating the patient and comorbidities together with diet and lifestyle modification and optimizing healthcare resources through a multidisciplinary team approach constitute key

features of the journal. This journal is indexed on American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/integrated-blood-pressure-control-journal>