COPD management costs according to the frequency of COPD exacerbations in UK primary care

Yogesh Suresh Punekar¹

Amit Shukla²

Hana Müllerova³

¹Global Health Outcomes, GlaxoSmithKline R&D, Uxbridge, UK; ²Worldwide Epidemiology, GlaxoSmithKline R&D, Upper Providence, PA, USA; ³Worldwide Epidemiology, GlaxoSmithKline R&D, Uxbridge, UK

Background: The economic burden of chronic obstructive pulmonary disease (COPD) exacerbations is significant, but the impact of other sources on the overall cost of COPD management is largely unknown. We aimed to estimate overall costs for patients experiencing none, one, or two or more exacerbations per year in the UK.

Methods: A retrospective cohort of prevalent COPD patients was identified in the Clinical Practice Research Datalink UK database. Patients with information recorded for at least 12 months before and after cohort entry date were included (first prevalent COPD diagnosis confirmed by spirometry on/after April 1, 2009). Patients were categorized as having none, one, or two or more moderate-to-severe COPD exacerbations in the 12 months after cohort entry and further classified by the Global initiative for chronic Obstructive Lung Disease (GOLD) category of airflow obstruction and the Medical Research Council dyspnea scale. Study outcomes included counts of general practitioner interactions, moderate-severe COPD exacerbations, and non-COPD hospitalizations. Estimated resource use costs were calculated using National Health Service reference costs for 2010–2011.

Results: The cohort comprised 58,589 patients (mean age 69.5 years, mean dyspnea grade 2.5, females 46.6%, current smokers 33.1%). The average total annual per patient cost of COPD management, excluding medications, was £2,108 for all patients and £1,523, £2,405, and £3,396 for patients experiencing no, one, or two or more moderate-to-severe exacerbations, respectively. General practitioner interactions contributed most to these annual costs, accounting for £1,062 (69.7%), £1,313 (54.6%), and £1,592 (46.9%) in patients with no, one, or two or more moderate-to-severe exacerbations, respectively.

Conclusion: Disease management strategies focused on reducing costs in primary care may help reduce total COPD costs significantly.

Keywords: chronic obstructive pulmonary disease, frequent exacerbations, infrequent exacerbations, health resources, health care costs

Introduction

Chronic obstructive pulmonary disease (COPD) is a highly prevalent¹ and debilitating disease that has a significant impact on patient quality of life and the costs borne by the health care system.^{2,3} An exacerbation of COPD is the most common condition that requires hospital admission,⁴ and contributes substantially to the related economic impact.^{5–7} Therefore, COPD exacerbations have been considered the most important target for reducing costs. A study in the UK National Health Service (NHS) in 2001 showed that the total excess primary care cost associated with acute exacerbations of chronic bronchitis in the COPD-diagnosed population was £35.7 million. This included the excess cost of all prescription medications, general practitioner (GP)

Correspondence: Yogesh Suresh Punekar
Global Health Outcomes,
GlaxoSmithKline, Stockley Park,
Uxbridge UB11 1BT, UK
Tel +44 020 8990 4786
Fax +44 020 8990 3505
Email yogesh.q.punekar@gsk.com

consultations, hospital emergency room visits, and inpatient hospital episodes.⁸ Similarly, a retrospective analysis of data from a large national health plan database in the US showed that COPD-related mean annual costs were 56% higher for patients with two or more exacerbations compared with the overall population of COPD patients.⁸ This finding has also been reflected in other studies conducted across Europe and the US, suggesting that prevention of exacerbations requiring hospitalization is the foremost factor critical in the development of a cost-effective disease management strategy.^{5-7,10,11}

In contrast, symptomatic patients who are at a low risk of exacerbation represent an unexplored population with respect to health care costs. These patients continue to report symptoms, potentially consuming significant primary care resources. Therefore, it is important to assess the severity of these symptoms and their impact on health care resource utilization. The annual administration of the Medical Research Council (MRC) dyspnea scale in the UK primary care setting instituted by the UK NHS Quality Outcomes Framework (since April 2009) allows such quantitative assessment of symptoms of dyspnea.¹² A previous study using this dyspnea recording has found a high prevalence of dyspnea in the primary care COPD population in the UK.¹³

This study evaluated the primary care COPD population and its costs using the Clinical Practice Research Datalink (CPRD) database. The study objectives were to estimate resource use and the associated costs among patients classified by no, one, and two or more exacerbations and to compare resource use and costs between different levels of exacerbations across the Global initiative for chronic Obstructive Lung Disease (GOLD) stages of airflow obstruction and MRC dyspnea categories.

Materials and methods

Study design

This was a retrospective cohort study of COPD patients identified from primary care in the UK using the CPRD. CPRD is the NHS observational data and interventional research service, jointly funded by the NHS National Institute for Health Research and the Medicines and Healthcare products Regulatory Agency.¹⁴ In this study, we used the primary care electronic medical records database part of the CPRD, formerly known as the General Practice Research Database.¹⁴ About 660 general practices participate in the CPRD; the database contains data for about 11 million patients, mostly from England and Wales. The COPD cohort study design and analysis methods (protocol WEUSKOP5903), prior to the cost

estimation and modeling described herein, were approved by the CPRD Independent Scientific Advisory Committee.

Patients aged 40 years or older with one or more records for diagnosis of COPD (Table S1) on or after April 1, 2009 were included. Date of the first recorded COPD diagnosis on or after April 1, 2009 was taken as the cohort entry date. Patients were required to have at least a 12 month history before and after cohort entry and a record of forced expiratory volume in one second/forced vital capacity (FEV_1/FVC) <0.7 . The observation period consisted of a fixed period of 12 months of follow-up after cohort entry. The corresponding 12 month data prior to cohort entry have been referred to as the pre-observation period. Health care resource use during the observation and pre-observation periods was captured as moderate-to-severe exacerbations, GP interactions, and non-COPD hospitalizations. Exacerbations were further split into moderate and severe events. A moderate exacerbation was defined as an episode characterized by management with antibiotics commonly used in COPD exacerbations and oral corticosteroids (within 5 days of initiation of antibiotics), and/or a medical diagnosis of COPD exacerbation. A severe exacerbation was defined as an episode characterized by hospitalization for COPD. These were selected using an algorithm that identified hospital admissions of specific COPD cause or general admissions with a record of COPD diagnosis or medication up to 3 days before and 7 days after the index hospitalization date. The first event qualifying as an exacerbation was considered as the start date of the episode. Thereafter, a 14 day moving window¹⁵ was applied to identify at least 2 weeks of an exacerbation event-free period to ensure that a relapse was not considered a separate exacerbation. GP interactions were recorded and classified as administrative contact, surgery correspondence, visit to a surgery nurse, in-person surgery visit, out-of-office GP visit, or a GP home visit. Non-COPD hospitalizations were defined as hospital admission for specific non-COPD cause or general hospital admissions (without a specific cause) with no record of COPD diagnosis or medication recorded around this event.

Patients were classified by their moderate-to-severe exacerbation frequency (no, one, or two or more) during the observation period. Further, exacerbation frequency and medications in the pre-observation period and comorbidities at baseline were recorded. All costs were calculated using 2011 prices (except the medication costs, which were taken from 2012) expressed as mean annual costs per patient, and summarized (Table 1). For patients with no, one, and two or more exacerbations, baseline characteristics and costs estimated during the observation period were

Table 1 Unit costs estimated for the resource use categories

Resource use item	Unit cost	Reference
Moderate exacerbation	£85.29	NHS reference costs 2010–2011
Severe exacerbation	£1,263.76	NHS reference costs 2010–2011
Hospital episode	£1,366.88	PSSRU 2011
GP practice in-person visit	£36	PSSRU 2011, Page 149
GP practice nurse visit	£13	(Based on 15.5-minute consultation) × £51 per hour of face to face contact
GP home visit	£121	PSSRU 2011, Page 149
GP out-of-office visit	£121	Assumed to be the same as home visits
GP practice administrative contact	£22	PSSRU 2011, Page 149; considered equivalent to telephone consultation
GP practice correspondence	£3	Based on 30% of indirect contact of 15.5-minute consultation (0.3×15.5) × £39 per hour of non-face to face contact

Abbreviations: GP, general practitioner; NHS, National Health Service; PSSRU, Personal Social Services Research Unit.

compared using the chi-square test and one-way analysis of variance.

Bivariate relationships with exacerbation frequency categories were evaluated using the Cochran–Mantel–Haenszel test (for categorical or ordinal variables), polyserial correlations (for continuous variables), or negative binomial modeling with exacerbation frequency during the observation period as the predictor variable while controlling for age and gender to assess a relationship with the count data.

The cost of a severe exacerbation was estimated using the relevant COPD-related Healthcare Resource Groups (HRG) codes (DZ21A–K for short stay and long stay) further weighted by the reported annual number of episodes from the NHS reference costs.¹⁶ In addition, it was assumed that 67% of patients would arrive at the hospital by ambulance, based on a clinical audit of COPD exacerbations in 2008.¹⁷ Therefore, 67% of the HRG cost for ambulance transportation (code PS06B) was added to the weighted COPD HRG costs. Costs of a moderate exacerbation were compiled based on the resource use stated in the report by the GOLD Strategy Group¹⁸ and combined with cost data from the NHS reference costs,¹⁶ the Personal Social Services Research Unit (PSSRU) 2011,¹⁹ and the British National Formulary 65.²⁰ This included a GP consultation lasting 11.7 minutes, an accident and emergency visit with no admission (in 29% of cases), and a prescription of prednisolone (30 mg) and co-amoxiclav (500 mg).²¹ The cost of a non-COPD hospitalization was

estimated as a weighted average of short-stay and long-stay hospital episodes from PSSRU costs.¹⁹

The costs of a GP visit in surgery and a GP visit at home were estimated to be £36 and £121, respectively, based on PSSRU costs.¹⁹ The cost of an out-of-office GP visit was assumed to be equivalent to a GP home visit, and the cost of a GP administrative contact (£22) was assumed to be equivalent to a phone consultation at a GP surgery. The cost of a nurse visit at a GP practice was estimated to be £13 based on a typical 15.5-minute face to face consultation, and the cost of a GP correspondence was estimated to be £3 based on 30% indirect costs of a typical non-face to face consultation.¹⁹

These unit costs were then applied to the corresponding annual resource use counts to calculate the total annual COPD management costs for the entire cohort and for patients with no, one, and two or more exacerbations. Costs estimated during the observation period were further summarized by the GOLD stage of airflow obstruction²² and the level of dyspnea estimated using the MRC dyspnea scale²³ captured nearest to the cohort entry.

Results

Patient disposition and demographic characteristics

The final analytic COPD cohort consisted of 58,589 patients (Figure S1) with a mean follow-up of 13.5 years in CPRD. The cohort had a mean age of 69.5 years; 46.6% of the patients were female and 33.1% were current smokers (Table 2). The mean ($\pm$ standard deviation) FEV₁ percent predicted value at baseline was 60.0 ± 20.5 , with a majority of patients in the GOLD 2 category. The mean MRC grade at baseline was 2.5 ± 1.0 .

The cohort consisted of 31,049 (52.9%), 14,189 (24.2%), and 13,351 (22.7%) patients experiencing no, one, and two or more episodes of exacerbation during the observation period, respectively (Table 2). Patients with a higher exacerbation frequency had lower percent predicted FEV₁ ($P < 0.0001$), more dyspnea ($P < 0.0001$), and a higher comorbidity index ($P < 0.0001$) at or near baseline. The resources collected in the pre-observation and observation periods for each of the three groups are presented in Table 3.

Cost of COPD treatment

The mean total annual COPD management cost per patient was £2,108 in the observation period. Of these costs, £257 was accounted for by exacerbations, £637 was from non-COPD hospitalizations, and £1,213 was due to GP interactions. The corresponding total COPD management

Table 2 Patient disposition and demographic characteristics at the start of the observation period, split by frequency of moderate-to-severe exacerbations during observation period

Patient variable	Moderate-severe exacerbations: none	Moderate-severe exacerbations: one	Moderate-severe exacerbations: two or more	P-value across exacerbation categories
^a All COPD, n	31,049	14,189	13,351	
Female, n (%)	13,619 (43.86)	6,831 (48.14)	6,832 (51.17)	<0.0001
Age, mean (SD)	69.3 (10.46)	69.77 (10.32)	69.44 (10.17)	0.0180
Current smokers, n (%)	10,531 (33.92)	4,667 (32.89)	4,173 (31.26)	<0.0001
BMI, mean (SD)	26.98 (5.72)	27.04 (5.95)	26.82 (6.04)	0.0431
FEV ₁ %pred, mean (SD)	62.83 (19.94)	59.39 (20.05)	54.15 (20.74)	<0.0001
MRC grade, mean (SD)	2.28 (0.99)	2.51 (1.03)	2.83 (1.08)	<0.0001
Charlson score, mean (SD)	2.48 (1.63)	2.56 (1.66)	2.58 (1.66)	<0.0001

Note: ^aNumber of patients experiencing no, one, and two or more moderate-to-severe exacerbations during the 12 month observation period.

Abbreviations: %pred, percent predicted; BMI, body mass index; COPD, chronic obstructive pulmonary disease; FEV₁, forced expiratory volume at one second; MRC, Medical Research Council dyspnea scale; SD, standard deviation.

cost in the pre-observation period was £1,826, comprising £200 due to exacerbations, £572 due to non-COPD hospitalizations, and £1,054 due to GP interactions.

When split by the frequency of moderate-to-severe exacerbations during the observation period, the total annual per patient COPD costs were £1,523, £2,405, and £3,396

for patients with no, one, and two or more exacerbations, respectively. GP interactions contributed most to these annual costs, accounting for £1,062 (69.7%) in patients with no exacerbations, £1,313 (54.6%) in patients with one exacerbation, and £1,592 (46.9%) in patients with two or more exacerbations (Figure 1). For this cohort, the pre-observation mean

Table 3 Health care utilization during the 12 month observation period before and after cohort entry

Patient variable	Moderate-severe exacerbations: none n=31,049	Moderate-severe exacerbations: one n=14,189	Moderate-severe exacerbations: two or more n=13,351	P-value across the exacerbation categories [#]
Resource category 12 months prior to start of observation period, n (rate) ^a				
Moderate exacerbations	9,226 (0.30)	8,804 (0.62)	19,523 (1.46)	<0.0001/<0.0001
Severe exacerbations	2,278 (0.07)	1,753 (0.12)	2,709 (0.20)	0.2065/<0.0001
Non-COPD hospitalizations	10,347 (0.33)	6,470 (0.46)	7,682 (0.58)	<0.0001/<0.0001
GP				
GP practice in-person visit	309,543 (9.96)	162,821 (11.47)	173,822 (13.01)	<0.0001/<0.0001
GP practice nurse visit	106,730 (3.43)	58,332 (4.11)	67,137 (5.02)	<0.0001/<0.0001
GP home visit	5,643 (0.18)	3,680 (0.25)	4,908 (0.36)	0.1763/<0.0001
GP out-of-office visit	2,704 (0.08)	2,079 (0.14)	2,624 (0.19)	<0.0001/<0.0001
GP practice administrative contact	719,613 (23.17)	371,698 (26.19)	391,748 (29.34)	<0.0001/<0.0001
GP practice correspondence	27,843 (0.89)	15,764 (1.11)	16,048 (1.20)	<0.0001/<0.0001
Resource category during 12 month observation period, n (rate) ^a				
Moderate exacerbations	–	10,952 (0.77)	32,302 (2.42)	NA
Severe exacerbations	–	3,237 (0.23)	5,776 (0.43)	NA
Non-COPD hospitalizations	10,310 (0.33)	7,383 (0.52)	9,631 (0.72)	<0.0001/<0.0001
GP				
GP practice in-person visit	341,138 (10.98)	190,325 (13.41)	212,879 (15.94)	<0.0001/<0.0001
GP practice nurse visit	128,755 (4.14)	80,210 (5.65)	96,652 (7.23)	<0.0001/<0.0001
GP home visit	6,036 (0.19)	5,110 (0.36)	7,950 (0.59)	0.2986/<0.0001
GP out-of-office visit	2,542 (0.08)	2,568 (0.18)	4,068 (0.30)	<0.0001/<0.0001
GP practice administrative contact	786,501 (25.33)	423,124 (29.82)	4,57,316 (34.25)	<0.0001/<0.0001
GP practice correspondence	22,009 (0.70)	13,554 (0.95)	14,187 (1.06)	<0.0001/<0.0001

Notes: ^aObserved frequency/total number of patients; [#]negative binomial regression adjusted for age and sex was used to test any difference in event counts between patients with one and two or more exacerbations events compared with those with no exacerbation event during the 12 month follow-up (referent group). The first P-value refers to a comparison of those with one event vs no event; the second P-value refers to a comparison of those with two or more events vs those with no events.

Abbreviations: COPD, chronic obstructive pulmonary disease; GP, general practitioner; NA, not applicable for categories where no events in the referent group were recorded.

Figure 1 Costs of COPD treatment (per patient per year) before and after cohort entry date.

Note: ^aClassification based on moderate-to-severe exacerbation frequency in the follow-up period.

Abbreviations: COPD, chronic obstructive pulmonary disease; GP, general practitioner.

total annual per patient COPD costs were £1,523, £1,928, and £2,420, respectively.

The estimated annual COPD management costs (excluding non-exacerbation-related medication costs) during the observation period by GOLD stages of airflow obstruction are presented in Table 4 and by level of dyspnea in Table 5. The mean total annual COPD management costs were £1,767 for

patients with mild dyspnea (MRC grade 1 or 2) and £2,534 for patients with moderate-severe dyspnea.

Similar cost escalations with an increase in dyspnea were also observed in all cost components, including exacerbations (£182 versus £342), non-COPD hospitalizations (£504 versus £787), and GP interactions (£1,081 versus £1,404). This upward trend in total annual costs and cost

Table 4 COPD management costs split by exacerbation frequency during the observation period and further by GOLD stages of airflow obstruction

Exacerbation frequency	GOLD categories	No of patients	Cost per patient per year			
			Exacerbations	Non-COPD hospitalizations	GP surgery contact	Total
All patients	Stage 1	8,844	£202.40	£621.90	£1,159.76	£1,984.06
	Stage 2	29,832	£232.61	£635.89	£1,218.50	£2,087.00
	Stage 3	15,497	£319.65	£672.90	£1,297.33	£2,289.88
	Stage 4	3,377	£445.53	£775.20	£1,418.44	£2,639.17
None	Stage 1	5,475	£0.00	£502.88	£1,026.41	£1,529.29
	Stage 2	16,815	£0.00	£468.84	£1,068.41	£1,537.25
	Stage 3	7,013	£0.00	£428.50	£1,073.42	£1,501.92
	Stage 4	1,152	£0.00	£405.51	£1,100.32	£1,505.83
One	Stage 1	1,971	£378.76	£731.44	£1,286.98	£2,397.18
	Stage 2	7,324	£365.97	£738.93	£1,307.32	£2,412.22
	Stage 3	3,855	£347.98	£710.18	£1,331.90	£2,390.06
	Stage 4	814	£351.51	£682.12	£1,308.08	£2,341.71
Two or more	Stage 1	1,398	£753.96	£937.95	£1,507.45	£3,199.36
	Stage 2	5,693	£754.80	£1,001.51	£1,551.82	£3,308.13
	Stage 3	4,629	£789.53	£1,018.99	£1,614.04	£3,422.57
	Stage 4	1,411	£871.16	£1,137.25	£1,747.82	£3,756.23

Abbreviations: COPD, chronic obstructive pulmonary disease; GOLD, Global initiative for chronic Obstructive Lung Disease; GP, general practitioner.

Table 5 COPD management costs split by exacerbation frequency during the observation period and further by levels of dyspnea

Exacerbation frequency	MRC dyspnea scale	Patients (n)	Cost per patient per year			
			Exacerbations	Non-COPD hospitalizations	GP surgery contact	Total
All COPD patients	MRC 1	9,475	£147.73	£456.67	£992.66	£1,597.06
	MRC 2	19,863	£198.50	£527.08	£1,122.87	£1,848.45
	MRC 3	13,251	£289.63	£701.10	£1,307.06	£2,297.79
	MRC 4	7,440	£395.54	£859.85	£1,489.57	£2,744.96
	MRC 5	1,612	£549.09	£1,189.76	£1,840.23	£3,579.08
None	MRC 1	6,251	£0.00	£367.70	£899.43	£1,267.14
	MRC 2	11,552	£0.00	£417.81	£1,006.53	£1,424.34
	MRC 3	6,478	£0.00	£526.29	£1,163.58	£1,689.88
	MRC 4	2,959	£0.00	£592.94	£1,257.82	£1,850.76
	MRC 5	524	£0.00	£763.10	£1,472.06	£2,235.16
One	MRC 1	2,059	£339.52	£560.84	£1,120.17	£2,020.53
	MRC 2	4,765	£340.54	£618.77	£1,214.69	£2,174.00
	MRC 3	3,487	£368.61	£758.89	£1,334.02	£2,461.51
	MRC 4	1,876	£407.57	£911.00	£1,514.72	£2,833.29
	MRC 5	398	£455.75	£1,202.40	£1,789.27	£3,447.42
Two or more	MRC 1	1,165	£605.03	£752.17	£1,269.79	£2,477.37
	MRC 2	3,546	£657.12	£761.34	£1,380.15	£2,633.15
	MRC 3	3,286	£781.61	£987.24	£1,563.77	£3,102.86
	MRC 4	2,605	£842.87	£1,130.34	£1,738.48	£3,487.93
	MRC 5	690	£1,033.85	£1,517.05	£2,158.54	£4,340.16

Abbreviations: COPD, chronic obstructive pulmonary disease; GP, general practitioner; MRC, Medical Research Council dyspnea scale.

components was observed regardless of exacerbation frequency. This increase in total annual cost between patients with mild and moderate-severe dyspnea was £396 for patients with no exacerbations, £518 for patients with one exacerbation, and £862 for patients with two or more exacerbations in the observation period.

The mean total annual COPD management costs were £2,063 for mild-moderate patients (GOLD stages 1 and 2) and £2,351 for severe patients (GOLD stages 3 and 4). When split by exacerbation frequency, total costs decreased with increase in severity among patients with no exacerbations (£1,535 versus £1,502) and patients with one exacerbation (£2,409 versus £2,382). In patients with two or more exacerbations, total costs increased with COPD severity (£3,287 versus £3,499). Among the cost components, GP interactions showed an upward trend with increase in COPD severity, regardless of the exacerbation frequency.

A majority of exacerbation costs were associated with severe exacerbations. For the entire cohort, the mean annual per patient cost of moderate and severe exacerbations was £63 and £194 in the observation period and £55 and £145 in the pre-observation period, respectively. Corresponding costs in the observation period were £67 and £295 for patients with one exacerbation and £214 and £567 for patients with two or more exacerbations.

Discussion

This study explored health care resource use and associated costs of COPD patients in the CPRD database. We focused on differences in costs by exacerbation frequency, further split by the GOLD stages of airflow obstruction and the levels of dyspnea. Patients were stratified into three categories of no, one, and two or more moderate-to-severe exacerbations based on an analysis published in the literature.²⁴

Our results show that the majority of the health care costs in COPD patients are attributable to GP interactions. In patients with two or more (frequent) exacerbations, these accounted for 47% of total COPD costs whilst contributing 70% and 55% in patients with no and one exacerbation, respectively. A study by McGuire et al assessed the excess cost of chronic bronchitis and acute exacerbations of chronic bronchitis in the UK primary care setting.⁸ They estimated that the excess cost of GP visits attributable to chronic bronchitis and acute exacerbations of chronic bronchitis was £6.5 million in England and Wales in 1997–1998. Our estimate of £23.3 million due to in-person GP visits and £61.8 million due to GP interactions in the CPRD sample alone is significantly higher.

A study reported by Wouters estimated the annual direct cost of COPD patients to be \$1,254 in the UK in 2002, with mean annual GP visits per person being 6.6.²⁵ The estimated cost in our study was £1,826 in the pre-observation period

and £2,108 during the observation period. We estimated the mean annual GP practice visits to be 12.7 among the entire cohort. This suggests a substantial increase in health care use in the primary care setting over the last 10 years. The Wouters study also reported 14% for patients with hospitalizations and 12% for emergency room visits.²⁵ We found 9.6% and 21.8% of patients reporting at least one hospitalization for COPD and non-COPD causes, respectively.

A study by Schermer et al estimated the costs associated with exacerbations in a primary care setting in the Netherlands. Their study reported annual costs of exacerbations as £40, £53, £61, and £92 among patients with no, mild, moderate, and severe airflow obstruction, respectively, in 2001.²⁶ The corresponding estimates in our study were £202, £233, £320, and £446, respectively. However, it is important to note that the severity classification used in the study by Schermer et al was based on the European Respiratory Society recommendation,²⁷ while ours was based on GOLD.²³ A study by Andersson et al showed that exacerbations account for 35%–45% of the total per capita health care costs for COPD.²⁸ In our study, which was mainly based on the primary care COPD population, exacerbations accounted for less than 25% of the costs, even for frequent exacerbators, highlighting a relatively small contribution of exacerbation costs to the overall COPD management costs. One possible explanation for the higher proportion of GP costs is the holistic assessment of GP interactions in our study. We included all forms of GP interactions, including non-GP-led interactions within and outside the surgery. Most other studies published in the literature have focused on surgery-based GP-led visits.

In our study, total COPD management costs and its cost components increased with increasing level of disease severity. This was more evident with increasing levels of dyspnea than with increasing levels of airflow obstruction, regardless of the exacerbation frequency. In part, this could be explained by more severe COPD patients having higher comorbidities and thus requiring more GP interactions and non-COPD hospitalization. This hypothesis, however, was not tested in the current study. A similar study by Tabberer et al estimated the incremental annual cost of moderate-to-severe dyspnea in general practice to be in excess of £250 per patient.²⁹ Our estimates of the corresponding mean increase in annual COPD management costs with each level increase in dyspnea were £242, £357, and £466 for patients with no, one, and two or more exacerbations, respectively. Overall, the resource use costs increased with higher levels of exacerbation frequency and dyspnea.

Therefore, efforts to better manage patients in primary care with an aim to reduce exacerbations and symptoms that drive GP visits will help reduce overall COPD management costs significantly.

Our study has several limitations. We excluded the cost of non-exacerbation-related medications from our cost calculations. A previous study has shown medication cost to be a major contributor to overall health care costs.⁸ This may significantly underestimate the annual COPD costs per patient in our analysis. We also used tariff-based cost estimates applicable across England and Wales. Whilst this improves the generalizability of our findings, the actual costs in local settings may differ from national estimates, thereby limiting their applicability to local health economies. Further, patients were included only if their information was available for at least 24 months. Such a condition of a longer stay in a cohort may introduce survival bias, given that patients with a higher disease severity and a higher risk of health care events do not survive for the full follow-up. In addition, the quality of spirometry data is unknown. One study indicated the possibility of a variable quality of spirometry measurement in a primary care population largely not included in the CPRD database.³⁰

Conclusion

Estimated COPD management costs increased with exacerbation frequency and level of dyspnea based on data from 12 months of follow-up in an observational cohort study of patients managed in primary care. Interactions with GP surgery accounted for a substantial proportion of costs regardless of the exacerbation frequency. Further, the costs of GP interactions, exacerbations, and non-COPD hospitalizations increased with increasing levels of dyspnea. Therefore, in a primary care setting, better disease management focused on holistic assessment and reduction of symptoms is required. This is particularly applicable to patients with a low risk of exacerbations, who continue to utilize a significant proportion of health care resources. Future strategies should follow published guidelines such as GOLD to include systematic measurement of symptoms and health status, allowing physicians to identify patients with unmet need and treat them accordingly to improve individual outcomes and the collective burden.

Disclosure

Editorial support, in the form of development, assembling tables and figures, collating author comments, copyediting, fact checking, and referencing, was performed

by Dr Annirudha Chillar from Cactus Communications Inc, and was funded by GlaxoSmithKline. The authors have no other conflicts of interest to report.

References

1. Buist AS, McBurnie MA, Vollmer WM, et al; BOLD Collaborative Research Group. International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet*. 2007;370(9589):741–750.
2. Effing T, Kerstjens H, van der Valk P, Zielhuis G, van der Palen J. (Cost)-effectiveness of self-treatment of exacerbations on the severity of exacerbations in patients with COPD: the COPE II study. *Thorax*. 2009;64(11):956–962.
3. Nurmatov U, Buckingham S, Kendall M, et al. Effectiveness of holistic interventions for people with severe chronic obstructive pulmonary disease: systematic review of controlled clinical trials. *PLoS One*. 2012;7(10):e46433.
4. Chetty M, MacKenzie M, Douglas G, Currie GP. Immediate and early discharge for patients with exacerbations of chronic obstructive pulmonary disease: is there a role in “real life”? *Int J Chron Obstruct Pulmon Dis*. 2006;1(4):401–407.
5. Jahnz-Rozyk K, Targowski T, From S. Costs of exacerbations of chronic obstructive pulmonary disease in primary and secondary care in 2007 – results of multicenter Polish study. *Pol Merkuriusz Lekarski*. 2009;26(153):208–214. Polish.
6. Nielsen R, Johannessen A, Benediktsdottir B, et al. Present and future costs of COPD in Iceland and Norway: results from the BOLD study. *Eur Respir J*. 2009;34(4):850–857.
7. Ornek T, Tor M, Altin R, et al. Clinical factors affecting the direct cost of patients hospitalized with acute exacerbation of chronic obstructive pulmonary disease. *Int J Med Sci*. 2012;9(4):285–290.
8. McGuire A, Irwin DE, Fenn P, et al. The excess cost of acute exacerbations of chronic bronchitis in patients aged 45 and older in England and Wales. *Value Health*. 2001;4(5):370–375.
9. Pasquale MK, Sun SX, Song F, Hartnett HJ, Stenkowski SA. Impact of exacerbations on health care cost and resource utilization in chronic obstructive pulmonary disease patients with chronic bronchitis from a predominantly Medicare population. *Int J Chron Obstruct Pulmon Dis*. 2012;7:757–764.
10. Dalal AA, Christensen L, Liu F, Riedel AA. Direct costs of chronic obstructive pulmonary disease among managed care patients. *Int J Chron Obstruct Pulmon Dis*. 2010;5:341–349.
11. Miravittles M, Murio C, Guerrero T, Gisbert R. Pharmacoeconomic evaluation of acute exacerbations of chronic bronchitis and COPD. *Chest*. 2002;121(5):1449–1455.
12. Fletcher CM, Elmes PC, Fairbairn AS, Wood CH. The significance of respiratory symptoms and the diagnosis of chronic bronchitis in a working population. *BMJ*. 1959;2(5147):257–266.
13. Mullerova H, Lu C, Tabberer M. Disease burden of dyspnea in a primary care COPD population. *Am J Respir Crit Care Med*. 2012;185:A1518.
14. Clinical Practice Research Datalink. Available from: <http://www.cprd.com/home/>. Accessed October 4, 2012.
15. Leidy NK, Wilcox TK, Jones PW, Roberts L, Powers JH, Sethi S. Standardizing measurement of chronic obstructive pulmonary disease exacerbations. Reliability and validity of a patient-reported diary. *Am J Respir Crit Care Med*. 2011;183(3):323–329.
16. Gov.UK. Policy paper. NHS reference costs 2010–2011. Available from: <https://www.gov.uk/government/publications/2010-11-reference-costs-publication>. Accessed October 4, 2012.
17. Buckingham RJ, Lowe D, Pursey NA, Roberts CM, Stone RA. Report of the National Chronic Obstructive Pulmonary Disease Audit 2008: clinical audit of COPD exacerbations admitted to acute NHS units across the UK: Royal College of Physicians, British Thoracic Society, British Lung Foundation, 2008. Available from: <http://www.brit-thoracic.org.uk/Portals/0/Clinical%20Information/COPD/NCROP/NCROPClinicalAudit.pdf>. Accessed November 17, 2013.
18. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management and prevention of COPD (2010). Available from: http://www.goldcopd.org/uploads/users/files/GOLDReport_April112011.pdf. Accessed November 17, 2012.
19. Personal Social Services Research Unit. Unit Costs of Health and Social Care 2011. Available from: <http://www.pssru.ac.uk/project-pages/unit-costs/2011/index.php>. Accessed October 4, 2012.
20. British National Formulary 65, 2012. Available from: <http://bnf.org/bnf/index.htm>. Accessed October 4, 2012.
21. Hertel N, Kotchie RW, Samyshkin Y, Radford M, Humphreys S, Jameson K. Cost-effectiveness of available treatment options for patients suffering from severe COPD in the UK: a fully incremental analysis. *Int J Chron Obstruct Pulmon Dis*. 2012;7:183–199.
22. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease (2006). Available from: http://www.goldcopd.org/uploads/users/files/GOLDReport2006_0122.pdf. Accessed October 4, 2012.
23. National Institute for Health and Care Excellence. Medical Research Council dyspnea scale. Available from: http://www.nice.org.uk/usingguidance/commissioningguides/pulmonaryrehabilitationsserviceforpatientswithcopd/mrc_dyspnoea_scale.jsp. Accessed June 20, 2011.
24. Hurst JR, Vestbo J, Anzueto A, et al. Susceptibility to exacerbation in chronic obstructive pulmonary disease. *N Engl J Med*. 2010;363(12):1128–1138.
25. Wouters EF. Economic analysis of the Confronting COPD survey: an overview of results. *Respir Med*. 2003;97 Suppl C:S3–S14.
26. Schermer TR, Saris CG, van den Bosch WJ, et al. Exacerbations and associated health care cost in patients with COPD in general practice. *Monaldi Arch Chest Dis*. 2006;65(3):133–140.
27. Siafakas NM, Vermeire P, Pride NB, et al. Optimal assessment and management of chronic obstructive pulmonary disease (COPD). The European Respiratory Society Task Force. *Eur Respir J*. 1995;8(8):1398–1420.
28. Andersson F, Borg S, Jansson SA, et al. The costs of exacerbations in chronic obstructive pulmonary disease (COPD). *Respir Med*. 2002;96(9):700–708.
29. Tabberer M, Lu C, Müllerova H. Incremental burden of dyspnea in COPD to a primary care COPD population. Presented at the Annual Congress of the American Thoracic Society, San Francisco, CA, USA, May 18–23, 2012.
30. Strong M, South G, Carlisle R. The UK Quality and Outcomes Framework pay-for-performance scheme and spirometry: rewarding quality or just quantity? A cross-sectional study in Rotherham, UK. *BMC Health Serv Res*. 2009;9:108.

Supplementary materials

Table S1 COPD management costs according to the frequency of COPD exacerbations in UK primary care

ReadCode description	ReadCode
COPD ReadCodes used to extract the study COPD cohort from the electronic medical records	
Acute exacerbation of COAD	H312200
Admitted COPD emergency	8H2R.00
Airways obstruction irreversible	663K.00
Chronic bronchitis with emphysema	49I E
COAD follow-up	66YL.I2
COPD	5199GP
COPD follow-up	66YL.I1
COPD self-management plan given	66YI.00
Centrilobular emphysema	H322.00
COPD with acute exacerbation, unspecified	H3yI.00
Chronic bullous emphysema	H320.00
Chronic bullous emphysema NOS	H320z00
COPD with acute lower respiratory infection	H3y0.00
COAD	H3...I1
COAD NOS	H3z..00
COPD	H3...00
COPD NOS	H3z..I1
COPD annual review	66YM.00
COPD clinical management plan	8CRI.00
COPD disturbs sleep	66Yg.00
COPD does not disturb sleep	66Yh.00
COPD follow-up	66YL.00
COPD monitor phone invite	9OI4.00
COPD monitoring	66YB.00
COPD monitoring 1st letter	9OI0.00
COPD monitoring 2nd letter	9OI1.00
COPD monitoring 3rd letter	9OI2.00
COPD monitoring administration	9OI..00
COPD monitoring by doctor	66YT.00
COPD monitoring by nurse	66YS.00
COPD monitoring due	66YD.00
COPD monitoring verb invite	9OI3.00
Emphysema pulmonary	492
Exacerbation COAD	5199GE
Emphysema	H32..00
Emphysema NOS	H32z.00
Emphysematous bronchitis	H312100
Health education – COPD	679V.00
Mild COPD	H36..00
Moderate COPD	H37..00
COAD	5199G

(Continued)

Table S1 (Continued)

Obstructive chronic bronchitis NOS	H312z00
Radiological emphysema	9906E
Severe COPD	H38..00
Very severe COPD	H39..00

Abbreviations: COPD, chronic obstructive pulmonary disease; COAD, chronic obstructive airways disease; NOS, not otherwise specified.

Figure S1 Study flow chart.

Abbreviations: COPD, chronic obstructive pulmonary disease; FEV₁, forced expiratory volume at one second; FVC, forced vital capacity.

International Journal of COPD

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-copd-journal>

Dovepress

This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.