Evaluation of the stability and antimicrobial activity of an ethanolic extract of *Libidibia ferrea*

Raquel de Oliveira Marreiro¹
 Maria Fulgência Costa Lima Bandeira¹
 Tatiane Pereira de Souza²
 Mailza Costa de Almeida¹
 Katiana Bendaham¹
 Gisely Naura Venâncio¹
 Isis Costa Rodrigues²
 Cristiane Nagai Coelho¹
 Patrícia Sâmea Lêdo Lima Milério¹
 Glauber Palma de Oliveira¹
 Nikeila Chacon de Oliveira Conde¹

¹School of Dentistry, Federal University of Amazonas, Manaus, Amazonas; ²School of Pharmacy, Federal University of Amazonas, Manaus, Amazonas, Brazil

Abstract: Biofilm is a dense, whitish, noncalcified aggregate of bacteria, with desquamated epithelial cells and food debris creating conditions for an imbalance of resident oral microflora and favoring the destruction of hard and soft tissues by development of caries and gingivitis. The aim of this study was to obtain and characterize an extract of *Libidibia ferrea*, ex *Caesalpinia ferrea* L. and to evaluate its feasibility for formulation as a mouthwash, according to current legislation. For this purpose, pH, sedimentation, density, and stability were evaluated, along with microbiological testing of the extract. The microbiological test was used to verify the presence of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, fungi, yeasts, coliforms, and minimum inhibitory concentrations of *Streptococcus mutans* and *Streptococcus oralis* strains. Characterization, microbiological evaluation, and minimum inhibitory concentration results were tabulated and described using descriptive statistics. The *L. ferrea* extract showed stable characteristics, product quality, and antibacterial activity against the microorganisms tested irrespective of experimental time intervals. According to these results, it can be concluded that formulation of a mouthwash containing *L. ferrea* extract to control biofilm is feasible, but further studies are needed.

Keywords: biofilm, extract, quality control

Introduction

Dental plaque is defined as a community of oral bacteria on a tooth surface in which microorganisms are found embedded in the extracellular polymeric substance and in intimate communication with each other via several different pathways, such as auto/coaggregation, metabolic communication, quorum sensing, and competent stimulation peptides.¹

Dental caries and periodontal disease are initiated by accumulation of plaque on the soft tissue and on the tooth surface, leading to tooth loss and gum problems.² A consequence of such biofilm growth is increased resistance to chemical antimicrobial agents and antibiotics,^{3,4} which has profound implications in the oral environment and in medicine.

Biofilm can be controlled by mechanical procedures or by chemical intervention. Mechanical control is the most effective method, but is not always done properly. Thus, antimicrobial substances have been used for chemical control of dental biofilm, working as aids to mechanical procedures and for specific conditions.⁵⁻⁷ Given the incidence of oral disease, increased bacterial resistance to antibiotics, the adverse affects of some antibacterial agents currently used in dentistry, and financial considerations in developing countries, there is a need for alternative prevention and treatment options

Correspondence: Gisely Naura Venâncio
 School of Dentistry, Federal University of Amazonas, Waldemar Pedrosa Avenue, 1539, Downtown, 69025-050, Manaus, Amazonas, Brazil
 Tel +92 8839 0595
 Email ginaura@gmail.com

that are safe, effective, and economical. Hence, the search for alternative products continues, and natural phytochemicals isolated from plants used as traditional medicines are considered good alternatives.⁸ The search for therapeutic applications of medicinal plants and their derivatives has been growing in recent years throughout the world. Several studies have been carried out in order to evaluate new therapeutic use of plants. The discovery of new antimicrobial components is relevant, given that bacterial and fungal infections of the oral cavity are a relatively common problem. For example, *Candida albicans* is opportunistic yeast commonly identified in denture stomatitis and other forms of oral candidosis, and *Streptococcus mutans* is associated with dental biofilm and one of the most important microorganisms involved in dental caries.⁹

Studies have been conducted to assess the activity of plants of the Amazon flora against the microorganisms present in dental biofilm.^{10–12} The significant global growth of phytotherapy as part of preventive and curative programs has encouraged assessment of the activity of a number of plant extracts. The ancient practice of members of a community using plants to re-establish health, common to all people but almost forgotten for decades, has become increasingly more intense in the civilized world in recent years.¹³ In the Amazon region of Brazil, the fruit of *Libidibia ferrea* is widely used as an antimicrobial and healing medicine in many situations, including oral infections.¹⁴ Based on these observations, the aim of this study was to evaluate the antibacterial activity of *L. ferrea* extract against microorganisms present in biofilm, in the hope of making it a future complementary method for chemical control of biofilm formation.

Materials and methods

The dry fruit of *L. ferrea* (228022-INPA), a species of the family Leguminosae caesalpinaceae, known as jucá, jucaína, and pau-ferro (Brazilian ironwood), was collected at the National Institute of Amazon Research and processed at the School of Pharmaceutical Sciences, Federal University of Amazonas. The alcoholic extract was prepared by maceration at room temperature for 72 hours using ethanol as the solvent and 10% (w/v) of raw vegetable material. After extraction, the extract was filtered and taken to the vacuum evaporator (RV10 basic rotary evaporator, IKA®-Werke GmbH and Co, Staufen, Germany) to remove the solvent. The residue was then stored in previously sterilized and dried amber bottles with lids.

Characterization of hydroalcoholic extract

Characterization of the extract was performed using three different storage environments, ie, an oven (37°C), room temperature (24°C–26°C), and a refrigerator ($\pm 4.8^\circ\text{C}$) at experimental time intervals of 0, 7, 30, 45, and 90 days.

Aliquots of the extract were withdrawn aseptically with the aid of an automatic pipette and the following were assessed: pH, using a pH meter (pHtek, PHS-3B®, Campinas, São Paulo, Brazil); sedimentation, by rotational speed in a centrifuge at 1,500 and 3,000 rpm for 5 minutes (Nova técnica®, Nt810, Piracicaba, São Paulo, Brazil); organoleptic characteristics (color, odor, consistency, and brightness), by visual-tactile assessment; and density, using the dry pycnometer method.¹⁵

Microbiological evaluation

The extract was analyzed for the existence of microorganisms to verify the presence or absence of *Staphylococcus aureus* and *Pseudomonas aeruginosa* in Plate Count Agar, Baird-Parker, and Cetrimide media and fungi and yeasts using Sabouraud Agar. The presence or absence of growth was analyzed by counting colony-forming units. Cultivation was done in triplicate for each experimental time interval and condition.

Escherichia coli broth was used at a proportion of 10 mL for each tube with a lid (15 × 150 mm) in inverted Durham tubes for total coliform count. For each type of culture medium used, a white group (without extract) was prepared and the whole experimental procedure was conducted in a class II biological safety cabinet.

Assessment of MIC by microdilution

The minimum inhibitory concentration (MIC) was determined using a previously reported method,¹⁶ and 100 μL of liquid culture medium (Mueller-Hinton) was distributed into sterile 96-well microplates. In row A of the plate, 100 μL of the extract solution was added and analyzed at a concentration of 100 μL per mL. Using a 12-channel microdilution pipette, serial dilutions were made by transferring 100 μL from row A to row B and so on until row G of the plate. The last row (row H) was left without any extract (test solution) and used as a control for bacterial growth. Next, 25 μL of the standardized bacterial suspension and indicator solution were inoculated into each plate well. The plate was sealed and incubated at 37°C for 24 hours and reading was performed visually. The MIC was defined as the lowest concentration of the extract capable of inhibiting bacterial growth, ie,

preventing color change. Five repetitions of the tests were performed. Two plates were used for the experiment against two types of bacterium (*S. mutans* and *S. oralis*). In each plate, two groups of *L. ferrea* at different experimental time intervals were tested (extract 1 at 20 days and extract 2 at 140 days).

The results for characterization, microbiological assessment, and MIC were tabulated and submitted to descriptive analysis. For assessment of pH, the non-parametric Kruskal-Wallis test and Epi-Info version 7 software for Windows, developed and distributed by the Centers for Disease Control and Prevention (<http://www.cdc.gov/epiinfo>), were used at a 5% level of significance.

Results

Characterization of the *L. ferrea* extract for pH, density, sedimentation, and stability was performed at baseline (T0) and at 7 (T7), 30 (T30), 45 (T45), and 90 (T90) days in three storage locations, ie, a refrigerator, an oven, and at room temperature of 24°C–26°C (Table 1).

As shown in Table 2, the mean pH values in the oven were 4.32 (T0), 4.21 (T7), 4.26 (T30), 3.86 (45), and 3.42 (T90). In the refrigerator, pH decreased to 4.32 (T0), 4.25 (T7), 4.60 (T30), 4.30 (T45), and 3.95 (T90). At room temperature, pH decreased similarly to 4.32 (T0), 4.25 (T7), 4.53 (T30), 4.17 (T45), and 3.91 (T90). Analysis of variance (5% significance level) and Tukey's test for multiple comparisons were used, and showed a statistically significant difference in pH in the room temperature group at T45 in comparison with the other groups (refrigerator and oven).

No sediment was found in the sedimentation test at 1,500 rpm. However, small amounts of sediment were found at T45 using the same test at 3,000 rpm in the oven (1 mm) and at room temperature (0.2 mm). At T90, sediment was found on storage in the refrigerator and oven (0.5 mm).

There was little variation in the density values, and those found for oven storage were 0.886 (T0), 0.883 (T7), 0.893 (T30), 0.889 (T45), and 0.892 (T90). In the refrigerator, the corresponding values were 0.886 (T0), 0.883 (T7), 0.893 (T30), 0.889 (T45), and 0.891 (T90). At room temperature,

density was 0.886 (T0), 0.883 (T7), 0.889 (T30), 0.890 (T45), and 0.891 (T90). The differences in density between the initial and final time intervals increased by 0.006 g/L in the oven and by 0.005 g/L in the refrigerator and at room temperature.

The organoleptic properties of color, brightness, odor, and consistency did not change, regardless of variations in location and storage time.

No contamination of microorganisms was found in the *L. ferrea* extract in the microbiological tests performed to verify the presence or absence of bacteria, fungi, yeasts, and total coliforms.

Tests for antibacterial activity of the extract (24 g of *L. ferrea* per 100 mL of 80% ethanol) against *S. mutans* and *S. oralis* showed that extract 1 (initial time interval) had bactericidal activity in seven dilutions (100 µL, 50 µL, 25 µL, 12.5 µL, 6.25 µL, 3.125 µL, and 1.5625 µL). At T2, the extract (final time interval, 120 days) showed bactericidal activity at a concentration of 3.125 µL.

Discussion

The desire for new products offering better biocompatibility and pharmacological activity with less toxicity has led to an increase in research on natural products for use in dentistry. Natural products with antimicrobial activity may be important therapeutic agents to prevent and treat caries, periodontal disease, and oral candidiasis.^{8,12,17}

Several studies have been conducted in dentistry to evaluate plant species, including *Echinacea purpurea*, *Copaifera multijuga*, *Lippia sidoides*, *Stryphnodendron barbatimao*, and *Pothomorphe umbellata*.^{4,10–12,18–21} The present study examined the characteristics of a hydroalcoholic extract of the plant species *L. ferrea*, aiming to obtain more information for future formulation of a natural mouthwash applicable in dentistry.

The results showed no statistically significant difference in pH within the 45-day experimental time interval between the room temperature and refrigerator groups, but did show a significant difference in the oven group. At the end of the experiment, the pH of the extract was below 5.5, which is considered a critical pH value for dissolution of enamel, and thus must be modified in other formulations.

In the sedimentation test, at all time intervals and at different rotations, the sediments (0.2 mm, 0.5 mm, and 1 mm) did not interfere with antibacterial activity. The observed variation in density was small and independent of experimental time interval or storage location.

Analyzing the different density values at the initial and final time intervals, an increase of 0.006 g/L was found in

Table 1 Mean pH values for the extract according to local storage conditions and trial duration

Storage location	Period (days)				
	0	7	30	45	90
Oven	4.32	4.21	4.26	3.86	3.42
Refrigerator	4.32	4.25	4.60	4.30	3.95
Room temperature	4.32	4.25	4.53	4.17	3.91

Table 2 Mean of pH difference according to types of storage

Period (days)	Groups									P*
	Room temperature (n=3)			Oven (n=3)			Refrigerator (n=3)			
	Mean	Q ₁	Q ₂	Mean	Q ₁	Q ₂	Mean	Q ₁	Q ₂	
0–7	–0.08	–0.21	0.08	–0.11	–0.19	–0.02	–0.05	–0.21	0.05	0.967
0–30	0.23	0.07	0.33	–0.07	–0.14	0.04	0.24	0.23	0.38	0.055
0–45	–0.16 ^a	–0.23	–0.06	–0.44 ^b	–0.56	–0.37	–0.01 ^a	–0.13	0.08	0.039 ^c
0–90	–0.43	–0.49	–0.30	–0.89	–0.98	–0.81	–0.37	–0.45	–0.27	0.061

Notes: *Kruskal-Wallis test; ^{a,b,c}statistically significant difference at a level of 5%.

Abbreviations: Q₁, first quartile; Q₂, second quartile.

oven storage conditions, while in the refrigerator and at room temperature, the increase was 0.005 g/L. These variations were due to formation of substances with higher molecular weight, loss of water, or volatility of the extract; however, such modifications were not sufficient to interfere in the evaluation of the other tests.

Identification and purity of the plant, as well as evaluation of its active ingredients, are essential when trying to identify good quality products.¹¹ In Brazil, despite wide consumption of products derived from plants, the products commercialized and consumed are not subjected to specific control.

In this study, the aim of the microbiological tests was to investigate growth of microorganisms (bacteria, fungi, yeasts, and total coliforms) in the extract. Under the conditions and types of culture medium tested, there was no indication of contamination by bacteria, fungi, yeast, or coliforms at any time interval or storage condition.

Fungal contamination of herbal drugs can lead to alteration and/or destruction of the active ingredients and loss of safety and efficacy; in addition, toxic substances (mycotoxins) may be produced, making these products unfit for consumption regardless of the level of contamination.^{22–24}

With regard to assessment of the antibacterial activity of the *L. ferrea* extract against the *S. mutans* and *S. oralis* strains using the microdilution test on microplates to determine MIC, the tests showed that extract 1 (initial time interval) had bactericidal activity in the seven dilutions (100 µL, 50 µL, 25 µL, 12.5 µL, 6.25 µL, 3.125 µL, and 1.5625 µL) established in the methodology. Extract 2 (final time interval, 120 days) showed bactericidal activity at a concentration of 3.125 µL. The results establish the antibacterial efficacy of the extract against the microorganisms tested after 4 months of storage. A previous study of the antimicrobial activity of *L. ferrea* against pathogens of the oral cavity is consistent with the present results.¹⁴

Different studies assessing the antimicrobial activity of natural products against various bacterial strains have

shown positive results for such substances with regard to inhibition of growth or development of bacteria, and this has also been shown in dentistry, suggesting that plant extracts could be an alternative therapeutic option for infections of the oral cavity, such as denture stomatitis, dental caries, and periodontitis.^{8,17,25} Based on the results, it is concluded that the *L. ferrea* extract has antibacterial activity against the microorganisms tested and satisfactory stability and quality, enabling the formulation of a mouthwash using this extract to control dental biofilm.

Acknowledgment

This study was supported by the Foundation for Research Support of the Amazonas State (FAPEAM).

Disclosure

The authors have no conflicts of interest to report in this work.

References

1. Rickard AH, Bachrach G, Davies D. Cell-cell communication in oral microbial communities. In: Rogers AH, editor. *Molecular Oral Microbiology*. Norfolk, UK: Caister Academic Press; 2008.
2. Cury JA. Controle químico da placa dental [Oral health promotion]. In: ABOPREV: Kriger L, editor. *Promoção de saúde bucal*. 3rd ed. São Paulo, Brazil: Artes Médicas; 2003.
3. Jyothi KS, Seshagiri M. In-vitro activity of saponins of *Bauhinia purpurea*, *Madhuca longifolia*, *Celastrus paniculatus* and *Semecarpus anacardium* on selected oral pathogens. *J Dent (Tehran)*. 2012;9:216–223.
4. Vahabi S, Fekrazad R, Ayremlou S, Taheri S, Zangeneh N. The effect of antimicrobial photodynamic therapy with radachlorin and toluidine blue on *Streptococcus mutans*: an in vitro study. *J Dent (Tehran)*. 2011;8:48–54.
5. Bugno A, Buzzo AA, Nakamura CT, Pereira TC, Matos D, Pinto TJA. Avaliação da contaminação microbiana em drogas vegetais [Evaluation of microbial contamination in herbal drugs]. *Rev Ciênc Farm Básica Apl*. 2005;41:491–497.
6. Fejerskov O, Kidd E. Cárie dentária – a doença e seu tratamento clínico [Dental caries – the disease and clinical treatment]. 1st ed. São Paulo, Brazil: Santos; 2005.
7. Moreira ACA, Santos TAM, Carneiro MC, Porto MR. Atividade de um enxaguatório bucal com clorexidina a 0,12 por cento sobre a microbiota sacarolítica da saliva [Activity of a mouthwash with chlorhexidine 0.12% on saccharotic saliva microbiotic]. *Rev Ciênc Méd Biol*. 2008;7:260–272.

8. Palombo EA. Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases. *Evid Based Complement Alternat Med*. 2011;2011:680354.
9. Pereira EM, Gomes RT, Freire NR, Aguiar EG, Brandão Md, Santos VR. *In vitro* antimicrobial activity of Brazilian medicinal plant extracts against pathogenic microorganisms of interest to dentistry. *Planta Med*. 2011;77:401–404.
10. Bandeira MFCL, Oliveira MRB, Pizzolito AC, Benattineto C, Jorge-Neto J. Study preliminary pharmacology of the *Copaifera multijuga* (copaiba oil). *J Bras Clin Estet Odontol*. 1999;3:39–41. Portuguese.
11. Marques AA, Sponchiado EC Jr, Garcia LF, Garrido AD, França SC, Lia RC. Morphological analysis of tissue reaction caused by a new endodontic paste in subcutaneous tissue of rats. *J Conserv Dent*. 2011;14:309–313.
12. Paixão CCB. Uso de plantas medicinais em pacientes portadores de afecções bucais [Medicinal plants in oral diseases]. *Odontol Clin – Cient*. 2002;1:1–4.
13. Veiga VF, Zunino L, Calixto JB, Patitucci ML, Pinto AC. Phytochemical and antiedematogenic studies of commercial copaiba oils available in Brazil. *Phytother Res*. 2001;15:476–480.
14. Sampaio FC, Pereira Mdo S, Dias CS, Costa VC, Conde NC, Buzalaf MA. *In vitro* antimicrobial activity of *Caesalpinia ferrea* Martius fruits against oral pathogens. *J Ethnopharmacol*. 2009;124:289–294.
15. *Farmacopéia Brasileira*. National Health Surveillance Agency. 4th ed. São Paulo, Brazil: 1988.
16. Telles MAS, Mosca A. Evaluation of the use of microtitre plate technique to determine the minimum inhibitory concentration of isoniazid in *Mycobacterium tuberculosis* strains. *Rev Inst Adolfo Lutz*. 2000;59:15–19.
17. Castilho AR, Murata RM, Pardi V. Natural products in Dentistry. *Health J*. 2007;11–19.
18. Couto GBL. Biocompatibility of the *Lippia sidoides* Cham (Verbenaceae) hidro-alcoholic essence. *Rev Cons Reg Odontol Pernambuco*. 2000;3:83–90. Portuguese.
19. Feres M, Figueiredo LC, Barreto IM, Coelho MH, Araújo MW, Cortelli SC. *In vitro* antimicrobial activity of plant extracts and propolis in saliva samples of healthy and periodontally-involved subjects. *J Int Acad Periodontol*. 2005;7:90–96.
20. Melo Júnior EJM, Raposo MJ, Santana AEG, Lisboa Neto JA, Diniz MFA. Estudo de plantas medicinais com atividade antimicrobiana sobre microrganismos presentes na alveolite [Study of medicinal plants with antimicrobial activity against microorganisms in alveolitis]. *Rev ABO Nacional*. 2000;8:4–9.
21. Vendola MCC. *Equinacea purpurea* na odontologia [*Equinacea purpurea* in Dentistry]. *Técs Estéts*. 2004;1:3–12.
22. Correa Júnior C, Ming LC, Scheffer MC. Cultivo de plantas medicinais e aromáticas e condimentares [Cultivation of medicinal and aromatic plants and herbs]. Jaboticabal: FUNEP, 1994.
23. Matos FJA. Plantas medicinais: guia de seleção e emprego de plantas usadas em Fitoterapia no Nordeste do Brasil [Medicinal plants: guide selection and use of plants used in herbal medicine in Northeast Brazil]. Fortaleza: Imprensa Universitária-UFC, 2000.
24. Sponchiado Júnior EC, Pereira JV, Pereira JO, França SC. *Pothomorphe umbellata* (Caapeba). *Rev Fitos*. 2007;3:31–37.
25. Maekawa LE, Valera MC, Oliveira LD, Carvalho CA, Camargo CH, Jorge AO. Effect of *Zingiber officinale* and propolis on microorganisms and endotoxins in root canals. *J Appl Oral Sci*. 2013;21:25–31.

Clinical, Cosmetic and Investigational Dentistry

Publish your work in this journal

Clinical, Cosmetic and Investigational Dentistry is an international, peer-reviewed, open access, online journal focusing on the latest clinical and experimental research in dentistry with specific emphasis on cosmetic interventions. Innovative developments in dental materials, techniques and devices that improve outcomes and patient satisfaction

Submit your manuscript here: <http://www.dovepress.com/clinical-cosmetic-and-investigational-dentistry-journal>

Dovepress

and preference will be highlighted. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.