Is combination of neutrophil to lymphocyte ratio and platelet lymphocyte ratio a useful predictor of postoperative survival in patients with esophageal squamous cell carcinoma?

Alpaslan Tanoglu¹
Ergenekon Karagoz²
Nurettin Yiyit³
Ufuk Berber⁴

¹Department of Gastroenterology,

²Department of Infectious

Diseases and Clinical Microbiology,

³Department of Thoracic Surgery,

⁴Department of Pathology, GATA
Haydarpasa Training Hospital, Uskudar,
Turkey

Dear editor

We read with interest the recent article entitled “Combination of neutrophil to lymphocyte ratio and platelet lymphocyte ratio is a useful predictor of postoperative survival in patients with esophageal squamous cell carcinoma” by Feng et al.¹ In their study, authors aimed to investigate the usefulness of a novel inflammation-based prognostic system, using the combination of neutrophil lymphocyte ratio (NLR) and platelet lymphocyte ratio (PLR), for predicting survival in patients with esophageal squamous cell carcinoma (ESCC). Finally, they concluded that combination of NLR and PLR is a useful predictor of postoperative survival in patients with ESCC and combination of these parameters is superior to NLR or PLR as a predictive factor in patients with ESCC. We would like to thank the authors for their contribution.

PLR has been recently suggested to be a marker of thrombotic and inflammatory condition, mainly in patients with malignancies.^{2,3} NLR is a readily available and inexpensive laboratory marker which is used to assess systemic inflammation. In literature, it was shown that diabetes mellitus, thyroid functional abnormalities, essential hypertension, valvular heart diseases, acute coronary syndromes, renal and/or hepatic failure, metabolic syndrome, and many inflammatory diseases may potentially affect the NLR.⁴⁻⁷ Thus, it would be more relevant if Feng et al had mentioned these NLR-affecting factors while evaluating the predictive role of NLR in postoperative survival of patients with ESCC. Besides, medication may alter NLR and/or PLR, so it would have been useful if the patients were described in greater detail in terms of antibiotic, anti-diabetic, anti-hypertensive drug use and/or other medications. In addition, it would also have been better if the authors indicated the elapsed time between taking the blood samples and measuring NLR and PLR, since waiting period prior to analysis may affect these parameters.

We believe that the findings of Feng et al¹ will lead to further studies concerning the predictive role of NLR and PLR for postoperative survival of patients with ESCC. But, it should be clearly kept in mind that NLR or PLR itself alone without other variables may not secure true information about postoperative survival of patients with ESCC. Finally we concluded that these parameters should be evaluated with other variables as mentioned above.

Correspondence: Alpaslan Tanoglu
Department of Gastroenterology,
GATA Haydarpasa Training Hospital,
34668, Uskudar, Turkey
Tel +90 216 542 2020
Fax +90 216 542 2007
Email alpaslantanoglu@yahoo.com

Disclosure

The authors have no conflicts of interest in this correspondence.

References

1. Feng JF, Huang Y, Liu JS. Combination of neutrophil lymphocyte ratio and platelet lymphocyte ratio is a useful predictor of postoperative survival in patients with esophageal squamous cell carcinoma. *Onco Targets Ther*. 2013;7;6:1605–1612.
2. Wang D, Yang JX, Cao DY, et al. Preoperative neutrophil-lymphocyte and platelet-lymphocyte ratios as independent predictors of cervical stromal involvement in surgically treated endometrioid adenocarcinoma. *OncoTargets Ther*. 2013;6:211–216.
3. Smith RA, Ghaneh P, Sutton R, Raraty M, Campbell F, Neoptolemos JP. Prognosis of resected ampullary adenocarcinoma by preoperative serum CA19-9 levels and platelet-lymphocyte ratio. *J Gastrointest Surg*. 2008;12(8):1422–1428.
4. Alkhouri N, Morris-Stiff G, Campbell C, et al. Neutrophil to lymphocyte ratio: A new marker for predicting steatohepatitis and fibrosis in patients with nonalcoholic fatty liver disease. *Liver Int*. 2012;32(2):297–302.
5. Balta S, Cakar M, Demirkol S, Arslan Z, Akhan M. Higher neutrophil to lymphocyte ratio in patients with metabolic syndrome. *Clin Appl Thromb Hemost*. 2013;19(5):579.
6. Karaman M, Balta S, Ay SA, et al. The Comparative Effects of Valsartan and Amlodipine on vWf Levels and N/L Ratio in Patients with Newly Diagnosed Hypertension. *Clin Exp Hypertens*. 2013;35(7):516–522.
7. Stotz M, Gerger A, Eisner F, et al. Increased neutrophil-lymphocyte ratio is a poor prognostic factor in patients with primary operable and inoperable pancreatic cancer. *Br J Cancer*. 2013;109:416–421.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.