A systematic review of interventions conducted in clinical or community settings to improve dual-task postural control in older adults

Maayan Agmon¹

Basia Belza²

Huong Q Nguyen^{2,3}

Rebecca G Logsdon²

Valerie E Kelly⁴

¹The Cheryl Spencer Department of Nursing, Faculty of Social Welfare and Health Studies, University of Haifa, Israel; ²School of Nursing, University of Washington, Seattle, WA, USA; ³Department of Research and Evaluation, Kaiser Permanente, CA, USA; ⁴School of Medicine, University of Washington, Seattle, WA, USA

Background: Injury due to falls is a major problem among older adults. Decrements in dual-task postural control performance (simultaneously performing two tasks, at least one of which requires postural control) have been associated with an increased risk of falling. Evidence-based interventions that can be used in clinical or community settings to improve dual-task postural control may help to reduce this risk.

Purpose: The aims of this systematic review are: 1) to identify clinical or community-based interventions that improved dual-task postural control among older adults; and 2) to identify the key elements of those interventions.

Data sources: Studies were obtained from a search conducted through October 2013 of the following electronic databases: PubMed, CINAHL, PsycINFO, and Web of Science.

Study selection: Randomized and nonrandomized controlled studies examining the effects of interventions aimed at improving dual-task postural control among community-dwelling older adults were selected.

Data extraction: All studies were evaluated based on methodological quality. Intervention characteristics including study purpose, study design, and sample size were identified, and effects of dual-task interventions on various postural control and cognitive outcomes were noted.

Data synthesis: Twenty-two studies fulfilled the selection criteria and were summarized in this review to identify characteristics of successful interventions.

Limitations: The ability to synthesize data was limited by the heterogeneity in participant characteristics, study designs, and outcome measures.

Conclusion: Dual-task postural control can be modified by specific training. There was little evidence that single-task training transferred to dual-task postural control performance. Further investigation of dual-task training using standardized outcome measurements is needed.

Keywords: physical therapy, balance, walking, motor learning, fall prevention

Introduction

In 2020, one out of five people in western countries will be 70 years of age or older.¹ Healthy aging is accompanied by changes in sensory and cognitive domains that may lead to balance and gait impairments.^{2,3} Balance and gait impairments, in turn, contribute to recurrent falls, which are related to increased mortality and morbidity.⁴ Thirty percent of adults over age 65, and 50% of those over age 85, are likely to have at least one fall.^{5,6} Consequently, finding effective ways to decrease falls in the elderly may reduce disability and increase life expectancy.⁷

Although falls are multifactorial,⁸ impaired postural control is one important factor contributing to falls. Postural control is defined as the ability to control the body's position in space for the purposes of stability and orientation,⁹ and is critical

Correspondence: Maayan Agmon
University of Haifa, Abba Hushi I,
Haifa, Israel
Tel +972 54 900 1609
Fax +972 4 828 8017
Email agmon.mn@gmail.com

during standing balance and walking tasks. Much research has focused on the interplay between postural control and cognition,¹⁰ using dual-task postural control paradigms to examine this relationship.¹¹ Dual-task performance refers to the ability to conduct two tasks simultaneously, with dual-task postural control referring to situations when at least one of the tasks involves postural control, such as walking while talking on the phone or while holding a tray. Evaluating dual-task performance is a complex process as it involves the evaluation of each task performed independently as well as in combination. One way of analyzing the performance is by calculating the dual-task cost, defined as the decline in dual-task compared to single-task performance of a task.^{12–14}

Changes associated with aging may lead to deterioration with the performance of each individual task as well as with the dual-task combination. For example, the gait pattern is affected by age, with reduced stride length and gait speed as well as increased lateral sway and stride to stride variability among older adults.¹⁵ Executive function is a set of cognitive skills required in order to plan, monitor, and conduct goal-directed complex actions,¹⁶ and an important aspect of executive function that tends to deteriorate with aging is the ability to divide or switch attention between the different tasks.^{17,18} This ability is critical for dual-task performance. In older adults, dual-task postural control deficits have been associated with declines in cognitive function^{19,20} and increased risk for falls.^{21–23}

Recent studies have demonstrated the potential for modification of dual-task performance among the elderly.^{24,25} Specifically, the ability to divide attention between two tasks in order to conduct them simultaneously is modifiable following training.^{26,27} An important aspect of effective motor learning is training specificity, which refers to training a specific task through repetitive exercises in order to achieve improvements in that task.²⁸ Training in dual-task performance is more complicated than training a discrete movement under single-task condition,²⁹ and the level of specificity required to improve dual-task performance is still unknown. Moreover, dual-task postural control performance is influenced by the types of tasks, their difficulty, and the outcome measured.^{30,31}

There are a growing number of interventions aimed at improving dual-task postural control in healthy older adults. Wollesen and Voelcker-Rehage²⁴ performed a systemic review of the dual-task literature to examine the effects of specific versus general training and task combination on dual-task performance. They concluded that dual-task training is more effective than single-task for improving dual-task

standing performance, whereas both dual-task and single-task training improved dual-task walking. The current review extends this effort by examining how the application of motor learning principles, such as training specificity, setting, dose, duration, and intensity, may impact the efficacy of dual-task interventions. A better understanding of the effective elements of previous training trials can inform future dual-task interventions designed to improve mobility and reduce fall risk in older adults. Thus, the aims of this systematic review are to: 1) examine the effectiveness of different interventions on dual-task postural control among healthy older adults; and 2) identify key elements of training protocols that effectively improve dual-task postural control in older adults.

Methods

Data sources and searches

A systematic search was performed of the following computerized electronic databases: PubMed (January 1966 through October 2013), CINAHL (January 1982 through October 2013), PsycINFO (January 1900 through October 2013), PEDro (January 1929 through October 2013), and Web of Science (January 1900 through October 2013). Search terms included combinations of the following key words: “dual-task”; “older adults” or “elderly”; “treatment” or “intervention” or “therapy” or “rehabilitation”; “gait” or “balance” or “postural control”. References found by a manual search in identified articles were also reviewed and included as appropriate.

Inclusion criteria and study selection

To be included in this systematic review, a study had to meet the following criteria: 1) participants defined as healthy adults aged 60 years or older; 2) interventions were conducted in a community or clinical setting; 3) interventions required a minimum of 180 minutes of training over at least 3 total days; 4) dual-task postural control was measured as an outcome; and 5) the publication was written in English. Exclusion criteria included participants with a specific neurologic disorder, such as Parkinson’s disease, Alzheimer’s disease, dementia, or stroke.

Two reviewers (MA, VEK) screened the abstract search results and decided independently, based on inclusion and exclusion criteria, which studies to include. Results were compared and, when reviewer decisions differed, the full article was reviewed and evaluated to obtain agreement.

Data extraction and quality assessment

Reviewers, who were not blinded to the author or the journal, assessed the quality of each included study in terms

of the grade of recommendation and the level of evidence provided using the scoring protocol developed by Portney and Watkins.³² This scale includes ten levels of evidence divided into four levels of recommendations. The highest level of evidence is a meta-analysis and the lowest level is expert opinion.

Studies were summarized according to the following characteristics: methodological quality and level of evidence; study design; sample size; sample characteristics (age and sex); key characteristics of the training protocols (training specificity, content, setting, intensity); assessment time points; outcome measures (for postural control task and concurrent cognitive or motor task); and results. Data synthesis using a meta-analysis was not possible because of the variety of study designs, methodologies, and outcomes measured.

Results

The literature search yielded 162 publications that were screened, with 26 publications reviewed in full. Twenty-two publications met the inclusion criteria and were included in this review (Figure 1). Publications were excluded based on the following criteria: 1) not intervention trials (93 publications); 2) population was not appropriate (eg, included younger adults or older adults with a neurologic disorder; 43 publications); 3) did not measure dual-task performance as an outcome (3 publications); or 4) not written in English (1 publication).

Methodological quality and level of evidence

Twenty-two publications were included in this review, with two publications from the same intervention trial (Table 1).

Figure 1 Flowchart of systematic literature search.

Table I Level of evidence (n=22)

Study	Level of evidence ³²	Design
Hiyamizu et al, 2012 ³³	1b	RCT
Li et al, 2010 ³⁴	1b	RCT
Melzer et al, 2013 ³⁵	1b	RCT
Mozolic et al, 2011 ³⁶	1b	RCT
Silsupadol et al, 2009A ²⁶	1b	RCT
Silsupadol et al, 2009B ²⁷	1b	RCT
Trombetti et al, 2011 ³⁷	1b	RCT
Yamada et al, 2011 ³⁸	1b	RCT
Donath et al, 2013 ³⁹	2b	Preliminary RCT
Granacher et al, 2010 ⁴⁰	2b	Preliminary RCT
Hall et al, 2009 ⁴¹	2b	Preliminary RCT
Pichierri et al, 2012 ⁴²	2b	Preliminary RCT
Plummer-D'Amato et al, 2012 ⁴³	2b	Preliminary RCT
Uemura et al, 2012 ⁴⁴	2b	Preliminary RCT
Verghese et al, 2010 ⁴⁵	2b	Preliminary RCT
You et al, 2009 ⁴⁶	2b	Preliminary RCT
Agmon et al, 2012 ⁴⁷	3	Uncontrolled pretest to posttest
Bisson et al, 2007 ⁴⁸	3	Controlled pretest to posttest
Lajoie et al, 2004 ⁴⁹	3	Controlled pretest to posttest
Melzer et al, 2009 ⁵⁰	3	Case control
Toulotte et al, 2006 ⁵¹	3	Uncontrolled pretest to posttest
Silsupadol et al, 2006 ⁵²	4	Case series

Notes: Levels of evidence: 1a, Systematic review of RCTs; 1b, Individual RCT with narrow confidence interval; 2a, Systematic review of cohort studies; 2b, Individual cohort study or low quality RCT; 3, Individual case-control or pretest-posttest study; 4, Case series or poor quality cohort and case-control studies; 5, Expert opinion.

Abbreviation: RCT, randomized clinical trial.

In terms of level of evidence, eight were considered to be level 1b (randomized clinical trials with narrow confidence intervals) and eight were considered to be 2b (low quality randomized clinical trials). These sixteen studies^{26,27,33–46} had some methodological weaknesses; only one study included an intention-to-treat analysis,³⁷ and only four (two from the same trial) incorporated blinded assessors.^{26,27,43,45} The other six were classified as level 3 (case-control or cohort)^{47–51} and level 4 (case series).⁵²

Sample characteristics

Studies were included in this review only if the participants were adults aged 60 years or older. Sample size ranged from three (one per group)⁵² to 134.³⁷ The participants were predominantly female.

Sample characteristics varied across studies. Physical and cognitive functioning and the tests used to assess these characteristics varied. Most studies evaluated cognitive status using the mini-mental state examination with

inclusion criteria typically based on scores of 24 out of 30 points.^{26,27,41,46,48,50–52} Physical status was defined using performance-based tests, such as the Berg Balance Scale,^{26,27,35} Tinetti Test,³⁷ Mini-Balance Evaluation Systems Test (Mini-BEST),⁴⁷ or the Dynamic Gait Index,⁴¹ as well as self-reported walking abilities.^{26,27,35,38,40,43–45,47–52} Fall history was specified only in four studies.^{41,46,51,52} Living situation also varied across studies, though most studies involved participants who lived independently in the community (Table 2).

Training parameters of interventions

The training protocols incorporated in these studies varied with respect to training specificity (Table 3). Both single-task and dual-task approaches were used, and dual-task training incorporated various task combinations. Of the studies incorporating dual-task interventions,^{26,27,33–38,43–46,52} eleven included postural control tasks as one of the tasks,^{26,27,33,35–38,43,44,46,52} while two trained dual-task performance on two cognitive tasks.^{34,45} The other nine studies^{39–42,47–51} assessed the effects of single-task postural control training on dual-task postural control.

The studies that trained dual-task postural control used different combinations of tasks, with different levels of difficulty in both the cognitive and the postural tasks. For example, Trombetti et al³⁷ combined walking as the postural control task with a variety of motor (eg, handling musical instruments) and cognitive (eg, responding to changes in the beat of the music) tasks. Plummer-D'Amato et al⁴³ trained three different postural control domains (balance, gait, and agility) in combination with four different cognitive tasks (random number generation, word association, backward recitation of words or number sequences, and working memory).

Because dual-task performance can be modified by focusing attention on one task or another, the effect of dual-task training may be influenced by the instructions provided.⁵³ In this review, only the three studies conducted by Silsupadol et al^{26,27,52} specifically examined the impact of different instructions on dual-task postural control. These studies compared the effect of two sets of instructions – variable priority instructions and fixed priority instructions – on dual-task performance following the same training. Variable priority instructions required the participant to focus on one task at a time (either the postural or the cognitive task) while the fixed priority instructions required the participant to focus on both the postural and cognitive tasks at all times.

Interventions were conducted either one-on-one^{26,27,36,45,46,48,49,52} or in group settings (four to 24 people per group),^{33–35,37–44,47,50,51} with varying levels of supervision.

Table 2 Studies included in the review (n=22)

Study	Sample size, n (sex, F/M)	Group (sample size, n): mean (SD) age, years	Sample characteristics		Cognitive	Living situation*	Fall history
			Physical				
Hiyamizu et al 2012 ³³	36 (26/10)	Intervention (17): 72.0 (5.1) Control (19): 71.2 (4.4)	Not specified	Not specified	Not specified	Community-dwelling	With or without risk of falls
Li et al, 2010 ³⁴	20 (13/7)	Intervention (10): 74.6 (5.7) Control (10): 77.7 (7.1)	Healthy older adults without balance problem		Not specified	Community-dwelling	Without
Melzer et al, 2013 ³⁵	66 (49/17)	Intervention (33): 78.3 (6.2) Control (33): 75.7 (6.6)	BBS >45 Independently ambulatory: cane acceptable, not walker		MMSE >24	Community-dwelling and in senior living facility	Not specified
Mozolic et al, 2011 ³⁶	66 (35/31)	Intervention (33): 69.4 (3.2) Control (33): 69.4 (2.5)	Not specified		MMSE ≥5th percentile for participant age and education level	Community-dwelling	Not specified
Silsupadol et al, 2009A ²⁶	21 (17/4)	Single-task (7): 74.7 (7.8) DT fixed priority (8): 74.3 (6.1) DT variable priority (6): 76.0 (4.6)	Balance impairment defined as BBS ≤52 and/or self-selected gait speed ≤1.1 m/s		MMSE >24	Community-dwelling	With or without
Silsupadol et al, 2009B ²⁷	21 (17/4)	Single-task (7): 74.7 (7.8) DT fixed priority (8): 74.3 (6.1) DT variable priority (6): 76.0 (4.6)	Balance impairment defined as BBS ≤52 and/or self-selected gait speed ≤1.1 m/s		MMSE >24	Community-dwelling	With or without
Trombetti et al, 2011 ³⁷	134 (129/5)	Early intervention (66): 75.0 (8.0) Delayed intervention (68): 76.0 (6.0)	Increased risk of falling defined by meeting at least one of the following criteria: 1) one or more self-reported falls after age of 65 years; 2) Tinetti Test >2 of 7; 3) one or two criteria of physical frailty	Not specified	Not specified	Community-dwelling	At risk of fall
Yamada et al, 2011 ³⁸	84 (65/19)	Intervention (41): 83.0 (6.7) Control (43): 82.9 (5.5)	Ability to walk independently		Rapid Dementia Screening Test ≤4	Community-dwelling	Not specified
Donath et al, 2013 ³⁹	18 (9/9)	Intervention (9): 71.4 (4.7) Control (9): 73.1 (6.7)	At least moderately active		Not specified	Community-dwelling	Not specified
Granacher et al, 2010 ⁴⁰	20 (14/6)	Intervention (11): 71.9 (4.8) Control (9): 74.9 (6.3)	Able to walk without an assistive device		Not specified	Independent, community-dwelling	Not specified
Hall et al, 2009 ⁴¹	15 (10/5)	Intervention (8) Control (7)	Dynamic Gait Index score ≤19		MMSE ≥24	Community-dwelling	With a history of falls or at risk of falls
Pichierri et al, 2012 ⁴²	15 (9/6)	All participants: 72.2 (7.7); mean and SD not stated by group Intervention (9): 83.6 (3.4) Control (6): 86.2 (4.8)	Able to stand upright for at least 5 minutes		MMSE >22	Care homes	Not specified
Plummer-D'Amato et al, 2012 ⁴³	17 (16/1)	Dual-Task (10): 76.6 (5.6) Single-Task (7): 76.6 (6.0)	Able to walk independently at least 0.5 m/s without an assistive device		Able to follow three steps command	Community-dwelling	Without
Uemura et al, 2012 ⁴⁴	15 (12/3)	Intervention (8): 82.4 (5.9) Control (7): 82.5 (6.8)	Able to walk independently or with cane, no regular exercise in the previous 12 months		Not specified	Community-dwelling	Not specified
Verghese et al, 2010 ⁴⁵	24 (15/9)	Intervention (12): 77.4 (7.0) Control (12): 79.9 (7.5)	Sedentary with gait velocity <1.0 m/s		MMSE >25	Community-dwelling	Not specified

(Continued)

Table 2 (Continued)

Study	Sample size, n (sex, F/M)	Group (sample size, n): mean (SD) age, years	Sample characteristics		Cognitive	Living situation*	Fall history
			Physical	Independently ambulatory			
You et al, 2009 ⁴⁶	13 (11/2)	Intervention (8) Control (5) All participants: 68.3 (6.5); mean and SD not stated by group	Independently ambulatory		MMSE ≥ 24	Independent, community-dwelling	At least one fall in the past year
Agmon et al, 2011 ⁴⁷	30 (27/3)	Intervention (30): 74.5 (7.9)	Able to walk 10 meters independently		Not specified	Community-dwelling	Not specified
Bisson et al, 2007 ⁴⁸	24 (14/10)	Virtual reality (12): 74.4 (3.7) Bio feedback (12): 74.4 (4.9)	Able to walk without an assistive device		MMSE ≥ 20	Community-dwelling	Without
Lajoie et al, 2004 ⁴⁹	24 (20/4)	Intervention (12): 70.3 Control (12): 71.4	Able to stand and walk without an aid		Not specified	Community-dwelling and residential care facilities	Not specified
Melzer et al, 2009 ⁵⁰	48 (Not specified)	SD not stated by group Reporting exercise (24): 8.1 (6.19) Reporting inactive (24): 78.9 (5.1)	Able to walk with no aids		MMSE > 24	Independent, in retirement homes	Not specified
Toulotte et al, 2006 ⁵¹	16 (16/0)	With history of falls (8): 71.1 (5.0) Without history of falls (8): 68.4 (4.5)	Able to walk without an assistive device		MMSE ≥ 24	Not specified	Half with a history of fall within 2 years, half without
Sisupadol et al, 2006 ⁵²	3 (2/1)	Single-task (1): 82 Dual-task fixed priority (1): 90 Dual-task variable priority (1): 93	Able to walk 9 meters without the assistance of another person		MMSE ≥ 24	Independent	With

Notes: *The papers defined the living situation of study's participants differently. Here we used their definition.

Abbreviations: BBS, Berg Balance Scale; F, female; M, male; MMSE, mini-mental state exam; m/s, meters per second; SD, standard deviation.

There is no indication that one approach was more effective than the other.

Outcome measures

Balance and walking under both single-task and dual-task conditions was assessed using a variety of measures (Table 4). Therefore, there was no common set of standardized measures that could be used to compare changes in dual-task postural control across the studies (Table 2). Measures of single-task balance and walking included the Berg Balance Scale, the Dynamic Gait Index, the Timed Up and Go test, postural sway, and various gait parameters (speed, gait stability, center of mass or center of pressure, and variability) assessed during both simple and complex walking tasks. In addition, a number of studies incorporated measures of balance confidence or self-efficacy (Activities-specific Balance Confidence Scale, Falls Efficacy Scale) or function (Late Life Function and Disability Index) to examine the effects of training protocols on balance self-efficacy and functioning in daily life.

Measures of dual-task balance and walking included postural sway and gait parameters (speed, stability, variability). For dual-task balance and walking, performance on a variety of concurrent cognitive and motor tasks was also assessed. Examples of motor tasks included separating two linked rings or throwing and catching a ball. Examples of cognitive tasks included arithmetic tasks (eg, serial-3 subtractions), working memory tasks (eg, n-back test), or choice reaction time tasks (eg, auditory Stroop test). Dual-task performance was also evaluated using the dual-task cost calculation. In the studies that trained dual-task postural control, some assessed the efficacy of the intervention using trained task combinations⁴⁰ while others measured at least one novel task.^{26,27,52}

Dual-task performance changes

From the 22 publications included in this review, 18 demonstrated improvement in some aspect of dual-task performance^{26,27,33–40,42,44–49,51,52} whereas four did not.^{41,43,50,54} Of those showing improvement, three studies^{26,27,52} demonstrated improvement for both the postural control task and the concurrent cognitive or motor task. Seven studies demonstrated improvements in the dual-task cost for either postural control tasks or cognitive tasks, but not both.^{37,40,46–49,51} The other eight did not measure dual-task cost and demonstrated improvement on only one aspect of the tasks combination.^{33–36,38,39,42,44}

Table 3 Study characteristics (n=22)

Study	Protocol		Frequency		Duration		Setting	Content (by group)
	Time (minutes)							
Hiyamizu et al, 2012 ³³	60		Two times weekly		12 weeks		Group setting	1) Intervention: calculations, visual searches, and verbal fluency tasks were performed simultaneously during the balance training 2) Control: strength, balance, and walking training
Li et al, 2010 ³⁴	60		Five sessions at least 2 days apart		Approximately 5 weeks		Group setting (four to six people)	1) Intervention: computerized cognitive dual-task training: visual discrimination tasks 2) Control: no intervention
Melzer et al, 2013 ³⁵	60		Two times weekly		24 weeks		Group setting	1) Intervention: balance exercise in five different levels including perturbation and dual-task exercise 2) Control: no intervention
Mozolic et al, 2011 ³⁶	60		One time weekly		8 weeks		Individual training	1) Intervention: cognitive training focused on visual and auditory selective attention 2) Control: education lectures
Silsupadol et al, 2009A ²⁶	45		Three times weekly		4 weeks		One on one	1) Single-task balance training 2) Dual-task balance training with fixed priority instructions
Silsupadol et al, 2009B ²⁷	45		Three times weekly		4 weeks		One on one	3) Dual-task balance training with variable priority instructions 1) Single task balance training 2) Dual-task balance training with fixed priority instructions
Trombetti et al, 2011 ³⁷	60		One time weekly		25 weeks		Group setting	3) Dual-task balance training with variable priority instructions 1) Multitask exercises based on Jaques-Dalcroze music education (eg, walking in time to music, responding to changes in music, walk and turn, exaggerated upper body movements)
Yamada et al, 2011 ³⁸	20		Two times weekly		24 weeks		Group setting	1) Intervention: DVD seated based exercise, 15 minutes regular exercise, and 5 minutes dual-task verbal fluency while sitting and stepping 2) Control: no intervention
Donath et al, 2013 ³⁹	60		Two times weekly		6 weeks		Group setting	1) Intervention: specific progressive fall training includes martial arts and different falls on different mats 2) Control: three educational lessons on fall prevention knowledge
Granacher et al, 2010 ⁴⁰	60		Three times weekly		6 weeks		Group setting	1) Balance training (standing on unstable surfaces, single leg stance with increased flexion, catching and throwing a ball) Intervention: Tai Chi
Hall et al, 2009 ⁴¹	90 60 control	Tai Chi	Two times weekly		12 weeks		Group setting	2) Control: health-related topics
Pichierri et al, 2012 ⁴²	60		Two times weekly		12 weeks		Group setting	1) Progressive resistance training, progressive postural balance training, progressive dance video gaming
Plummer-D'Amato et al, 2012 ⁴³	45		One time weekly		4 weeks		Group setting	1) Intervention: balance, gait, and agility. The balance and gait activity combined with simultaneously cognitive tasks (number generation, word association, backward recitation, and working memory) 2) Control: balance, gait, and agility

(Continued)

Table 3 (Continued)

Study	Protocol		Setting		Content (by group)
	Time (minutes)	Frequency	Duration		
Uemura et al, 2012 ⁴⁴	35	One time weekly	24 weeks	Group setting for 30 minutes plus 5 minutes dual-task training	The intervention and control received 30 minutes of regular exercises. At the end of each session, each group received 5 minutes of different dual-task exercises: 1) Intervention: dual-task switch exercises 2) Control: steady state gait under dual-task
Verghese et al, 2010 ⁴⁵	24	Three times weekly at least 1 day apart	8 weeks	Individual training	1) Intervention: computerized cognitive training: auditory and cross modality tasks aimed at training attention and executive function 2) Control: received information about the importance of physical activity 1) Walking with cognitive task (memorizing and computing tasks) 2) Walking while listening to music 1) Enhance Fitness training
You et al, 2009 ⁴⁶	30	Five times weekly	18 total sessions over 6 weeks	One on one	1) Virtual reality ("juggling" a virtual ball) 2) Visual biofeedback (COP position)
Agmon et al, 2012 ⁴⁷	60	Three times weekly	6 weeks	Group setting	1) Computerized feedback and postural training 1) General training (Tai Chi, Feldenkreis, walking)
Bisson et al, 2007 ⁴⁸	30	Two times weekly	10 weeks	One on one	1) Exercise program: strengthening, flexibility, static balance, dynamic balance 1) Single task balance training
Lajoie et al, 2004 ⁴⁹	60	Two times weekly	8 week	One on one	2) Dual-task balance training with fixed priority instructions
Melzer et al, 2009 ⁵⁰	60	Three times weekly	3 years	Group setting	3) Dual-task balance training with variable priority instructions
Toulotte et al, 2006 ⁵¹	60	Two times weekly	3 months	Group setting	
Silsupadol et al, 2006 ⁵²	45	Three times weekly	4 weeks	One on one	

Abbreviation: COP, center of pressure.

Table 4 Study outcomes (n=22)

Study	Assessment points	Dual-task performance measures	Results	Dual-task improvement
Hiyamizu et al, 2012 ³³	1) Pretest 2) Posttest	1) Sway length of COG while performing Stroop test 2) The rate of Stroop task (accurate number/total numbers × 100) while maintaining standing position on a force plate Each physical outcome was tested concurrently with n-back working memory task: 1) Single support standing balance with eyes open and closed (four trials each) and involved standing on the dominant leg for 10 seconds per trial 2) Double-support standing balance under three conditions: stable platform, visual surround sway, and platform sway 3) Sit to stand test 4) Gait speed Voluntary step execution test under dual-task	1) Nonsignificant improvement 2) Significant improvement in the intervention group 1) Significant improvement	Yes, cognitive under dual-task
Li et al, 2010 ³⁴	1) Pretest 2) Posttest			Yes, motor under dual-task
Melzer et al, 2013 ³⁵	1) Pretest 2) Posttest		2) Significant improvement 3) Nonsignificant improvement 4) Nonsignificant improvement Significant improvement	Yes, motor
Mozolic et al, 2011 ³⁶	1) Pretest 2) Posttest	Walking while reciting words, measured number of recited words	Number of words recited significantly improved Dual-task cost did not improve	Yes, cognitive
Silsupadol et al, 2009A ²⁶	1) Pretest 2) Posttest	1) Narrow-base walking with serial-3 subtractions (trained dual-task condition): a) Walking: COM-AJC b) Serial-3 subtractions: response rate and number of errors	a) Improved for all groups with greatest improvement for those trained with variable priority instructions b) Response rate improved only in the VP group: number of errors did not differ between groups	Yes, both cognitive and walking tasks (for the trained task only) under dual-task
Silsupadol et al, 2009B ²⁷	1) Pretest 2) After 2 weeks 3) Posttest 4) 12 weeks after the intervention	2) Obstacle crossing with auditory Stroop task (novel dual-task condition): a) Walking: COM-AJC b) Auditory Stroop task: verbal reaction time Gait speed (while responding to addition/subtraction questions)	a) Did not improve for any group b) Did not improve for any group Improved for the trained dual-task groups (variable priority and fixed priority) but not for the control group (single-task)	Yes, motor under dual-task
Trombetti et al, 2011 ³⁷	1) Pretest 2) Posttest 3) 12 months	Walking in self-selected, slow and fast speeds while counting backward by one from 50: 1) Gait speed: gait velocity, stride length, cadence 2) Dynamic balance: double support %, step width 3) Gait variability: stride time variability, stride length variability	1) Stride length improved in the intervention group after 6 months 2) No significant changes 3) Decrease in stride length variability after 6 months, maintained at 12 months	Yes, motor under dual-task

(Continued)

Table 4 (Continued)

Study	Assessment points	Dual-task performance measures	Results	Dual-task improvement
Yamada et al, 2011 ³⁸	1) Pretest 2) Posttest	Gait speed while walking 10 meters carrying a ball on a tray. Dual-task cost calculated for both tasks	Walking time under manual task improved	Yes, motor
Donath et al, 2013 ³⁹	1) Pretest 2) Posttest	1) AP and ML (COP) under motor dual-task 2) Gait variability with motor dual-task	Improvement only for the intervention group in both dual-tasks measure	Yes, motor
Granacher et al, 2010 ⁴⁰	1) Pretest 2) Posttest	1) Gait velocity under dual and triple tasks 2) Cognitive performance while walking 3) Motor performance while walking (contact ring time)	1) Nonsignificant improvement 2) Nonsignificant improvement 3) Significant improvement in the intervention group but not in the control group	Yes, motor under dual-task
Hall et al, 2009 ⁴¹	1) Pretest 2) Posttest	Dual-task cost for postural and cognitive measures	Nonsignificant improvement	No
Pichierri et al, 2012 ⁴²	1) Pretest 2) Posttest	Voluntary step execution test under dual-task condition	Significant improvement of initiation time of forward and backward steps execution	Yes, motor
Plummer-D'Amato et al, 2012 ⁴³	1) Pretest 2) Posttest	6 minutes obstacle negotiation under single and dual-task (dual-task cost calculated for each task)	Nonsignificant improvement	No improvement
Uemura et al, 2012 ⁴⁴	1) Pretest 2) Posttest	1) Gait initiation under dual-task 2) Steady state gait (10 meters) under dual-task	1) Improvement only for the intervention group 2) Improvement for both groups but more effective for the intervention group	Yes, motor under dual-task
Verghese et al, 2010 ⁴⁵	1) Pretest 2) Posttest	Gait velocity while talking: reciting the alternate alphabet	Significant improvement	Yes, motor under dual-task
You et al, 2009 ⁴⁶	1) Pretest 2) Posttest	1) Gait velocity under dual-task 2) AP and ML (COP) 3) Memory performance data	1) Nonsignificant improvement in the intervention group, but significant improvement in the control group 2) Nonsignificant improvement in AP and ML 3) Significant improvement in the intervention group but not in the control group	Yes, cognitive under dual-task
Agmon et al, 2012 ⁴⁷	1) Pretest 2) Posttest	1) Timed Up and Go cognitive and 1 minute walk with verbal fluency a) Gait speed and walking under two tests b) Dual-task cost for walking c) Dual-task cost cognitive	a) Significant improvement for both tests b) Nonsignificant improvement for both tests c) Significant improvement for the Timed Up and Go and decline for 1 minute walk with verbal fluency	No
Bisson et al, 2007 ⁴⁸	1) Pretest 2) Posttest 3) 1 month after intervention	1) Reaction time under dual-task condition 2) Postural sway under dual-task conditions	1) Significant improvement in both groups at posttest and 1 month follow-up 2) No training effect	Yes, cognitive under dual-task
Lajoie et al, 2004 ⁴⁹	1) Pretest 2) Posttest 3) 2 weeks after intervention	1) Reaction time under dual-task condition	1) Significant improvement in the intervention group at posttest and 2 week follow-up compared to pretest, but not in the control group	Yes, cognitive under dual-task
Melzer et al, 2009 ⁵⁰	1) Single posttest measurement	2) Postural sway under dual-task condition Voluntary step execution test under dual-task	2) Nonsignificant improvement Only the preparation phase of dual-task step execution was faster in the exercise group compared to the control group	No improvement

Toulotte et al, 2006 ⁵¹	1) 3 months before training (T0) 2) 2 days before training (T1) 3) 2 days after training (T2) 4) 3 months after training (T3)	1) Timed Up and Go while holding glass full of water: a) Walking speed b) Stride time c) Single support time d) Cadence e) Stride length	a) T0 to T1: no change for either group; T1 to T2: increased for both groups; T2 to T3: decreased for both groups b) T0 to T1: no change for either group; T1 to T2: decreased for both groups; T2 to T3: increased for both groups c) T0 to T1: increased for both groups; T1 to T2: decreased for both groups; T2 to T3: increased for both groups d) T0 to T1: no change for either group; T1 to T2: increased for both groups; T2 to T3: decreased for nonfallers e) T0 to T1: no change for either group; T1 to T2: increased for both groups; T2 to T3: decreased for both groups	Yes, motor under dual-task
Silsupadol et al, 2006 ⁵²	1) Pretest 2) After 2 weeks 3) 12 weeks after intervention	Timed Up and Go cognitive	Significant improvement in all three participants. Greater improvement in the dual-task training participants	Yes, both cognitive and motor under dual-task

Abbreviations: AP and ML COP, anterior–posterior and medial–lateral center of pressure; COM-AJC, center of mass–ankle joint center inclination; VP, variable priority; COG, center of gravity.

Retention

Most studies measured outcomes immediately before and after the intervention, with only five studies examining retention of improvements at different time points after the end of the intervention.^{27,37,48,49,52} Two studies demonstrated improvements in dual-task cognitive performance that were retained at 2 weeks^{48,49} and 1 month postintervention.⁴⁸ Three studies showed some degree of retention in dual-task motor performance for periods ranging from 2 months to 6 months postintervention.^{27,37,52}

Transfer

Eight studies assessed whether training benefits transferred to untrained tasks.^{26,34,36,38,41,45,47,52} Three studies^{26,38,52} examined the effect of dual-task postural control training on novel or untrained dual-task postural control tasks. Silsupadol et al^{26,52} showed no transfer effect, while Yamada et al³⁸ demonstrated a transfer effect. Two studies^{41,47} examined whether single training benefits transferred to dual-task postural control, and showed negative results. Three studies^{34,36,45} measured the effect of cognitive dual-task training on dual-task performance involving a postural control task and showed transfer to some aspects of dual-task performance (see Table 3).

Discussion

This investigation of the literature on dual-task training demonstrates the potential to increase postural control, thereby improving balance and walking ability in older adults. Furthermore, this systematic review builds on previous research by examining specific training parameters that may impact the efficacy of dual-task interventions.

Training specificity

Overall, evidence supports the effectiveness of specific training to improve dual-task postural control among healthy older adults. Training specificity is a key element of motor learning.⁵⁵ However, the definition of training specificity is not obvious within the dual-task paradigm since the intended outcome of interventions could include either an improved ability to divide attention between both tasks or to preferentially improve performance of the postural control task. The majority of studies that incorporated direct dual-task training demonstrated improvement on some aspect of dual-task postural control, with only one exception.⁴³ Interventions that trained single-task postural control demonstrated improvement on measures of single-task balance and walking but not on dual-task postural control, with one exception.³⁹ Thus, training dual-task performance specifically, rather than just

single-task performance, appears to be a crucial element for interventions that aim to improve dual-task performance. This notion was supported by several studies with the highest level of evidence included in the review.^{26,27,33–35,37,38}

Training content

The interventions that directly trained dual-task postural control employed a variety of task combinations with different levels of difficulty in the postural control and concurrent cognitive or motor tasks. Some task combinations required mainly mathematical skills for the cognitive task^{26,27,52} whereas others required verbal and memory skills⁴¹ or auditory skills.³⁷ For the postural control task, most studies used walking^{26,27,40,46} or standing^{35,42,49,50} whereas a few used more complicated tasks such as walking within a narrow path²⁶ or obstacle crossing.⁴³ Two studies used two motor tasks such as standing while catching and throwing a ball,⁴⁰ and walking while holding a tray.³⁹ The only study⁴³ that specifically trained dual-task performance but did not show specific dual-task performance improvements identified a lack of specificity of outcome measures relative to the trained tasks as well as insufficient training hours (4 hours total) as potential explanations for this finding. While the current research suggests that a variety of trained task combinations can improve dual-task performance, future studies should compare how different combinations of tasks impact the efficacy of training.

The impact of specific task combinations on dual-task performance has been widely discussed. Recent reviews^{30,31} have examined the effect of different concurrent tasks on walking. Al-Yahya et al³¹ reported that tasks involving internal interference (ie, requiring top-down processing and driven by factors that are internal to the participant⁵⁶ such as verbal fluency or mathematical tasks) have a greater influence on gait parameters than external interference or bottom-up tasks (eg, reaction time tasks). Chu et al³⁰ assessed the predictive value of different task combinations for predicting falls. Their meta-analysis indicated that the combination of a mental tracking task and walking is a good predictor for falls among the elderly. Among the studies in this review, only seven studies used this combination in either the training protocol^{26,27,38,43,46,52} or outcome measure.^{26,27,36,43,45,47,52} Since fall prevention is an important goal of dual-task interventions, future studies should consider incorporating mental tracking tasks in combination with walking in their protocol and/or outcome measurement.

Instructions and feedback

Variable priority instructions, in which participants were asked to shift their attention back and forth between tasks,

appeared to be more effective for improving performance than fixed priority instructions, in which participants focused on either the postural control task or the concurrent cognitive or motor task.^{26,27,52} However, this direct comparison of instructions was limited to only three studies, one of which was a case series with only three participants. Thus, determining the most effective instructions for dual-task training merits further investigation.⁵⁷

Moreover, the effect of feedback⁵⁵ was not explored by any of these investigations, and different forms of feedback may influence motor learning differently among older adults.⁵⁸ Feedback focused on knowledge of results (for example, how many meters someone walks) is more effective than feedback focused on knowledge of performance (the nature of the movement). Tailored feedback during dual-task training could target each task separately or both tasks simultaneously. Nevertheless, even the effect of feedback during motor learning of a single task among the elderly was not clear.⁵⁹ Several issues should be addressed regarding the optimal use of feedback during dual-task training, including: 1) whether feedback is more effective than the absence of feedback during dual-task training among the elderly; and 2) whether feedback on one task at a time is more effective than feedback on both tasks simultaneously during dual-task training.

Training parameters reflecting motor learning

Several parameters of dual-task training may promote motor learning. The setting of training may influence efficacy. The interventions reviewed here were conducted in two distinct settings: group training and one-on-one training. Similar rates of success were found in both settings; ten out of 13 interventions conducted in a group setting^{26,27,33,35,37–40,42,44,51} demonstrated improvement in some aspects of dual-task, while six out of ten conducted in one-on-one settings^{34,36,45,46,48,49} demonstrated successful outcomes. The dose of intervention is an important factor influencing motor learning. The total training hours varied from 5 hours⁴⁸ to 25 hours³⁷ and were spread from 1 week to over 25 weeks.³⁷ For studies that showed improvement in dual-task performance, the length of training sessions ranged from 20 minutes³⁸ to 60 minutes.^{33–37,39–42,47,49–51} Among studies with the highest methodological quality, Silsupadol et al^{26,27} conducted the most intensive training, with three sessions per week for 1 month, but Trombetti et al³⁷ conducted the longest intervention, with one session per week for 25 weeks. Both of these studies demonstrated some degree of improvement in dual-task walking, with

some retention of benefits, providing support for a high training dose. Among the studies in this review, the variability in training dose, including session intensity, session frequency, and training duration, makes determining the optimal dose of dual-task training challenging.

Outcome measures

There is no gold standard for dual-task assessment, and the studies in this review used various dual-task combinations for their outcome measures. For example, the postural control tasks included obstacle crossing,^{26,43} over ground walking,^{34,40,44,45} or standing on a force plate.⁴⁸ Within similar tasks, a variety of different parameters were assessed. For example, the parameters used to assess performance during walking included frontal plane inclination angle,²⁶ gait variability,^{37,39} or gait speed.^{27,40,45}

Recent research emphasizes the importance of calculating dual-task cost to understand the underlying mechanism of improvement in dual-task performance⁷ and as a sensitive means of assessing fall risk.^{13,14} Dual-task cost is often calculated using the formula:

$$(\text{dual-task} - \text{single task})/\text{single task} \times 100 \quad [1]$$

and expressed as percentage. Using dual-task cost elucidates the mechanisms underlying the improvement by demonstrating whether improvements are achieved in both tasks or whether improvements in one task occur at the expense of the other task.⁷ Among the studies included in the current review, only three^{41,43,47} calculated the dual-task cost; therefore, comparisons between studies are limited. Agmon et al⁴⁷ showed that there was a change in the trade-off between cognitive and motor costs between pre- and postintervention. This finding demonstrates changes in prioritization between tasks, but not actual improvement with training.

Retention

Retention of motor skills for up to 1 year has been demonstrated in humans in a laboratory setting.⁶⁰ Among the studies demonstrating the highest level of evidence, Trombetti et al³⁷ demonstrated the longest retention (12 months), although this was only a partial retention of improvements. Future studies should investigate which practice conditions promote optimal retention as well as the effects of interventions that incorporate ongoing or maintenance programs.

Transfer

Some studies included in this review demonstrated evidence of transfer from trained tasks to novel tasks, but this finding

was not uniform. Several studies demonstrated that dual-task postural control training transferred to improvements in novel dual-task combinations.^{26,38,52} Interestingly, three studies that trained cognitive dual-task performance demonstrated improvements on dual-task postural control.^{34,36,45} Participants in these studies were trained, while sitting, on tasks that required switching and dividing attention, and the impact on dual-task walking^{36,45} and standing³⁴ was assessed. These studies illuminate the potential to improve dual-task postural control by training two cognitive tasks; a protocol that emphasizes the ability to divide or rapidly shift attention. However, these findings were not consistent and may depend on the level of difficulty of the trained tasks compared to the measured task.⁴³

Future directions

This review highlights several questions that merit further exploration. First, although the existing research provides support for the ability to improve dual-task performance, particularly following response to specific dual-task training, it is not clear what effects these improvements have on function in daily life or fall risk. Outcome measures could be expanded to include functional measurements of dual-task performance relevant to daily life, such as putting on a shirt while standing or walking while talking on the phone. In order to understand the influence of dual-task interventions on falls prevention, future studies should incorporate prospective falls assessments over longer-term follow-up periods.

Second, there is a need to further examine the effect of different motor learning parameters, and the interaction between them, on dual-task acquisition, retention, and transfer. These might include the influence of instructions or different modes of feedback, the specificity of training, and the effect of dose on the response to training. In addition, further exploration is needed to determine the efficacy of training within subgroups of older adults, such as those with and without a history of falls or with different cognitive abilities.

Third, Li et al⁶¹ emphasized the importance of adopting an ecological perspective when training and measuring dual-task performance. Thus, finding new ways to address dual-task performance in valid ecological environments needs to be explored. Recently, Mirelman et al⁶² suggested a treadmill with a virtual reality protocol aimed at fall reduction. Such protocols should be tested first on subjects in clinical settings, followed by testing in home-based users. Home-based virtual reality training has the potential to reach larger populations in a complex and safe environment.⁶³

Finally, in order to strengthen the evidence base for improving dual-task postural control, future studies should include larger, more representative samples and use a standard set of outcome measures to allow cross-study comparison. Outcome measures should include walking speed and stride-to-stride variability,⁷ standard cognitive tasks (such as those that require mental tracking³⁰ or internal interference processing),³¹ and calculation of dual-task cost for both tasks.⁶⁴

Limitations

Sixteen randomized clinical trials^{26,27,33–46} evaluated the effectiveness of training on dual-task performance on a total of 516 subjects with heterogeneous measures that precluded quantitative synthesis or meta-analysis. Furthermore, the quality of evidence in the present review was mixed, with a risk of bias because some studies did not use randomization. The sample was predominately female, limiting the ability to investigate sex differences in intervention efficacy.

Studies incorporated different intervention protocols and various outcome measures to assess the effectiveness of the interventions. This variability made it difficult to identify specific recommendations about the optimal content or duration of dual-task interventions. The lack of long-term follow-up limits the ability to determine whether the benefits of these interventions were retained, as well as the ability to understand the impact of dual-task postural control training on fall risk.

Conclusion

A synthesis of research examining the effect of different interventions on dual-task postural control suggests that interventions training for balance under single-task conditions can improve balance under single-task conditions, but this improvement does not transfer to dual-task performance.^{41,47,50} Instead, dual-task training appears to be necessary to improve dual-task performance. While variability amongst studies makes it difficult to identify optimal parameters of interventions, it appears that effective interventions can be conducted in either group or one-on-one settings, with a variety of task combinations incorporated into the intervention. The shortest training schedule of 20 minutes twice a week for 24 weeks³⁸ as well as only five sessions of 1 hour each³⁴ demonstrated improvement in some aspects of dual-task performance.

Future investigations of interventions to improve dual-task postural control should include focused dual-task training and address tasks that have the highest correlation with fall risk. Moreover, long term follow-up with regard to

fall occurrence and daily function should be incorporated in order to better understand whether improved dual-task postural control impacts these areas. Future research should also focus on motor learning elements that may extend the retention of dual-task training benefits in order to determine the most effective protocols. Finally, in order to achieve comparability between interventions, an agreed-upon set of outcome measures should be defined and dual-task cost calculation should be included.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing*. 2006;35 Suppl 2:ii37–ii41.
2. Maki BE, Cheng KC, Mansfield A, et al. Preventing falls in older adults: new interventions to promote more effective change-in-support balance reactions. *J Electromyogr Kinesiol*. 2008;18(2):243–254.
3. Inzitari M, Baldereschi M, Di Carlo A, et al. Impaired attention predicts motor performance decline in older community-dwellers with normal baseline mobility: results from the Italian Longitudinal Study on Aging (ILSA). *J Gerontol A Biol Sci Med Sci*. 2007;62(8):837–843.
4. Wood KM, Edwards JD, Clay OJ, Wadley VG, Roenker DL, Ball KK. Sensory and cognitive factors influencing functional ability in older adults. *Gerontology*. 2005;51(2):131–141.
5. Campbell A, Reinken J, Allan B, Martinez G. Falls in old age: a study of frequency and related clinical factors. *Age Ageing*. 1981;10(4): 264–270.
6. Nevitt MC, Cummings SR, Kidd S, Black D. Risk factors for recurrent nonsyncopal falls. *JAMA*. 1989;261(18):2663–2668.
7. Montero-Odasso M, Verghese J, Beauchet O, Hausdorff JM. Gait and cognition: a complementary approach to understanding brain function and the risk of falling. *J Am Geriatr Soc*. 2012;60(11):2127–2136.
8. Fuller GF. Falls in the elderly. *Am Fam Physician*. 2000;61(7): 2159–2168.
9. Shumway-Cook A, Silver IF, LeMier M, York S, Cummings P, Koepsell TD. Effectiveness of a community-based multifactorial intervention on falls and fall risk factors in community-living older adults: a randomized, controlled trial. *J Gerontol A Biol Sci Med Sci*. 2007;62(12): 1420–1427.
10. Woollacott M, Shumway-Cook A. Attention and the control of posture and gait: a review of an emerging area of research. *Gait Posture*. 2002;16(1):1–14.
11. Lundin-Olsson L, Nyberg L, Gustafson Y. “Stops walking when talking” as a predictor of falls in elderly people. *Lancet*. 1997;349(9052):617.
12. Dumas M, Rapp MA, Krampe RT. Working memory and postural control: adult age differences in potential for improvement, task priority, and dual tasking. *J Gerontol B Psychol Sci Soc Sci*. 2009;64(2): 193–201.
13. Yamada M, Aoyama T, Arai H, et al. Dual task walk is a reliable predictor of falls in robust elderly adults. *J Am Geriatr Soc*. 2011;59(1): 163–164.
14. Beauchet O, Annweiler C, Dubost V, et al. Stops walking when talking: a predictor of falls in older adults? *Eur J Neurol*. 2009;16(7):786–795.
15. Hausdorff JM, Rios DA, Edelberg HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil*. 2001;82(8):1050.
16. Lezak M, editor. *Neuropsychological Assessment*. New York, NY: Oxford University Press; 2004.
17. Royall DR, Palmer R, Chiodo LK, Polk MJ. Declining executive control in normal aging predicts change in functional status: the Freedom House Study. *J Am Geriatr Soc*. 2004;52(3):346–352.

18. Springer S, Giladi N, Peretz C, Yogev G, Simon ES, Hausdorff JM. Dual-tasking effects on gait variability: The role of aging, falls, and executive function. *Mov Disord*. 2006;21(7):950–957.
19. van Iersel MB, Kessels RP, Bloem BR, Verbeek AL, Rikkert MG. Executive functions are associated with gait and balance in community-living elderly people. *J Gerontol A Biol Sci Med Sci*. 2008;63(12):1344–1349.
20. Coppin AK, Shumway-Cook A, Saczynski JS, et al. Association of executive function and performance of dual-task physical tests among older adults: analyses from the InChianti study. *Age Ageing*. 2006;35(6):619–624.
21. Zijlstra A, Ufkes T, Skelton D, Lundin-Olsson L, Zijlstra W. Do dual tasks have an added value over single tasks for balance assessment in fall prevention programs? A mini-review. *Gerontology*. 2008;54(1):40–49.
22. Nordin E, Moe-Nilssen R, Ramnemark A, Lundin-Olsson L. Changes in step-width during dual-task walking predicts falls. *Gait Posture*. 2010;32(1):92–97.
23. Muhaidat J, Kerr A, Evans JJ, Pilling M, Skelton DA. Validity of simple gait-related dual-task tests in predicting falls in community-dwelling older adults. *Arch Phys Med Rehabil*. 2014;95(1):58–64.
24. Wollesen B, Voelcker-Rehage C. Training effects on motor–cognitive dual-task performance in older adults. *Eur Rev Aging Phys Act*. 2013;1–20.
25. Beurskens R, Bock O. Age-related deficits of dual-task walking: a review. *Neural Plast*. 2012;2012:131608.
26. Silsupadol P, Lugade V, Shumway-Cook A, et al. Training-related changes in dual-task walking performance of elderly persons with balance impairment: a double-blind, randomized controlled trial. *Gait Posture*. 2009;29(4):634–639.
27. Silsupadol P, Shumway-Cook A, Lugade V, et al. Effects of single-task versus dual-task training on balance performance in older adults: a double-blind, randomized controlled trial. *Arch Phys Med Rehabil*. 2009;90(3):381–387.
28. Shumway-Cook A, Woollacott MH. Aging and postural control. In: Shumway-Cook A, Woollacott MH, editors. *Motor Control Translating Research into Clinical Practice*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2007:212–232.
29. Wulf G. *Attention and Motor Skill Learning*. Champaign, IL: Human Kinetics Publishers; 2007.
30. Chu YH, Tang PF, Peng YC, Chen HY. Meta analysis of type and complexity of a secondary task during walking on the prediction of elderly falls. *Geriatr Gerontol Int*. 2013;13(2):289–297.
31. Al-Yahya E, Dawes H, Smith L, Dennis A, Howells K, Cockburn J. Cognitive motor interference while walking: a systematic review and meta-analysis. *Neurosci Biobehav Rev*. 2011;35(3):715–728.
32. Portney LG, Watkins MP. *Foundations of Clinical Research: Applications to Practice*. 3rd ed: Upper Saddle River, NJ: Pearson and Prentice Hall; 2009.
33. Hiyamizu M, Morioka S, Shomoto K, Shimada T. Effects of dual task balance training on dual task performance ability in elderly people: a randomized controlled trial. *Clin Rehabil*. 2012;26(1):58–67.
34. Li KZ, Roudaia E, Lussier M, Bherer L, Leroux A, McKinley PA. Benefits of cognitive dual-task training on balance performance in healthy older adults. *J Gerontol A Biol Sci Med Sci*. 2010;65(12):1344–1352.
35. Melzer I, Oddsson LLe. Improving balance control and self-reported lower extremity function in community-dwelling older adults: a randomized control trial. *Clin Rehabil*. 2013;27(3):195–206.
36. Mozolic JL, Long AB, Morgan AR, Rawley-Payne M, Laurienti PJ. A cognitive training intervention improves modality-specific attention in a randomized controlled trial of healthy older adults. *Neurobiol Aging*. 2011;32(4):655–668.
37. Trombetti A, Hars M, Herrmann FR, Kressig RW, Ferrari S, Rizzoli R. Effect of music-based multitask training on gait, balance, and fall risk in elderly people: a randomized controlled trial. *Arch Intern Med*. 2011;171(6):525–533.
38. Yamada M, Aoyama T, Hikita Y, et al. Effects of a DVD-based seated dual-task stepping exercise on the fall risk factors among community-dwelling elderly adults. *Telemed J E Health*. 2011;17(10):768–772.
39. Donath L, Faude O, Bridenbaugh SA, et al. Transfer effects of fall training on balance performance and spatio-temporal gait parameters in healthy community-dwelling seniors: a pilot study. *J Aging Phys Act*. Epub July 22, 2013.
40. Granacher U, Muehlbauer T, Bridenbaugh S, Bleiker E, Wehrle A, Kressig RW. Balance training and multi-task performance in seniors. *Int J Sports Med*. 2010;31(5):353–358.
41. Hall CD, Miszko T, Wolf SL. Effects of Tai Chi intervention on dual-task ability in older adults: a pilot study. *Arch Phys Med Rehabil*. 2009;90(3):525–529.
42. Pichierri G, Coppe A, Lorenzetti S, Murer K, de Bruin ED. The effect of a cognitive-motor intervention on voluntary step execution under single and dual task conditions in older adults: a randomized controlled pilot study. *Clin Interv Aging*. 2012;7:175–184.
43. Plummer-D'Amato P, Cohen Z, Daee NA, Lawson SE, Lizotte MR, Padilla A. Effects of once weekly dual-task training in older adults: A pilot randomized controlled trial. *Geriatr Gerontol Int*. 2012;12(4):622–629.
44. Uemura K, Yamada M, Nagai K, et al. Effects of dual-task switch exercise on gait and gait initiation performance in older adults: Preliminary results of a randomized controlled trial. *Arch Gerontol Geriatr*. 2012;54(2):e167–e171.
45. Verghese J, Mahoney J, Ambrose AF, Wang C, Holtzer R. Effect of cognitive remediation on gait in sedentary seniors. *J Gerontol A Biol Sci Med Sci*. 2010;65(12):1338–1343.
46. You JH, Shetty A, Jones T, Shields K, Belay Y, Brown D. Effects of dual-task cognitive-gait intervention on memory and gait dynamics in older adults with a history of falls: a preliminary investigation. *Neuro Rehabilitation*. 2009;24(2):193–198.
47. Agmon M, Kelly VE, Logsdon RG, Nguyen H, Belza B. The effect of enhanced fitness (EF) training on dual-task walking in older adults. *J Appl Gerontol*. Epub December 11, 2012.
48. Bisson E, Contant B, Sveistrup H, Lajoie Y. Functional balance and dual-task reaction times in older adults are improved by virtual reality and biofeedback training. *Cyberpsychol Behav*. 2007;10(1):16–23.
49. Lajoie Y. Effect of computerized feedback postural training on postural and attentional demands in older adults. *Aging Clin Exp Res*. 2004;16(5):363–368.
50. Melzer I, Marx R, Kurz I. Regular exercise in the elderly is effective to preserve the speed of voluntary stepping under single-task condition but not under dual-task condition. *Gerontology*. 2009;55(1):49–57.
51. Toulotte C, Thevenon A, Fabre C. Effects of training and detraining on the static and dynamic balance in elderly fallers and non-fallers: a pilot study. *Disabil Rehabil*. 2006;28(2):125–133.
52. Silsupadol P, Siu KC, Shumway-Cook A, Woollacott MH. Training of balance under single- and dual-task conditions in older adults with balance impairment. *Phys Ther*. 2006;86(2):269–281.
53. Kramer AF, Larish JF, Strayer DL. Training for attentional control in dual task setting: A comparison of young and old adults. *J Exp Psychol Appl*. 1995;1(1):50–76.
54. Hall E, Hylemon P, Vlahcevic Z, et al. Overexpression of CYP27 in hepatic and extrahepatic cells: role in the regulation of cholesterol homeostasis. *Am J Physiol Gastrointest Liver Physiol*. 2001;281(1):G293–G301.
55. Magill RA. *Motor Learning and Control: Concepts and Applications*. Volume 11. New York: McGraw-Hill; 2007.
56. Awh E, Belopolsky AV, Theeuwes J. Top-down versus bottom-up attentional control: A failed theoretical dichotomy. *Trends Cogn Sci*. 2012;16(8):437–443.
57. Del Rosario N. *Reasons to Switch: The Effects of Priority and Information Presentation on Dual-Task Interleaving Strategies [master's dissertation]*. London: University College London; 2009.

58. Shea CH, Wulf G. Enhancing motor learning through external-focus instructions and feedback. *Hum Mov Sci.* 1999;18(4):553–571.
59. Voelcker-Rehage C. Motor-skill learning in older adults – a review of studies on age-related differences. *Eur Rev Aging Phys Act.* 2008;5(1):5–16.
60. Dayan E, Cohen LG. Neuroplasticity subserving motor skill learning. *Neuron.* 2011;72(3):443–454.
61. Li KZH, Krampe RT, Bondar A. An ecological approach to studying aging and dual-task performance. *Cognitive Limitations in Aging and Psychopathology.* 2005:190–218.
62. Mirelman A, Rochester L, Reelick M, et al. V-TIME: a treadmill training program augmented by virtual reality to decrease fall risk in older adults: study design of a randomized controlled trial. *BMC Neurol.* 2013;13(1):15.
63. Holmes JD, Gu ML, Johnson AM, Jenkins ME. The effects of a home-based virtual reality rehabilitation program on balance among individuals with Parkinson's disease. *Phys Occup Ther Geriatr.* 2013;31(3):241–253.
64. Kelly VE, Janke AA, Shumway-Cook A. Effects of instructed focus and task difficulty on concurrent walking and cognitive task performance in healthy young adults. *Exp Brain Res.* 2010;207(1–2):65–73.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, the American Chemical Society's 'Chemical Abstracts

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress

Service' (CAS), Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.