An observational study of bimatoprost 0.01% in patients on prior intraocular pressure-lowering therapy: the Canadian Lumigan® RC Early Analysis Review (CLEAR) trial

Andrew C Crichton¹
Donald R Nixon²
Susan Simonyi³
Meetu Bhogal³
Christopher S Sigouin⁴
Marino J Discepolo⁵
Cindy ML Hutnik⁶
Darryl C Baptiste³
David B Yan⁷

On behalf of the CLEAR Study Group

¹Division of Ophthalmology, University of Calgary, Calgary, AB, Canada;

²Private practice, Barrie, ON, Canada;

³Medical Affairs, Allergan Inc., Markham, ON, Canada; ⁴CLINWest Research, Burlington, ON, Canada; ⁵Department of Ophthalmology, McGill University, Montreal, QC, Canada; ⁶Department of Ophthalmology and Pathology, Ivey Eye Institute, London, ON, Canada;

⁷Department of Ophthalmology, University of Toronto, Toronto, ON, Canada

Purpose: To evaluate the ocular hyperemia and intraocular pressure (IOP)-lowering efficacy of bimatoprost 0.01% in subjects with elevated IOP due to primary open-angle glaucoma (POAG) or ocular hypertension (OHT) in a real-world clinical setting.

Subjects and methods: This open-label, 12-week, observational study was conducted at 67 centers in Canada. Subjects with elevated IOP due to POAG or OHT instilled bimatoprost 0.01% as monotherapy once daily. Ocular hyperemia was graded by the investigator at baseline, week 6, and week 12 using a standardized photographic 5-point grading scale. Change in IOP from baseline was also evaluated at these time points. This analysis includes the subgroup of 268 subjects who had been previously treated with latanoprost 0.005%, bimatoprost 0.03%, travoprost 0.004%, and travoprost 0.004% with SofZia™ or nonselective beta-adrenergic receptor blockers prior to the study.

Results: After 12 weeks of treatment with 0.01% bimatoprost, ocular hyperemia was graded as none-to-mild hyperemia (grades 0, +0.5, or +1) for 94.1% of subjects and as moderate-to-severe hyperemia (grades +2 or +3) for 5.9%. No statistically significant shifts in ocular hyperemia ratings were observed at week 12 for any of the prior IOP-lowering therapies except bimatoprost 0.03%, in which 20.8% of subjects experienced an improvement. The mean percentage change from baseline IOP at week 12 following the switch to bimatoprost 0.01% monotherapy ranged from $-2.3\% \pm 17.3\%$ to $-26.3\% \pm 12.4\%$. Furthermore, the decreased mean percentage change from baseline IOP was statistically significant across all prior IOP-lowering medications, except for bimatoprost 0.03% at the 6- and 12-week visits and travoprost 0.004% at the 6-week visit.

Conclusion: This observational study demonstrates that bimatoprost 0.01% was well tolerated among POAG and OHT subjects who switched from prior IOP-lowering medication. Furthermore, a switch in ocular hypertensive treatment to bimatoprost 0.01% was associated with an additional 10%–15% reduction in IOP.

Keywords: glaucoma, intraocular pressure, hyperemia, bimatoprost

Introduction

Prostaglandin analogs (PGAs) and prostamides (PMs) are considered first-line therapies for the treatment of glaucoma and ocular hypertension (OHT), because they effectively lower intraocular pressure (IOP) and have a good safety profile.¹ The most common adverse event (AE) associated with both PGAs and PMs is conjunctival hyperemia. This side effect may be bothersome to patients to the extent that it reduces adherence to treatment.²

Correspondence: Andrew Crichton
Division of Ophthalmology, Faculty of Medicine, University of Calgary,
102-49 Richard Way SW, Calgary,
AB T3E 7M8, Canada
Tel +1 403 245 3730
Fax +1 403 245 1058
Email drcricht@telusplanet.net

Bimatoprost 0.03% (Lumigan®; Allergan, Inc., Irvine, CA, USA) is a PM approved for the safe and effective lowering of IOP in patients with open-angle glaucoma (OAG) or OHT.³⁻⁵ Several studies have found that bimatoprost 0.03% leads to greater reductions in IOP than latanoprost 0.005% (Xalatan®, Pfizer, Inc., New York, NY, USA) or travoprost 0.004% (Travatan® original version preserved with benzalkonium chloride [BAK]; Alcon Laboratories, Inc., Fort Worth, TX, USA), and may be useful for patients who do not respond to latanoprost.^{4,6,7} However, in comparison to PGA options, bimatoprost 0.03% is associated with a higher rate of conjunctival hyperemia, which in some cases may lead to discontinuation.⁴ Bimatoprost 0.01% (Lumigan® RC; Allergan, Inc., Irvine, CA, USA) was developed with the goal of creating a formulation of bimatoprost that would maintain the IOP-lowering efficacy achieved with bimatoprost 0.03% while offering an improved overall safety profile, particularly, improved ocular surface tolerability.³

In a Phase III, randomized, double-masked, multicenter clinical trial, bimatoprost 0.01% proved to have equivalent IOP-lowering efficacy to bimatoprost 0.03% through 12 months of treatment, while also showing improved tolerability, including less frequent and severe conjunctival hyperemia.³ Although these data provided evidence of the safety and efficacy of bimatoprost 0.01%, these findings were derived from a glaucoma and OHT patient population that met defined inclusion criteria that may not represent the diversity of patients typical of a given clinical practice. As a result, the “real-world” safety and efficacy of bimatoprost 0.01% as an ocular hypotensive therapy may be different from the results reported by Katz et al³ if bimatoprost 0.01% is administered to a more diverse patient population. The aim of the present study – the Canadian Lumigan® RC Early Analysis Review (CLEAR) Trial – was to determine the external validity of bimatoprost 0.01% in a real-world clinical setting. We reported the safety and efficacy of bimatoprost 0.01% in treatment-naïve primary OAG (POAG) and OHT subjects in a separate paper.⁸ Herein, we report the safety and efficacy of switching POAG and OHT subjects from a single IOP-lowering therapy to bimatoprost 0.01% monotherapy.

Subjects and methods

Study design

This 12-week, open-label, non-controlled, prospective, observational, multicenter (67 Canadian centers) study evaluated the occurrence and severity of ocular hyperemia following a switch to bimatoprost 0.01% in subjects with POAG or OHT. This study was approved by the respective institutional review

boards or independent ethics committees at each site, and followed the regulations of the Personal Information Protection and Electronic Documents Act and the Good Clinical Practice guidelines of the Declaration of Helsinki.

Study population

Eligible subjects were at least 18 years of age, diagnosed with elevated IOP due to either POAG or OHT, and determined by the treating physician to require treatment with bimatoprost 0.01% either as first-line monotherapy, as a switch of prior IOP-lowering medication, or as adjunctive IOP-lowering medication. This paper focuses on the subgroup of subjects who switched to bimatoprost 0.01% from a prior IOP-lowering medication. Subjects who were willing to return for all scheduled study visits signed an informed consent prior to or concurrent with the screening visit.

Subjects were excluded if they had hypersensitivity to BAK, any PGA, PM, or any other component of the study medication, or had any other abnormal ocular condition or symptom that would prevent them from entering the trial according to the treating physician's judgment. Female subjects who were pregnant, nursing, or of childbearing potential and not using adequate contraception, as well as any subject with an inability to adhere to the treatment or study visit plan, or subjects who had participated in any other clinical trial involving an investigational product of a new chemical entity within 6 months prior to the screening or baseline visit were also excluded.

Treatment

Bimatoprost 0.01% was obtained by the patient through commercial means, and not as investigational study drug provided by Allergan, Inc. At the investigator's discretion, the patient's prior IOP-lowering medication was switched to bimatoprost 0.01% (baseline visit), either because the patient required additional IOP lowering or the patient could not tolerate the prior IOP-lowering therapy. All subjects were instructed to self-instill bimatoprost 0.01% into the affected eye(s) in the evening at approximately 8 pm. No washout period was applied on switching to bimatoprost 0.01% to reflect more closely real-world practice patterns. Other therapy considered necessary for the patient's welfare was given at the discretion of the treating physician and was documented.

Outcome variables

Outcome variables were evaluated at baseline and at weeks 6 and 12 at 10 am (± 2 hours). The primary outcome variable was the occurrence and severity of ocular hyperemia at week 12, using a photographic 5-point grading

scale: 0= none – normal; +0.5= trace – trace flush reddish pink; +1= mild – mild flush, reddish color; +2= moderate – bright red color; +3= severe – deep, bright, diffuse redness.³

Secondary outcome variables included IOP change and IOP percentage change from baseline at weeks 6 and 12. Each participating site had an assigned examiner to measure IOP at each study visit. Although the type of tonometer was not specified in the protocol, all sites were instructed to use the same tonometer throughout the study for consistency. AEs were documented throughout the study, including AE seriousness and severity, action taken and relationship of the AE to treatment with bimatoprost 0.01%.

Statistical analyses

All subjects who provided informed consent and completed the screening/baseline visit (visit 1) were included in the intent-to-treat analysis. The intent-to-treat analysis used only data from the eye with the higher IOP at baseline (worse eye). If both eyes had the same IOP at baseline, the right eye was used.

Hyperemia grading was reported individually as 0, +0.5, +1, +2, and +3. Subsequently, the hyperemia scores were collapsed into two categories: 1) none to mild = subjects with hyperemia scores of 0, +0.5, or +1, and 2) moderate to severe = subjects with hyperemia scores of +2 or +3. These data were summarized as frequency counts and percentages at baseline and at weeks 6 and 12. The changes in collapsed hyperemia grading response from baseline at weeks 6 and 12 were summarized as improved, no change or worsened.

An improvement in hyperemia following a switch of IOP-lowering therapy to bimatoprost 0.01% was defined as a change in hyperemia grouping from moderate-to-severe hyperemia to none-to-mild hyperemia. No change in hyperemia was defined as staying within the same hyperemia grouping at baseline and on treatment with bimatoprost 0.01%. A worsening in hyperemia following a switch of IOP-lowering therapy to bimatoprost 0.01% was defined as a change in hyperemia grouping from none-to-mild hyperemia to moderate-to-severe hyperemia. Two-sided McNemar tests were used to evaluate treatment effects. IOP change and IOP percentage change from baseline were analyzed at weeks 6 and 12 using two-sided paired *t*-tests.

The present study was exploratory in nature, and as such no formal sample-size calculations were carried out. SAS for Windows version 9.2 (SAS Institute, Cary, NC, USA) was used for the analysis. All statistical tests were performed as two-sided tests with the significance level set at $P \leq 0.05$. *P*-values were not adjusted for multiple comparisons.

Results

Demographics

A total of 1,137 subjects were enrolled in the CLEAR Trial across 67 Canadian centers. Of the total population of subjects, 522 were naive to IOP-lowering treatment, and these were considered in a separate paper.⁸ The remaining 615 subjects had been treated with a prior IOP-lowering therapy and were switched to bimatoprost 0.01% as adjunctive IOP-lowering therapy ($n=165$) or as monotherapy ($n=450$). A complete list of the 450 subjects who switched prior IOP-lowering therapy to receive bimatoprost 0.01% monotherapy is presented in Table 1.

The present report focuses on the subset of 268 subjects who were switched from a prior IOP-lowering monotherapy to bimatoprost 0.01% monotherapy and had a sample size greater than 15 (ie, latanoprost 0.005% $n=113$, bimatoprost 0.03% $n=63$, travoprost 0.004% $n=16$, travoprost 0.004% with SofZia™ [Travatan Z®; Alcon Laboratories, Inc.] $n=36$, and nonselective beta-adrenergic blockers $n=40$). Thirty-two of the 268 subjects discontinued for the following reasons: lost to follow-up, $n=13$ (4.9%); ocular AE, $n=5$ (1.9%); other AE, $n=2$ (0.7%); and reasons not related to AEs, $n=12$ (4.5%). Of the five monotherapy-switch subjects who discontinued the study due to ocular AEs, a total of nine ocular AEs related to bimatoprost 0.01% monotherapy were reported (Table 2).

Of the 268 subjects who received bimatoprost 0.01% monotherapy, 142 (53.0%) were female (mean age 67.5 years) and 126 (47.0%) were male (mean age 67.3 years). Common associated medical comorbidities among the enrolled 268 switch patients included hypertension ($n=125$, 46.6%), cardiovascular disease ($n=48$, 17.9%), diabetes ($n=25$, 9.3%), asthma ($n=22$, 8.2%), and pulmonary disease ($n=15$, 5.6%).

Ocular hyperemia

Among those who were switched from a prior IOP-lowering therapy to bimatoprost 0.01% monotherapy, the baseline numbers (and percentages) of subjects presenting with each hyperemia grade (ie, 0, +0.5, +1, +2, and +3) are presented in Table 3.

The numbers (and percentages) of subjects who were switched from either latanoprost 0.005%, travoprost 0.004%, travoprost 0.004% with SofZia™ or a nonselective beta-adrenergic receptor blocker to bimatoprost 0.01% and demonstrated none-to-mild or moderate-to-severe hyperemia at baseline, week 6, and week 12 are presented in Table 4.

The percentages of subjects who switched from one prior IOP-lowering therapy and showed improved, no change, or

Table 1 Switch-subject disposition

Prior intraocular pressure (IOP)-lowering therapy	Baseline n	Week 6 n (discontinued)	Week 12 n (discontinued)
Adrenergic agonists	5	4 (1)	4 (1)
• Alphagan®	1	1 (0)	1 (0)
• Alphagan® P	3	2 (1)	2 (1)
• Generic brimonidine tartrate	1	1 (0)	1 (0)
Selective beta-adrenergic receptor blockers	3	2 (1)	2 (1)
• Betoptic®	2	2 (0)	2 (0)
• Generic betaxolol hydrochloride	1	0 (1)	0 (1)
Nonselective beta-adrenergic receptor blockers	40	37 (3)	35 (5)
• Betagan®	1	1 (0)	1 (0)
• Generic levobunolol hydrochloride	3	3 (0)	3 (0)
• Generic timolol maleate	10	8 (2)	6 (4)
• Timoptic®	23	22 (1)	22 (1)
• Timoptic XE® 0.5%	3	3 (0)	3 (0)
Carbonic anhydrase inhibitors	6	6 (0)	6 (0)
• Azopt®	4	4 (0)	4 (0)
• Trusopt®	2	2 (0)	2 (0)
Prostaglandin analogs and prostamides	228	216 (12)	202 (26)
• Lumigan® 0.03%	63	58 (5)	53 (10)
• Travatan®	16	16 (0)	16 (0)
• Travatan Z®	36	36 (0)	35 (1)
• Xalatan®	113	106 (7)	98 (15)
Non-prostaglandin analog combinations	29	27 (2)	27 (2)
• Azarga®	1	0 (1)	0 (1)
• Combigan®	15	14 (1)	14 (1)
• Cosopt®	13	13 (0)	13 (0)
Prostaglandin analog combinations	18	16 (2)	16 (2)
• DuoTrav®	12	10 (2)	10 (2)
• Xalacom®	6	6 (0)	6 (0)
More than one prior IOP-lowering therapy	21	21 (0)	21 (0)
Unavailable prior IOP-lowering therapy	100	95 (5)	87 (13)

Notes: The bold IOP-lowering therapies represent the subset of 268 subjects who switched from prior IOP-lowering monotherapy to bimatoprost 0.01% monotherapy and were the focus of analysis in this report. Manufacturer details are as follows: Alphagan®, Alphagan® P, Betagan®, Combigan®, Lumigan® (Allergan, Inc., Irvine, CA, USA); Azarga®, Betoptic®, Azopt®, DuoTrav®, Travatan®, Travatan Z® (Alcon Laboratories, Inc., Fort Worth, TX, USA); Timoptic®, Timoptic XE®, Trusopt®, Cosopt® (Merck & Co., Inc., Whitehouse Station, NJ, USA); Xalacom®, Xalatan® (Pfizer, Inc., New York, NY, USA).

worsened ocular hyperemia grading at the week 6 and week 12 visits are displayed in Table 5. At week 6, 96.2% of subjects who were previously treated with latanoprost 0.005% showed no change, and 3.8% demonstrated worsened hyperemia grade. This change in ocular hyperemia grade was statistically significant ($P=0.046$). By week 12, the change in ocular hyperemia

grade for subjects previously treated with latanoprost 0.005% was no longer statistically significant ($P=0.180$). Subjects previously treated with bimatoprost 0.03% who switched to bimatoprost 0.01% demonstrated improved (19.0% and 20.8%), no change (77.6% and 77.4%), or worsened (3.4% and 1.9%) ocular hyperemia grades at week 6 and week 12,

Table 2 Early discontinuation due to treatment-related ocular adverse events

Adverse event	Subjects, n (%)	Events, n
Eye pruritus	1 (0.4)	2
Ocular hyperemia	1 (0.4)	2
Drug ineffective	1 (0.4)	1
Foreign body sensation in eyes	1 (0.4)	1
Erythema	1 (0.4)	1
Sensation of foreign body	1 (0.4)	1
Conjunctivitis	1 (0.4)	1

Table 3 Overall rates of occurrence and severity of ocular hyperemia

Hyperemia grade	Number (%) of subjects with each hyperemia grade		
	Baseline	Week 6	Week 12
n (missing)	268 (0)	253 (15)	237 (31)
0	93 (34.7%)	91 (36.0%)	89 (37.6%)
+0.5	80 (29.9%)	91 (36.0%)	94 (39.7%)
+1	66 (24.6%)	51 (20.2%)	40 (16.9%)
+2	25 (9.3%)	19 (7.5%)	13 (5.5%)
+3	4 (1.5%)	1 (0.4%)	1 (0.4%)

Table 4 Rates of ocular hyperemia by prior intraocular pressure-lowering therapy

Prior intraocular pressure-lowering therapy	Number (%) of subjects with none-to-mild hyperemia/moderate-to-severe hyperemia					
	Baseline		Week 6		Week 12	
	None-to-mild	Moderate-to-severe	None-to-mild	Moderate-to-severe	None-to-mild	Moderate-to-severe
Latanoprost 0.005%	108 (95.6)	5 (4.4)	100 (94.3)	6 (5.7)	93 (94.9)	5 (5.1)
Bimatoprost 0.03%	45 (71.4)	18 (28.6)	52 (89.7)	6 (10.3)	49 (92.5)	4 (7.5)
Travoprost 0.004%	13 (81.3)	3 (18.8)	14 (87.5)	2 (12.5)	15 (93.8)	1 (6.3)
Travoprost 0.004% with SofZia™	33 (91.7)	3 (8.3)	33 (91.7)	3 (8.3)	32 (91.4)	3 (8.6)
Nonselective beta-adrenergic receptor blockers	40 (100.0)	0 (0.0)	34 (91.9)	3 (8.1)	34 (97.1)	1 (2.9)

Notes: None-to-mild hyperemia = grades 0, +0.5, or +1; moderate-to-severe hyperemia = grades +2 or +3. Bimatoprost (Lumigan®, Allergan, Inc., Irvine, CA, USA); Latanoprost (Xalatan®, Pfizer, Inc., New York, NY, USA); Travoprost (Travatan®, Alcon Laboratories, Inc., Fort Worth, TX, USA); Travoprost with SofZia™ (Travatan Z®, Alcon Laboratories, Inc.).

respectively. These changes in ocular hyperemia grade observed at both study visits were statistically significant ($P=0.013$ and $P=0.004$ at week 6 and week 12, respectively). Analysis of ocular hyperemia following the switch to bimatoprost 0.01% from the other prior IOP-lowering therapies (travoprost 0.004% original version preserved with BAK, travoprost 0.004% preserved with SofZia™, and nonselective beta-adrenergic receptor blockers) failed to reveal any statistically significant changes at the week 6 and week 12 study visits (Table 5).

Intraocular pressure

The mean IOP at baseline and weeks 6 and 12 and the mean percentage changes in IOP from baseline following switching to bimatoprost 0.01% at week 6 and week 12 are shown in Table 6. The additional decrease in mean percentage change in IOP achieved by switching to bimatoprost 0.01% monotherapy was statistically significant at week 6 and week 12 for all prior IOP-lowering therapies, except bimatoprost 0.03% at week 6 and week 12 and travoprost 0.004% (original version preserved with BAK) at week 6 (Table 6).

Safety

Thirteen subjects reported 22 AEs, for an overall incidence of 4.9% (13 of 268). A total of 19 AEs reported by eleven subjects were considered related to bimatoprost 0.01%, for

an incidence of 4.1% (eleven of 268). The most frequent reported treatment-related AEs associated with bimatoprost 0.01% were ocular hyperemia (1.1%), eye irritation (0.7%), and skin hyperpigmentation (0.7%). No serious AEs were observed in this study population.

Discussion

An ideal first-line IOP-lowering treatment for glaucoma and/or OHT should provide superior IOP-lowering efficacy with minimal side effects. In Canada, PGAs and PMs are recognized as first-line agents because of their superior efficacy and infrequent systemic side effects compared to other classes of topical ocular hypotensive therapies.¹ Nonetheless, the most common AE experienced by PGA and PM users is hyperemia, a nonspecific clinical term implying vasodilation and increased blood supply to the conjunctival blood vessels that has been shown to be an important factor affecting patient adherence.^{2,9} While bimatoprost has been shown to have a unique pharmacology with respect to lowering IOP, bimatoprost and the PGAs are believed to share a common noninflammatory signaling mechanism leading to vasodilation.^{9–11}

In the present study, the switch to bimatoprost 0.01% from prior IOP-lowering medication resulted in 94.1% and 5.9% of subjects presenting with none-to-mild and moderate-to-severe hyperemia, respectively, after 12 weeks.

Table 5 Change in ocular hyperemia following switch from prior intraocular pressure-lowering therapy

	Week 6, improved	Week 6, no change	Week 6, worsened	P-value	Week 12, improved	Week 12, no change	Week 12, worsened	P-value
Latanoprost	0.0%	96.2%	3.8%	0.046	1.0%	94.9%	4.1%	0.180
Bimatoprost 0.03%	19.0%	77.6%	3.4%	0.013	20.8%	77.4%	1.9%	0.004
Travoprost 0.004%	12.5%	81.3%	6.3%	0.564	12.5%	87.5%	0.0%	0.157
Travoprost 0.004% with SofZia™	8.3%	83.3%	8.3%	1.000	8.6%	82.9%	8.6%	1.000
Nonselective beta-adrenergic receptor blockers	0.0%	91.9%	8.1%	0.083	0.0%	97.1%	2.9%	0.317

Notes: Bimatoprost (Lumigan®, Allergan, Inc., Irvine, CA, USA); Latanoprost (Xalatan®, Pfizer, Inc., New York, NY, USA); Travoprost (Travatan®, Alcon Laboratories, Inc., Fort Worth, TX, USA); Travoprost with SofZia™ (Travatan Z®, Alcon Laboratories, Inc.).

Table 6 Intraocular pressure (IOP) values by prior IOP-lowering therapy

Prior IOP-lowering Therapy	Baseline	Week 6			Week 12		
	Mean IOP, mmHg $\pm$ SD	Mean IOP, mmHg $\pm$ SD	Mean % change in IOP $\pm$ SD	P-value*	Mean IOP, mmHg $\pm$ SD	Mean % change in IOP $\pm$ SD	P-value*
Latanoprost 0.005%	19.8 $\pm$ 3.7 (n=113)	16.8 $\pm$ 4.1 (n=106)	-15.1 $\pm$ 14.5 (n=106)	<0.001	16.7 $\pm$ 4.3 (n=98)	-15.2 $\pm$ 15.6 (n=98)	<0.001
Bimatoprost 0.03%	17.9 $\pm$ 3.6 (n=63)	17.0 $\pm$ 3.7 (n=58)	-3.4 $\pm$ 16.0 (n=58)	0.114	17.4 $\pm$ 4.3 (n=53)	-2.3 $\pm$ 17.3 (n=53)	0.342
Travoprost 0.004%	20.8 $\pm$ 4.5 (n=16)	18.4 $\pm$ 3.5 (n=16)	-8.9 $\pm$ 19.9 (n=16)	0.094	18.0 $\pm$ 3.7 (n=16)	-12.5 $\pm$ 14.4 (n=16)	0.004
Travoprost 0.004% with SofZia™	19.9 $\pm$ 4.0 (n=36)	18.0 $\pm$ 4.2 (n=36)	-9.4 $\pm$ 12.2 (n=36)	<0.001	17.8 $\pm$ 3.3 (n=35)	-9.5 $\pm$ 14.0 (n=35)	<0.001
Nonselective beta-adrenergic receptor blockers	21.6 $\pm$ 4.1 (n=40)	17.1 $\pm$ 3.9 (n=37)	-20.6 $\pm$ 14.8 (n=37)	<0.001	15.9 $\pm$ 3.6 (n=35)	-26.3 $\pm$ 12.4 (n=35)	<0.001

Notes: *P-values calculated for % change from baseline IOP. Bimatoprost (Lumigan®, Allergan, Inc., Irvine, CA, USA); Latanoprost (Xalatan®, Pfizer, Inc., New York, NY, USA); Travoprost (Travatan®, Alcon Laboratories, Inc., Fort Worth, TX, USA); Travoprost with SofZia™ (Travatan Z®, Alcon Laboratories, Inc.).

Abbreviation: SD, standard deviation.

Of the prior IOP-lowering treatments, bimatoprost 0.03% was associated with the greatest percentage of subjects presenting with moderate-to-severe hyperemia at baseline (28.6%). Following the switch from bimatoprost 0.03% to bimatoprost 0.01%, the occurrence of moderate-to-severe hyperemia was reduced to 10.3% and 7.5% at week 6 and week 12, respectively. These results suggest that bimatoprost 0.01% was well tolerated following the switch from prior IOP-lowering medications, a finding that corroborates others.¹²⁻¹⁴ Additionally, our findings concerning the occurrence and severity of ocular hyperemia following the switch to bimatoprost 0.01% correlate well with the randomized, controlled trial results reported by Katz et al, which demonstrated a threefold-lower incidence of moderate-to-severe hyperemia in the bimatoprost 0.01% group compared to the bimatoprost 0.03% group.³

In terms of IOP-lowering efficacy, a number of studies have now been published comparing various PGAs and PMs, including latanoprost 0.005% with travoprost 0.004% and bimatoprost 0.03%. In 2003, a multicenter study involving 411 patients demonstrated that latanoprost 0.005%, travoprost 0.004%, and bimatoprost 0.03% were comparable in their ability to lower IOP, although differences were found in associated rates of conjunctival hyperemia.¹⁵ A recent observational study found that subjects who were switched to bimatoprost 0.01% from either latanoprost 0.005%, travoprost 0.004% preserved with 0.015% BAK (original version), tafluprost, or bimatoprost 0.03% achieved an additional IOP decrease of 2.8 mmHg, 3.1 mmHg, 2.8 mmHg, or 1.0 mmHg, respectively.¹⁶ It should be noted that for both the report by Pfennigsdorf et al¹⁶ and the present analysis, travoprost preserved with BAK was available in Europe and Canada, as well as in Argentina, Brazil,

Chile, China, Colombia, India, Malaysia, Mexico, Taiwan, and Venezuela. As a result, the total number of patients who switched to bimatoprost 0.01% from prior travoprost preserved with BAK was low relative to other prior IOP-lowering therapies included in our analysis, due to the discontinuation of the original version of travoprost partway through study enrollment. A unique aspect of the present report is the inclusion of safety and efficacy data for the switch from prior travoprost preserved with SofZia™ to bimatoprost 0.01%, which demonstrated no statistically significant changes in hyperemia grade along with a -9.5% $\pm$ 14.0% change in IOP ($P<0.001$) at week 12.

In the present study, all of the patient subgroups on prior IOP-lowering therapies showed statistically significant improvements in IOP following the switch to bimatoprost 0.01% at week 12, except for the subgroup initially treated with bimatoprost 0.03%. The subgroup of patients previously treated with bimatoprost 0.03% was associated with the lowest IOP at baseline, which may explain why the IOP reductions following the switch to bimatoprost 0.01% were not statistically significant at the week 6 and 12 visits. Likely, the hyperemia side effect profile of bimatoprost 0.03% was the reason the investigators decided to switch the patient's IOP-lowering regimen to bimatoprost 0.01%.

Some limitations of the present study should be acknowledged. Due to the open-label design of this study, the results presented may have been influenced by observer bias by the participating physician in the subjective grading of ocular hyperemia and the measurement of IOP, since neither the physician nor patients were masked to study treatment. The present study did not include a washout period between treatments to more closely resemble real-world practice. By not including a washout period, it is difficult to discern rates of

ocular hyperemia caused by bimatoprost 0.01% and the prior IOP-lowering therapy. With 64 examiners involved in this study, we acknowledge the opportunity for interexaminer variability with hyperemia and IOP measurements. A Hawthorne effect for enhanced patient adherence to the study therapy may have contributed to the observed decrease in IOP when patients switched from prior therapy. Subject dropout can also affect a study's results. Thirty-two of the 268 subjects discontinued for the following reasons: lost to follow-up, $n=13$ (4.9%); ocular AE, $n=5$ (1.9%); other AE, $n=2$ (0.7%); and reasons not related to AEs, $n=12$ (4.5%). Of the five monotherapy-switch subjects who discontinued the study due to ocular AEs, a total of nine ocular AEs were related to bimatoprost 0.01% monotherapy (Table 2). Exclusion of subjects with ocular AEs may have understated the primary outcome of this study, namely occurrence and severity of hyperemia attributed to a switch from prior IOP-lowering therapy to bimatoprost 0.01% monotherapy. Lastly, regression to the mean happens when a reduction in IOP occurs spontaneously rather than therapeutically. Since the present study did not include a second baseline IOP measurement prior to enrollment selection, bias causing a regression to the mean may have partially contributed to the significant IOP reductions observed.

Overall, the results of this observational study provide real-world clinical evidence to support bimatoprost 0.01% as a suitable treatment option for POAG and OHT subjects. Additionally, further reduction in IOP may be achieved when switching from a prior IOP-lowering monotherapy to bimatoprost 0.01%.

Acknowledgments

The authors acknowledge the editorial assistance of Mary Ann Chapman, PhD and Jaspreet Grewal MSc, COMT, CCRP. This study was supported by Allergan Inc. The CLEAR Study Group: Iqbal Ike K Ahmed, Mississauga, ON; Rama Behki, Gatineau, QC; Robert Beldavs, Winnipeg, MB; Francois Bellefeuille, Trois-Rivieres, QC; Dan Belliveau, Halifax, NS; Raj Bindlish, Oakville, ON; Fadi Calotti, Brantford, ON; Jeff Chambers, Kelowna, BC; Jason Cherry, Richmond, BC; Robert Chevrier, Gloucester, ON; Alan Coffey, Westmount, QC; Andrew C Crichton, Calgary, AB; Marino J Discepola, Montreal, QC; Gordon Douglas, Calgary, AB; Stephane Dupont, Gramby, QC; Chaim Edelstein, Montreal, QC; Stephen Fichman, Montreal, QC; Bryce Ford, Calgary, AB; Michel Giunta, Sherbrooke, QC; Sheldon Goldhar, Scarborough, ON; Larry Green, Sidney, BC; Francine Guay, Lévis, QC; Eugene Hladky, Montreal, QC; Glen Hoar, Comox, BC;

Muhammad Humayun, Dartmouth, NS; Cindy ML Hutnik, London, ON; Norman Hwang, Burnaby, BC; Fahim Ibrahim, Point Edward, ON; Carolyn Isbister, West Vancouver, BC; Helene Joyal, St-Jean-Sur-Richelieu, QC; Barry Kattleman, Montreal, QC; Baseer Khan, Vaughan, ON; Elie Khouri, Saint-Laurent, QC; Thomas Klein, Mississauga, ON; Alain Lachance, Saint-Georges, QC; Hesham Lakosha, Halifax, NS; Alain Lamoureux, Sorel, QC; Gisele Li, Montreal, QC; Derek Lui, Woodstock, ON; Mark Lukasik, Owen Sound, ON; Sam Markowitz, Toronto, ON; Jean Mayer, St-Charles Borromeo, QC; Mohan Merchea, London, ON; Michael Miller, Mississauga, ON; Kam Mohaseb, North Vancouver, BC; Paul Murphy, Saskatoon, SK; Rashmi Nigam, Winnipeg, MB; Donald R Nixon, Barrie, ON; Patrick O'Keeffe, North Battleford, SK; Manuel Perrier, Lachine, QC; Carl Peters, Penticton, BC; Theodore Rabinovitch, Downsview, ON; Aaron Rifkind, Hamilton, ON; Suleiman Sefau, Charlottetown, PEI; Rob Semeniuk, Penticton, BC; Lesya Shuba, Halifax, NS; Barry Silver, Dorval, QC; Zakaria Tadrous, Abbotsford, BC; Eric Tam, Brampton, ON; Alexander Tan, Halifax, NS; Andrew Taylor, Niagara Falls, ON; Patricia Teal, Fort Erie, ON; David B Yan, Mississauga, ON; Natasha Yepes, Kitchener, ON, Canada.

Disclosure

Publication of this article was sponsored by Allergan Inc. Susan Simonyi, Meetu Bhogal, and Darryl C Baptiste are employees of Allergan Inc. Christopher S Sigouin is an employee of ClinWest, which received fees for assisting with statistical analysis. Andrew C Crichton, Donald R Nixon, Marino J Discepola, Cindy ML Hutnik and David B Yan declare no potential conflicts of interest in this work.

References

1. Canadian Ophthalmological Society Glaucoma Clinical Practice Guideline Expert Committee. Canadian Ophthalmological Society evidence-based clinical practice guidelines for the management of glaucoma in the adult eye. *Can J Ophthalmol*. 2009;44 Suppl 1:S7–S93.
2. Zimmerman TJ, Hahn SR, Gelb L, Tan H, Kim EE. The impact of ocular adverse effects in patients treated with topical prostaglandin analogs: changes in prescription patterns and patient persistence. *J Ocul Pharmacol Ther*. 2009;25:145–152.
3. Katz LJ, Cohen JS, Batoosingh AL, Felix C, Shu V, Schiffman RM. Twelve-month, randomized, controlled trial of bimatoprost 0.01%, 0.0125%, and 0.03% in patients with glaucoma or ocular hypertension. *Am J Ophthalmol*. 2010;149:661–671. e1.
4. Aptel F, Cucherat M, Denis P. Efficacy and tolerability of prostaglandin analogs: a meta-analysis of randomized controlled clinical trials. *J Glaucoma*. 2008;17:667–673.
5. Noecker RS, Dirks MS, Choplin NT, Bernstein P, Batoosingh AL, Whitcup SM. A six-month randomized clinical trial comparing the intraocular pressure-lowering efficacy of bimatoprost and latanoprost in patients with ocular hypertension or glaucoma. *Am J Ophthalmol*. 2003;135:55–63.

6. Goldberg I, Li XY, Selaru P, Paggiarino D. A 5-year, randomized, open-label safety study of latanoprost and usual care in patients with open-angle glaucoma or ocular hypertension. *Eur J Ophthalmol*. 2008;18:408–416.
7. Law SK, Song BJ, Fang E, Caprioli J. Feasibility and efficacy of a mass switch from latanoprost to bimatoprost in glaucoma patients in a prepaid Health Maintenance Organization. *Ophthalmology*. 2005;112: 2123–2130.
8. Nixon DR, Simonyi S, Bhogal M, et al. An observational study of bimatoprost 0.01% in treatment-naïve patients with primary open angle glaucoma or ocular hypertension: the CLEAR Trial. *Clin Ophthalmol*. 2012;6:2097–2103.
9. Chen J, Dinh T, Woodward DF, et al. Bimatoprost: mechanism of ocular surface hyperemia associated with topical therapy. *Cardiovasc Drug Rev*. 2005;3:231–246.
10. Woodward DF, Krauss AH, Chen J, et al. The pharmacology of bimatoprost (Lumigan). *Surv Ophthalmol*. 2001;45 Suppl 4:S337–S345.
11. Astin M, Stjernschantz J, Selén G. Role of nitric oxide in PGF2 α -induced ocular hyperemia. *Exp Eye Res*. 1994;59:401–408.
12. Kurtz S, Mann O. Incidence of hyperemia associated with bimatoprost treatment in naïve subjects and in subjects previously treated with latanoprost. *Eur J Ophthalmol*. 2009;19:400–403.
13. Abelson MB, Mroz M, Rosner SA, et al. Multicenter, open-label evaluation of hyperemia associated with use of bimatoprost in adults with open-angle glaucoma or ocular hypertension. *Adv Ther*. 2003;20: 1–13.
14. Craven ER, Liu CC, Batoosingh A, Schiffman RM, Whitcup SM. A randomized, controlled comparison of macroscopic conjunctival hyperemia in patients treated with bimatoprost 0.01% or vehicle who were previously controlled on latanoprost. *Clin Ophthalmol*. 2010;4: 1433–1440.
15. Parrish RK, Palmberg P, Sheu WP, XLT Study Group. A comparison of latanoprost, bimatoprost, and travoprost in patients with elevated intraocular pressure: a 12-week, randomized, masked-evaluator multicenter study. *Am J Ophthalmol*. 2003;135:688–703.
16. Pfennigsdorf S, Ramez O, von Kistowski G, et al. Multicenter, prospective, open-label, observational study of bimatoprost 0.01% in patients with primary open-angle glaucoma or ocular hypertension. *Clin Ophthalmol*. 2012;6:739–746.

Clinical Ophthalmology

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on

Submit your manuscript here: <http://www.dovepress.com/clinical-ophthalmology-journal>

Dovepress

PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.