

Association between dentures and the rate of falls in dementia

Sima Ataollahi Eshkoor¹
Tengku Aizan Hamid¹
Siti Sa'adiah Hassan Nudin²
Chan Yoke Mun¹

¹Institute of Gerontology, Universiti Putra Malaysia, Serdang, ²Institute for Behavioral Research, Kuala Lumpur, Malaysia

Background: Poor oral health, chronic diseases, functional decline, and low cognitive ability can increase the risk of falls in the elderly.

Objectives: The current study aimed to show the effects of oral health, diabetes mellitus (DM), hypertension (HT), heart disease, functional status, and sociodemographic factors on the risk of falls in elderly with dementia.

Materials and methods: The sample comprised 1,210 Malaysian elderly who were demented and noninstitutionalized. This study was a national cross-sectional survey entitled "Determinants of Health Status among Older Malaysians". The effects of age, ethnicity, sex differences, marital status, educational level, oral health, DM, HT, heart disease, and functional status on the risk of falls were evaluated. The multiple logistic regression model was used to estimate the effects of contributing variables on the risk of falls in samples.

Results: The prevalence of falls was approximately 17% in subjects. It was found that age (odds ratio [OR] 1.02), non-Malay ethnicity (OR 1.66), heart disease (OR 1.92), and functional decline (OR 1.58) significantly increased the risk of falls in respondents ($P < 0.05$). Furthermore, having teeth (OR 0.59) and dentures (OR 0.66) significantly decreased the rate of falls ($P < 0.05$).

Conclusion: It was concluded that age, non-Malay ethnicity, functional decline, heart disease, and oral health significantly affected falls in dementia.

Keywords: chronic diseases, dementia, fall, functional decline, oral health

Introduction

Dementia is the most common disabling disease in older people.¹ There are different types, such as Alzheimer's disease (AD), vascular dementia, dementia with Lewy bodies, and frontotemporal dementia.² Falls are common in the elderly,³⁻⁵ and can occur due to the negative effects of medications, medical conditions, poor nutrition, psychological problems, physical impairment, sensory disturbances, and cognitive impairment.⁶ Dementia increases the risk of falls in the elderly⁷ where cognitive decline^{8,9} impairs judgment, gait, and visuospatial perception, reducing the ability to recognize and avoid hazards.⁷ Furthermore, older people often suffer from chronic illnesses, including hypertension (HT), diabetes mellitus (DM), and vascular diseases.¹⁰ DM,¹¹⁻¹⁴ HT,¹⁵ and heart disease¹⁶ may increase the risk of falls in the elderly. DM is one of the most common chronic diseases in elderly¹² presenting with glucose intolerance.¹⁷ The prevalence and morbidities associated with DM are on the rise among older adults. The increased risk of falls in diabetic patients is due to dehydration,¹³ retinopathy, peripheral artery disease, peripheral polyneuropathy, cerebral vasculopathy, autonomic neuropathy, and poor muscle quality.¹¹

Correspondence: Tengku Aizan Hamid
Institute of Gerontology,
Universiti Putra Malaysia, Serdang,
Selangor 43400, Malaysia
Email aizan@upm.edu.my

Age can adversely influence the heart in older adults. Age-related changes in the cardiovascular system can occur due to the intrinsic cardiac aspects of senescence, primary cardiac disease, and the effects of comorbid conditions.¹⁸ The effects of cardiovascular problems on the greater risk of falls in the elderly are associated with intrinsic cardiac abnormalities (structure, rhythms), neurally mediated disorders, and some miscellaneous factors, such as pulmonary embolism and transient ischemic attacks.¹⁶ HT is also a condition that can strongly increase the rate of cardiovascular diseases,¹⁹ which consequently enhance the risk of falls in the elderly.¹⁵

A higher risk of falls in the elderly can be related to mobility dysfunction and balance problems. Functional decline influences mobility and balance skills in terms of time taken to complete the task measured by Get Up and Go Test (GUGT).²⁰ Oral health is also an important component of overall health, well-being, and quality of life in older individuals.²¹ Poor oral functioning can lead to systemic illnesses and life-threatening conditions in older adults.²² There is a positive relation between dental problems and the risk of falls among the elderly with dementia. Dental problems can affect postural control, which in turn increases the risk of falls.²³ Since the increased risk of falls is associated with morbidity, disability, social isolation, and lower quality of life in the elderly,³ the evaluation of correspondence factors has become a focus. This study aimed to investigate the effects of oral health, chronic diseases, functional status, and sociodemographic factors on the risk of falls in elderly with dementia.

Materials and methods

This project was registered in the National Medical Research Register (project code NMRR-09-443-4148). Approval and permission for conducting the study were received from the Ethical Committee of the Ministry of Health. This project was a national cross-sectional survey design entitled “Determinants of Health Status among Older Malaysians” carried out in cooperation with the Health Behavioral Research Institute, National Institute of Health, Ministry of Health, and Institute of Gerontology, Universiti Putra Malaysia.

The research involved 1,210 elderly with dementia who were Malaysian aged 60 years and above residing in noninstitutional places. The elderly living in institutions and bedridden were excluded. Samples represented the Malaysian population in terms of age, and were collected from Peninsular Malaysia, which was divided into four zones

of north, south, west, and central. Data were collected by trained interviewers who had prior experience as interviewers in projects conducted by the Institute of Gerontology. The average duration of interviews was about 60 minutes. In this study, the association between age, ethnicity, sex differences, marital status, educational level, DM, HT, heart disease, oral health, and functional status with the risk of falls was evaluated in respondents. Falls were defined according to the *International Classification of Diseases* (ICD-9),²⁴ which excluded falls due to loss of consciousness, epileptic seizure, and sudden onset of paralysis, eg, stroke. History of falls referred to having had at least one fall during the day or night during the past 6 months. Subjects who had a Mini-Mental State Examination (MMSE) mark less than 26 were considered to have dementia.²⁵ The MMSE is a clinical diagnostic tool used for both research and clinical practice in neurology and psychiatry. It is also one of the most widely used screening and outcome measures in cognition assessments.²⁶

Functional status was measured by the GUGT for rising from a chair, walking 3 meters, turning, and returning to the chair. The GUGT is a reliable and validated instrument used to test balance and gait.^{4,27} Functional decline was defined as a time of completion of more than 20 seconds.²⁸ In the current study, the measurement of oral health was based on a self-report of having teeth and dentures.²⁹ DM, HT, and heart disease were assessed by physicians.

Statistical analysis

The prevalence of falls was computed for total samples with regard to their age, ethnicity, sex, marital status, educational level, oral health, functional status, DM, HT, and heart disease. Bivariate analysis was carried out to examine the relation between each variable and the risk of falls using a series of χ^2 tests. Multivariate logistic regression analysis was used to test the association between age, ethnicity, educational level, marital status, oral health, functional status, DM, HT, heart disease, and the risk of falls in elderly with dementia. Odds ratios (ORs) with 95% confidence intervals (CIs) were computed. The critical level for rejection of the null hypothesis was considered to be a *P*-value of 5%, two-tailed. All analyses were performed using SPSS version 20.0 (IBM, Armonk, NY, USA).

Results

Analysis was run on data collected from 1,210 respondents who were elderly Malaysians with dementia. The mean age of respondents was 71 years (minimum 60±7.38 and maximum

103±7.38 years). The prevalence of falls was 17% among subjects (95% CI 15.01–19.24) (Table 1). The results indicated that the risk of falls in respondents with DM (20.4%) was higher than those without DM (16.3%). Subjects with heart disease (27.4%) experienced further risk of falls compared to those without heart disease (16.2%). The risk of falls in HT samples (18.9%) was higher than normotensive samples (15.7%). The results showed that the prevalence of falls was 22.9% in subjects with functional decline and 13.6% in the subjects with normal functional status. The findings indicated that respondents without teeth (18.7%) and dentures (18.6%) had a higher fall rate compared to those having teeth (10.9%) and dentures (14.4%). It was found that the risk of falls in females (18.3%) was higher than males (14.8%). In addition, the rate of falls was 14.7% in respondents with education and 18.5% in those with no education.

The prevalence of falls was found to be as high as 19.4% in single subjects and 14.2% in married subjects. Among the samples, 13.3% of Malay and 20.2% of non-Malay ethnicities experienced falls. A series of χ^2 tests evaluated the association of each variable with the risk of falls in samples. The findings indicated that ethnicity ($\chi^2=10.20$, $P=0.001$), heart disease ($\chi^2=7.78$, $P=0.006$), functional decline ($\chi^2=14.44$, $P=0.001$), marital status ($\chi^2=5.75$, $P=0.010$), and having teeth ($\chi^2=8.50$, $P=0.002$) or dentures ($\chi^2=3.60$, $P=0.034$) were significantly associated with the risk of falls among respondents ($P<0.05$). The results indicated that DM, HT, educational level, and sex differences were unrelated to the risk of falls ($P>0.05$) (Table 2). The logistic regression analysis showed that age ($P=0.041$), ethnicity ($P=0.003$), heart disease ($P=0.011$), functional decline ($P=0.033$), and having teeth ($P=0.021$) or dentures ($P=0.021$) were significantly associated with the risk of falls in samples. It was found that age (OR 1.02, 95% CI 1.00–1.05), non-Malay ethnicity (OR 1.66, 95% CI 1.19–2.33), functional decline (OR 1.58, 95% CI 1.04–2.41), and heart disease (OR 1.92, 95% CI 1.16–3.19) significantly increased the risk of falls in subjects ($P<0.05$). The results showed that DM, HT, sex differences, marital status, and educational level were not significant predictors for the risk of falls in subjects ($P>0.05$). Having teeth (OR 0.58, 95% CI 0.37–0.91) or dentures (OR 0.66, 95% CI 0.47–0.94) significantly reduced the risk of falls (Table 3).

Table 1 Prevalence of falls among 1,210 elderly with dementia

Falls	n	n (%)	95% CI
Yes	206	17	15.01–19.24
No	1,004	83	80.76–84.99

Abbreviation: CI, confidence interval.

Table 2 Prevalence of falls and associations with sociodemographic factors

	Total	n ^a	n%	95% CI	χ^2	P-value*
Diabetes mellitus						
No	993	162	16.3	14.14–18.74	2.07	0.092
Yes	216	44	20.4	15.54–26.24		
Heart disease						
No	1,114	180	16.2	14.12–18.44	7.78	0.006
Yes	95	26	27.4	19.42–37.08		
Hypertension						
No	695	109	15.7	13.17–18.57	2.13	0.084
Yes	514	97	18.9	15.72–22.48		
Functional test						
<20 seconds**	559	76	13.6	11.01–16.69	14.44	0.001
≥20 seconds	389	89	22.9	18.98–27.31		
Teeth						
No	949	177	18.7	16.3–21.25	8.50	0.002
Yes	256	28	10.9	7.68–15.36		
Dentures						
No	753	140	18.6	15.97–21.53	3.60	0.034
Yes	451	65	14.4	11.47–17.95		
Teeth and dentures						
No	535	114	21.31	18.05–24.98	12.46	<0.001
Yes	675	92	13.63	11.25–16.43		
Sex differences						
Male	438	65	14.8	11.82–18.47	2.32	0.074
Female	772	141	18.3	15.69–21.14		
Marital status						
Single	665	129	19.4	16.57–22.58	5.75	0.010
Married	543	77	14.2	11.5–17.37		
Ethnicity						
Malay**	550	73	13.3	10.69–16.36	10.20	0.001
Non-Malay	658	133	20.2	17.32–23.45		
Education						
No	775	143	18.5	15.88–21.33	2.82	0.054
Yes	430	63	14.7	11.62–18.31		

Notes: *Significant at the 0.05 level using the χ^2 test; **reference group; ^anumber of subjects who reported at least one fall. Missing data reduced the number of all subjects in some variables.

Abbreviation: CI, confidence interval.

Discussion

As falls can cause much burden and cost in patients with dementia and their caregivers,^{30,31} the assessment of risk factors has been a major focus in further investigations. It has been well documented that age-related changes in the body,³² DM,^{11–14} HT,¹⁵ heart disease,^{3,16,33} functional status,²⁰ and oral health all affect the risk of falls in the elderly.

This study was designed to identify the correspondence of age, ethnicity, sex differences, marital status, educational level, functional status, DM, HT, heart disease, and having teeth or dentures with the rate of falls among Malaysian elderly with dementia. In the current study, age, ethnicity, heart disease, functional decline, and having teeth or dentures significantly affected the risk of falls. This project studied

Table 3 Prevalence of falls and associations derived by logistic regression analysis

	B	SE	P-value*	OR	95% CI for OR	
					Lower	Upper
Diabetes mellitus	0.245	0.208	0.238	1.28	0.85	1.92
Heart disease	0.654	0.258	0.011	1.92	1.16	3.19
Hypertension	0.147	0.168	0.382	1.16	0.83	1.61
Functional test	0.457	0.215	0.033	1.58	1.04	2.41
Teeth	-0.531	0.230	0.019	0.58	0.37	0.91
Dentures	-0.409	0.177	0.022	0.67	0.47	0.94
Teeth and dentures	-0.423	0.162	0.009	0.66	0.48	0.90
Sex differences	-0.225	0.185	0.225	0.80	0.56	1.15
Educational level	0.046	0.185	0.803	1.05	0.73	1.50
Marital status	-0.104	0.180	0.562	0.90	0.63	1.28
Ethnicity	0.507	0.172	0.003	1.66	1.19	2.33
Age	0.024	0.012	0.041	1.02	1.00	1.05

Notes: *Significant at the 0.05 level using the logistic regression analysis. Hosmer–Lemeshow test, $\chi^2_6=6.56$; $P=0.584$.

Abbreviations: SE, standard error; OR, odds ratio; CI, confidence interval.

the effect of heart disease on the risk of falls in the elderly with dementia for the first time. Heart disease increased the risk of falls in subjects, which was probably due to medication use,^{34,35} sleep problems,^{36–38} and cardiac arrhythmias.³³ Medical conditions like postural hypotension as a side effect of medications and malnutrition can cause a higher rate of falls in the elderly with heart problems.³⁴ The association of heart disease with the impairment of cognition, disability, and gait disorders³⁹ potentially increased the rate of falls as well. The increased rate of falls in the elderly with DM and HT can be explained by the association of these problems with cardiovascular diseases.⁴⁰ This study was in line with other reports^{8,27,41} documenting functional decline as a risk factor for further falls in the elderly. It seems that balance problems, insufficient mobility, cognitive impairment,^{8,27} medications, orthostatic blood pressure, physical problems, psychological conditions, and vision problems⁸ enhance the risk of falls in elderly with dementia.

The increased rate of falls in samples without teeth or dentures was possibly due to the profound effects of teeth or dentures on systemic health.²¹ It has been established that having teeth or dentures affects nutrition, medical situations,^{42,43} social interactions, and morbidity rates.⁴² Therefore, the correlation between oral health and the risk of falls may depend on disturbance levels in food intake, masticatory ability, and behavioral and psychological status.⁴⁴

The results showed that DM and HT had no significant effects on the risk of falls in respondents, which is probably associated with the confounding effects of heart disease, functional decline, and sociodemographic factors.

Contrary to some reports,^{45–47} our study showed that ethnicity could affect the risk of falls.^{7,8,48} Such a discrepancy was possibly due to differences in lifestyle, cultures, and predisposition.⁴⁹ Indeed, lifestyle, cultural, economic, and environmental factors in a specific ethnic group are related to expression of variances in such diseases as hypertension, stroke, and ischemic heart disease, which in turn may enhance or attenuate the risk of falls.

Our study was a confirmation of previous reports indicating the presence of a link between age and the risk of falls.^{7,50–52} Such a relation can be due to age-related changes in gait, vision, memory, strength and physiological systems,^{8,48,53,54} which potentially predispose the elderly with dementia to a higher risk of falls. The findings showed that marital status, educational level, and sex differences were unrelated to the risk of falls in respondents. It is important to note that our results confirmed an earlier finding⁵⁵ contrary to reports indicating the effects of sex differences on the risk of falls.^{7,8,56} Our study did not confirm the presence of correlation between the risk of falls and marital status⁷ or educational level.^{7,8,53,57} The lack of such relevance can be related to the possible effects of confounding factors, such as social supports,⁵⁸ income,⁵⁹ supplement intake,⁸ sleep problems,^{30,60,61} environmental conditions,^{62,63} functional decline, heart problems, and oral health, which may affect the rate of falls. These factors may correlate with the physical, psychological, and cognitive changes in the patients with dementia. Therefore, the reduction of attention, cognition, mood, and motor performances can increase the risk of falls in dementia. An appropriate environment helps to overcome deficits in physical fitness and cognitive abilities posed by dementia, and thus reduces the risk of falls in elderly patients with dementia. Therefore, these factors can interact between physical activity and cognition when considering the risk of falls in respondents.⁵

Conclusion

This study faced some limitations. One problem was related to difficulty in collecting accurate self-reported data about the rate of falls and oral health from elderly with dementia, which can cause bias. Furthermore, the higher prevalence of comorbidities in these patients can limit such studies. The cross-sectional design of the study was another restriction factor that made it difficult to find the exact effective and noneffective factors. Accordingly, more standardized interviews and definitions are needed for further investigations about falls in dementia. However, this study could highlight some risk factors contributing to the risk of falls in elderly with dementia. It was concluded

that age, non-Malay ethnicity, heart disease, and functional decline increased the risk of falls in respondents. On the other hand, having teeth or dentures decreased the likelihood of falls. As falls can cause much burden, morbidity, and mortality among those patients, more studies are required to find ways to prevent and reduce the rate of falls. Furthermore, such data can help caregivers to reduce the risk of falls and unhappiness in such patients. In addition, this knowledge can provide an opportunity for policy makers and health care professionals to introduce suitable interventions in order to control the rate of falls among elderly with dementia.

Acknowledgments

The authors gratefully acknowledge the cooperation of all volunteers who participated in this study. The authors wish to acknowledge the financial support from the Ministry of Health for this study. In addition, the authors also thank all staff for their efforts in data collection.

Disclosure

The authors report no conflicts of interest in this work.

References

- Nikmat AW, Hawthorne G, Al-Mashoor S. Dementia in Malaysia: issues and challenges. *ASEAN J Psychiatry*. 2011;12:1–7.
- Kalapatapu RK. Dementia: a focused review. *Psychiatr Times*. 2010;27:1–12.
- Lawlor DA, Patel R, Ebrahim S. Association between falls in elderly women and chronic diseases and drug use: cross sectional study. *BMJ*. 2003;327:712–717.
- Hendriks MR, Bleijlevens MH, van Haastregt J, et al. A multidisciplinary fall prevention program for elderly persons: a feasibility study. *Geriatr Nurs*. 2008;29:186–196.
- Eshkoor SA, Hamid TA, Nudin SS, Mun CY. The effects of sleep quality, physical activity, and environmental quality on the risk of falls in dementia. *Am J Alzheimers Dis Other Dement*. 2013;28:403–407.
- Sirkin AJ, Rosner NG. Hypertensive management in the elderly patient at risk for falls. *J Am Acad Nurse Pract*. 2009;21:402–408.
- van Doorn C, Gruber-Baldini AL, Zimmerman S, et al. Dementia as a risk factor for falls and fall injuries among nursing home residents. *J Am Geriatr Soc*. 2003;51:1213–1218.
- Rubenstein LZ, Josephson KR. Falls and their prevention in elderly people: what does the evidence show? *Med Clin N Am*. 2006;90:807–824.
- Chen YM, Hwang SJ, Chen LK, Chen DY, Lan CF. Risk factors for falls among elderly men in a veterans home. *J Chin Med Assoc*. 2008;71:180–185.
- Heath JM, Stuart MR. Prescribing exercise for frail elders. *J Am Board Fam Med*. 2002;15:218–228.
- Lu FP, Lin KP, Kuo HK. Diabetes and the risk of multi-system aging phenotypes: a systematic review and meta-analysis. *PLoS One*. 2009;4:e4144.
- Maurer MS, Burcham J, Cheng H. Diabetes mellitus is associated with an increased risk of falls in elderly residents of a long-term care facility. *J Gerontol A Biol Sci Med Sci*. 2005;60:1157–1162.
- Laubscher T, Regier L, Bareham J. Diabetes in the frail elderly: individualization of glycemic management. *Can Fam Physician*. 2012;58:543–546.
- Azidah A, Hasniza H, Zunaina E. Prevalence of falls and its associated factors among elderly diabetes in a tertiary center, Malaysia. *Curr Gerontol Geriatr Res*. 2012;2012:1–5.
- Gangavati A, Hajjar I, Quach L, et al. Hypertension, orthostatic hypotension, and the risk of falls in a community-dwelling elderly population: the Maintenance of Balance, Independent Living, Intellect, and Zest in the Elderly of Boston study. *J Am Geriatr Soc*. 2011;59:383–389.
- Carey BJ, Potter JF. Cardiovascular causes of falls. *Age Ageing*. 2001;30:19–24.
- Sharma V, Aggarwal S, Sharma A. Diabetes in elderly. *J Endocrinol Metab*. 2011;1:9–13.
- Duncan AK, Vittone J, Fleming KC, Smith HC. Cardiovascular disease in elderly patients. *Mayo Clin Proc*. 1996;71:184–196.
- Babatsikou F, Zavatsanou A. Epidemiology of hypertension in the elderly. *Health Sci J*. 2010;4:24–30.
- Chiu A, Au-Yeung S, Lo SK. A comparison of four functional tests in discriminating fallers from non-fallers in older people. *Disabil Rehabil*. 2003;25:45–50.
- Coleman P. Improving oral health care for the frail elderly: a review of widespread problems and best practices. *Geriatr Nurs*. 2002;23:189–199.
- Miegel K, Wachtel T. Improving the oral health of older people in long-term residential care: a review of the literature. *Int J Older People Nurs*. 2009;4:97–113.
- Yoshida M, Kikutani T, Okada G, Kawamura T, Kimura M, Akagawa Y. The effect of tooth loss on body balance control among community-dwelling elderly persons. *Int J Prosthodont*. 2009;22:136–139.
- Masud T, Morris RO. Epidemiology of falls. *Age Ageing*. 2001;30:3–7.
- Folstein MF, Folstein SE, McHugh PR. “Mini-mental state”. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12:189–198.
- Sanchez CE, Ogilvy CS, Carter BS. Outcomes studies in cerebrovascular neurosurgery. *Neurosurg Focus*. 2007;22:E11.
- Kose N, Cuvalci S, Ekici G, Otman AS, Karakaya MG. The risk factors of fall and their correlation with balance, depression, cognitive impairment and mobility skills in elderly nursing home residents. *Saudi Med J*. 2005;26:978–981.
- Podsiadlo D, Richardson S. The timed “Up and Go”: a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991;39:142–148.
- Yamamoto T, Kondo K, Hirai H, Nakade M, Aida J, Hirata Y. Association between self-reported dental health status and onset of dementia: a 4-year prospective cohort study of older Japanese adults from the Aichi Gerontological Evaluation Study (AGES) project. *Psychosom Med*. 2012;74:241–248.
- McCall WV. Sleep in the elderly: burden, diagnosis, and treatment. *Prim Care Companion J Clin Psychiatry*. 2004;6:9–20.
- Wu CY, Su TP, Fang CL, Chang MY. Sleep quality among community-dwelling elderly people and its demographic, mental, and physical correlates. *J Chin Med Assoc*. 2012;75:75–80.
- Schleicher MM, Wedam L, Wu G. Review of tai chi as an effective exercise on falls prevention in elderly. *Res Sports Med*. 2012;20:37–58.
- Tan MP, Kenny RA. Cardiovascular assessment of falls in older people. *Clin Interv Aging*. 2006;1:57–66.
- Cheng JW, Nayar M. A review of heart failure management in the elderly population. *Am J Geriatr Pharmacother*. 2009;7:233–249.
- Hernandez RK, Farwell W, Cantor MD, Lawler EV. Cholinesterase inhibitors and incidence of bradycardia in patients with dementia in the Veterans Affairs New England Healthcare System. *J Am Geriatr Soc*. 2009;57:1997–2003.
- Caska CM, Hendrickson BE, Wong MH, et al. Anger expression and sleep quality in patients with coronary heart disease: findings from the Heart and Soul Study. *Psychosom Med*. 2009;71:280–285.
- Wolk R, Gami AS, Garcia-Touchard A, Somers VK. Sleep and cardiovascular disease. *Curr Probl Cardiol*. 2005;30:625–662.

38. Heckman GA, Patterson CJ, Demers C, St Onge J, Turpie ID, McKelvie RS. Heart failure and cognitive impairment: challenges and opportunities. *Clin Interv Aging*. 2007;2:209–218.
39. Taylor J, Stott DJ. Chronic heart failure and cognitive impairment: co-existence of conditions or true association? *Eur J Heart Fail*. 2002;4:7–9.
40. Sowers JR, Epstein M, Frohlich ED. Diabetes, hypertension, and cardiovascular disease an update. *Hypertension*. 2001;37:1053–1059.
41. Langley F, Mackintosh SFH. Functional balance assessment of older community dwelling adults: a systematic review of the literature. *Internet J Allied Health Sci Pract*. 2007;5:1–11.
42. Gil-Montoya JA, de Mello AL, Cardenas CB, Lopez IG. Oral health protocol for the dependent institutionalized elderly. *Geriatr Nurs*. 2006;27:95–101.
43. Jablonski RA, Kolanowski A, Therrien B, Mahoney EK, Kassab C, Leslie DL. Reducing care-resistant behaviors during oral hygiene in persons with dementia. *BMC Oral Health*. 2011;11:30.
44. Fujihara I, Sadamori S, Abekura H, Akagawa Y. Relationship between behavioral and psychological symptoms of dementia and oral health status in the elderly with vascular dementia. *Gerodontology*. 2013;30:157–161.
45. Faulkner KA, Cauley JA, Zmuda JM, et al. Ethnic differences in the frequency and circumstances of falling in older community-dwelling women. *J Am Geriatr Soc*. 2005;53:1774–1779.
46. Means KM, O'Sullivan PS, Rodell DE. Balance, mobility, and falls among elderly African American women. *Am J Phys Med Rehabil*. 2000;79:30–39.
47. Hanlon JT, Landerman LR, Fillenbaum GG, Studenski S. Falls in African American and white community-dwelling elderly residents. *J Gerontol A Biol Sci Med Sci*. 2002;57:M473–M478.
48. Fuller GF. Falls in the elderly. *Am Fam Physician*. 2000;61:2159–2168.
49. Jean-Louis G, Magai C, Cohen C, et al. Ethnic differences in self-reported sleep problems in older adults. *Sleep*. 2001;24:926–936.
50. Anderson K. Falls in the elderly. *J R Coll Physicians Edinb*. 2008;38:138–143.
51. Riefkohl EZ, Bieber H, Burlingame M, Lowenthal D. Medications and falls in the elderly: a review of the evidence and practical considerations. *Pharm Ther*. 2003;28:724–733.
52. Stevens M, Holman CD, Bennett N, De Klerk N. Preventing falls in older people: outcome evaluation of a randomized controlled trial. *J Am Geriatr Soc*. 2002;49:1448–1455.
53. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing*. 2006;35:ii37–ii41.
54. Akyol A. Falls in the elderly: what can be done? *Int Nurs Rev*. 2007;54:191–196.
55. Bueno-Cavanillas A, Padilla-Ruiz F, Jiménez-Moleón JJ, Peinado-Alonso CA, Gálvez-Vargas R. Risk factors in falls among the elderly according to extrinsic and intrinsic precipitating causes. *Eur J Epidemiol*. 2000;16:849–859.
56. Al-Aama T. Falls in the elderly: spectrum and prevention. *Can Fam Physician*. 2011;57:771–776.
57. Ness KK, Gurney JG, Ice GH. Screening, education, and associated behavioral responses to reduce risk for falls among people over age 65 years attending a community health fair. *Phys Ther*. 2003;83:631–637.
58. Faulkner KA, Cauley JA, Zmuda JM, Griffin JM, Nevitt MC. Is social integration associated with the risk of falling in older community-dwelling women? *J Gerontol A Biol Sci Med Sci*. 2003;58:M954–M959.
59. Mukamal KJ, Mittleman MA, Longstreth WT Jr, Newman AB, Fried LP, Siscovick DS. Self-reported alcohol consumption and falls in older adults: cross-sectional and longitudinal analyses of the cardiovascular health study. *J Am Geriatr Soc*. 2004;52:1174–1179.
60. Teo JSH, Briffa NK, Devine A, Dhaliwal SS, Prince RL. Do sleep problems or urinary incontinence predict falls in elderly women? *Aust J Physiother*. 2006;52:19–24.
61. Roepke SK, Ancoli-Israel S. Sleep disorders in the elderly. *Indian J Med Res*. 2010;131:302–310.
62. Van Hoof J, Kort HS, Duijnste MS, Rutten PG, Hensen JL. The indoor environment and the integrated design of homes for older people with dementia. *Build Environ*. 2010;45:1244–1261.
63. Phair L, Heath H. Environments and older people with dementia. *Ment Health Pract*. 2001;4:32–38.

Medical Devices: Evidence and Research

Publish your work in this journal

Medical Devices: Evidence and Research is an international, peer-reviewed, open access journal that focuses on the evidence, technology, research, and expert opinion supporting the use and application of medical devices in the diagnosis, treatment and management of clinical conditions and physiological processes. The identification of novel

devices and optimal use of existing devices which will lead to improved clinical outcomes and more effective patient management and safety is a key feature. The manuscript management system is completely online and includes a quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from authors.

Submit your manuscript here: <http://www.dovepress.com/medical-devices-evidence-and-research-journal>

Dovepress