Epidemiology of stroke in a rural community in Southeastern Nigeria

Kelechi O Enwereji¹
Maduaburochukwu C
Nwosu¹
Adesola Ogunniyi²
Paul O Nwani¹
Azuoma L Asomugha¹
Ezinna E Enwereji³

¹Neurology Unit, Department of Medicine, Nnamdi Azikiwe University Teaching Hospital, Nnewi, Anambra State, Nigeria; ²Neurology Unit, Department of Medicine, University College Hospital Ibadan, Oyo State, Nigeria; ³Department of Community Medicine/Nursing Sciences, College of Medicine, Abia State University, Uturu, Abia State, Nigeria

Background: The prevalence and incidence of stroke vary from community to community worldwide. Nonetheless, not much is known about the current epidemiology of stroke in rural Nigeria and indeed Africa.

Methods: We carried out a two-phase door-to-door survey in a rural, predominantly low-income, community in Anambra, Southeastern Nigeria. We used a modified World Health Organization (WHO) protocol for detecting neurological diseases in the first phase, and a stroke-specific questionnaire and neurological examination in the second phase. An equal number of sex- and age-matched stroke-negative subjects were examined.

Results: We identified ten stroke subjects in the study. The crude prevalence of stroke in rural Nigeria was 1.63 (95% confidence interval [CI] 0.78–3.00) per 1,000 population. The crude prevalence of stroke in males was 1.99 (95% CI 0.73–4.33) per 1,000, while that for females was 1.28 (95% CI 0.35–3.28) per 1,000 population. The peak age-specific prevalence of stroke was 12.08 (95% CI 3.92–28.19) per 1,000, while after adjustment to WHO world population, the peak was 1.0 (95% CI 0.33–2.33) per 1,000.

Conclusion: The prevalence of stroke was found to be higher than previously documented in rural Nigeria, with a slightly higher prevalence in males than females. This is, however, comparable to data from rural Africa.

Keywords: Africa, developing country, prevalence

Introduction

Stroke has existed since medieval times and considerable progress has been made in the knowledge of its nature and epidemiology.^{1,2} It is the second leading cause of death worldwide.³ The prevalence and incidence of stroke vary from community to community and from time to time worldwide. However, not much is known about the burden of stroke in sub-Saharan Africa.

The crude prevalence of stroke reported in a rural study in Southwestern Nigeria was 0.58 per 1,000 population,⁴ while the incidence of stroke was found to be 26 per 100,000 in a Southwestern urban register-based study in Nigeria.⁵ These studies were conducted more than two decades ago. Another relatively recent prevalence study in a Southwestern Nigerian urban community was found to be 1.14 per 1,000 population.⁶ The current epidemiology of stroke in rural Nigeria is yet unknown.

Therefore, the purpose of this study was to determine the current epidemiology of stroke in a rural community in Southeastern Nigeria while maintaining a continuous register for the incidence of stroke.

Correspondence: Kelechi O Enwereji
Neurology Unit, Department of Medicine,
Nnamdi Azikiwe University Teaching
Hospital, Nnewi, Anambra State, Nigeria
Tel +234 803 386 1317
Email doublekay2001@yahoo.com

Methodology

This study was conducted in Ukpò, a predominantly low-income rural community with a population of 17,840 and 3,820 households (based on data from The National Population Commission of Nigeria on the 2006 national population census). Ukpò is the headquarters of Dunukofia Local Government Area in Anambra State, Southeastern region of Nigeria. It is a community with undulating terrain caused by erosion, which makes it not easily accessible by some vehicles. The community's source of clean water is a privately owned bore-hole. Electricity is sporadic, resulting in a few inhabitants using electricity-generating sets, while most use kerosene lanterns. The residents of the community are essentially monogamists, with few practicing polygamy. The majority are Christians, indigenous, and of the Igbo tribe. The community is stable and does not tolerate intertribal or interreligious marriages. The predominant occupation is peasant farming, and their diet is tuber-based. They are served by one fully functional primary health care center located at the center of the community. There is some labor migration out of the community. They usually return during festive periods and maintain households in the community. They do, however, return home when they become retired or are unable to work due to ill health.

Ethical clearance was obtained from the ethical committee of the Nnamdi Azikiwe University Teaching Hospital (NAUTH), Nnewi prior to commencement of the study. The study was a descriptive community-based door-to-door survey. It was conducted in two phases, preceded by a sensitization meeting with the community leaders (the Igwe-in-council) and a 2-day workshop with the research assistants. The village town criers, church leaders, and schools were engaged to make regular announcements weeks before the commencement of the survey. The research assistants were educated on the research procedures and the use of research instruments. This comprised a detailed review of the questionnaire, demonstration of interview technique, and back-demonstration by the research assistants. The 2-day workshop was fully attended by clinical medical students and other research assistants recruited for the first phase of the study. The medical students further participated in pre-testing the questionnaires on patients attending the medical outpatient clinic at the Nnamdi Azikiwe University Teaching Hospital, Nnewi on three consecutive clinic days and familiarized themselves with the research instrument.

Consent procedure

Informed oral consent was first obtained from the Honourable Commissioner for Health in the state, subsequently the

traditional ruler of the community and his council of chiefs, then the heads of households and individual subjects. Furthermore, informed consent was obtained from spouses or close relatives of those subjects who were incapable of giving their informed oral consent. However, an informed written consent was obtained from those who participated in the second phase of the study.

Phase I: census and door-to-door survey using modified World Health Organization (WHO) questionnaire

A census of the community was conducted in computer-generated random samples of 33% of the total enumeration areas 3 weeks prior to the survey to determine the average number of persons per enumeration area. The persons in each enumeration area were listed according to households using the name of the most prominent member. There was an average of 252 persons per enumeration area, estimating to a total Ukpò population of 18,648 using the 74 enumeration areas as obtained from the National Population Commission. Therefore 33% of the population, which was approximately 6,150 persons (25 enumeration areas), was selected for interview from the estimated Ukpò population using a table of computer-generated randomization. All the persons resident in the 25 randomly selected enumeration areas at the time of the study were included in both the census and the survey.

The first phase was conducted January 14–28, 2011. It was a door-to-door survey carried out by eight teams of trained research assistants, each comprising a health attendant, two medical students with at least 2 years of clinical experience, and a literate Ukpò resident as their liaison officer. The teams of research assistants administered the screening instrument on a total of 6,150 Ukpò residents. The screening instrument was the modified WHO questionnaire for detecting neurological diseases, and it was interviewer-administered to detect persons with probable stroke. It was adapted from the WHO protocol for detecting neurological diseases for the purpose of this study. The modifications included rearranging the order of the questions by bringing forth less probing questions before the more probing ones to improve participant cooperation, and exclusion of the physical examination components of the protocol to reduce administration time as the second phase of the study also had a physical examination component. Also, the questions for detecting tremors and Parkinsonism were removed for the above reasons. The questionnaire was translated to the local dialect and back-translated to English by two bilingual Igbo West African Examination Council examiners. The translated version was crosschecked by the neurologists

who are versed in the Igbo language before back-translation and adjustments were made to remove or improve ambiguous questions. These were fully explained to the research assistants. The research instrument (stroke-specific questionnaire) was validated in the Medical Out-Patient Department of NAUTH Nnewi using 20 patients and 20 students as controls. This yielded 20 as true negative, 8 as false positive, 12 as true positive, and 0 as false negative. Therefore, the sensitivity was 100% while the specificity was 71%.

The cases identified in this phase of the study as having probable stroke were enlisted for the second phase of the study. A positive response to any of questions 4, 7, and 8 in the screening instrument was considered as probable stroke and a formal invitation extended to the person to come to the Neuroepidemiology outpatient clinic in NAUTH Health Centre Ukpò for the second phase of the study.

Phase 2: screening with stroke-specific questionnaire and examination of participants

The second phase of the study was conducted in the outpatient clinic of the NAUTH Health Centre Ukpò by means of a semi-structured stroke-specific questionnaire and focused clinical examination (Supplementary material; Figures S1–S4). The radial pulse and all other peripheral pulses were checked, blood pressure measured, carotid bruits listened for, and the heart checked for evidence of likely embolic sources and hypertensive end-organ damage. A neurologic examination was done independently on each subject to detect residual paralysis, hemisensory loss, exaggerated deep tendon jerks, and an extensor plantar response by two consultant neurologists blinded to each other's results. Where there was disagreement, both neurologists reexamined the subject together and agreed on a common finding. An equal number of stroke-negative controls were randomly selected from those who screened negative for stroke to match subjects by age (± 2 years) and sex. These controls were screened and examined for risk factors, as was done for the stroke subjects. The diagnosis of stroke was clinical, based on the WHO definition.

Data analysis

Data were confidentially collected with study codes, recorded, cleaned, and analyzed using Epi Info™ (Centers for Disease Control and Prevention, Atlanta, GA, USA) 3.5.1 2008 statistical software. Continuous variables were presented as means with standard deviations, while categorical variables were presented as proportions. Crude, age-specific, and age-adjusted prevalences (WHO World Standard Population)⁷

were calculated alongside their confidence intervals (CIs). Student's *t*-test was used to compare means, while Fisher's exact test was used to compare categorical variables. *P*-value < 0.05 was considered statistically significant.

Results

Demography of study population

A total of 6,150 persons were screened in the first phase of the study. Table 1 shows age and sex distribution of the 6,150 persons, comprising 3,017 (49.06%) males and 3,133 (50.94%) females, with a male to female ratio of approximately 1:1. The ages of the respondents ranged from 1 month to 103 years (mean 29.4 ± 20.8 years). Those aged less than 45 years were 75.27% of the screened population, while those aged above 45 years comprised 24.73% of the population.

A total of 20 persons were recruited for the second phase (post-stroke-specific questionnaire) of the study. They were made up of ten subjects and ten controls that were matched for age and sex. The mean age of subjects was 60.7 ± 11.4 years (range 47–86 years), while the mean age of controls was 60.4 ± 11.9 years (range 45–84 years). There was no statistical significance between the age of the subjects and that of controls ($P = 0.853$).

Prevalence of stroke

Table 2 shows that the crude prevalence at the time of the survey on January 14, 2011 (point prevalence) of stroke was 1.63 (95% CI 0.78–3.00) per 1,000 population. The peak age-specific prevalence of stroke was found to be 1.0 (95% CI 0.33–2.33) when age-adjusted per 1,000 WHO world population. It also showed a higher crude prevalence of stroke in males of 1.99 (95% CI 0.73–4.33) per 1,000 than in females, which was 1.28 (95% CI 0.35–3.28) per 1,000 population (relative risk = 1.55; $P = 0.44$).

Table 1 Age and sex distribution of the 6,150 persons screened

Age group in years	Male		Female		Total	
	N	%	N	%	N	%
<5	407	13.49	350	11.17	757	12.31
5–14	813	26.95	686	21.89	1,499	24.37
15–24	537	17.80	572	18.26	1,109	18.03
25–34	326	10.81	411	13.12	737	11.98
35–44	218	7.22	310	9.90	528	8.58
45–54	227	7.52	267	8.52	494	8.03
55–64	199	6.59	215	6.86	414	6.74
65–74	160	5.31	169	5.40	329	5.35
≥ 75	130	4.31	153	4.88	283	4.61
Total	3,017	100	3,133	100	6,150	100

Table 2 Sex-specific, age-specific, and age-adjusted (WHO World Population) prevalence of stroke

Age group (in years)	Total population			Male		Female	
	Number of strokes/ population	Stroke prevalence (n/1,000)	Age-adjusted prevalence (n/1,000)	Number of strokes/ population	Stroke prevalence (n/1,000)	Number of strokes/ population	Stroke prevalence (n/1,000)
<45	0/4,630	0	0	0/2,301	0	0/2,329	0
45–54	2/494	4.05	0.46	1/227	4.41	1/267	3.75
55–64	5/414	12.08	1.00	2/199	10.05	3/215	13.95
65–74	2/329	6.08	0.31	2/160	12.50	0/169	0
≥75	1/283	3.53	0.11	1/130	7.69	0/153	0
Total	10/6,150	1.63	1.63	6/3,017	1.99	4/3,133	1.28

Abbreviation: WHO, World Health Organization.

Discussion

Previous studies in both developed and developing countries have reported crude prevalence of stroke ranging from 0 to 10.2 per 1,000 population, and the burden of stroke has also been projected to be on the rise.^{3,6} The crude prevalence of stroke in this study was comparatively higher than reported in previous community-based studies in Nigeria (1.63 versus 1.14, 0.58, and 0.68 per 1,000)^{4,6,8} but lower than reported in Aiyete, Southwestern Nigeria (4.43 per 1,000).⁹ Furthermore, we standardized our results to the WHO World Standard Population to compare better with data from other studies. In those aged above 7 years, our study's age-adjusted prevalence remained higher than those of Danesi et al⁶ and Osuntokun et al⁴ (1.70 versus 1.10 and 0.53 per 1,000 population, respectively). Apart from the Danesi et al study, which was conducted less than one decade ago, all the other Nigerian studies were done more than two decades ago. However, unlike ours and the earlier Nigerian studies which were conducted in predominantly low-income

rural communities, Danesi et al's study was conducted in a mixed-income, urban community. Also, the proportion of the population at particular risk for stroke (45 years and above) was higher in our study than that of Danesi et al (24.73% versus 15.24%).⁶ This may be partly explained by the common practice in Nigeria of urban to rural migration after retirement of about 67% of retirees.¹⁰

Furthermore, our results were compared with other populations as detailed in Table 3. In a rural South African survey, the age-adjusted prevalence among populations aged above 15 years was higher than ours (3.3 versus 1.90 per 1,000).¹¹ Also, an Asian study in urban Singapore reported a higher age-adjusted prevalence than ours in the population aged 65 years and above (6.31 versus 0.40 per 1,000).¹² Nonetheless, the study in urban Nigeria when compared with ours showed a lower age-adjusted prevalence in the population 15 years and above of 1.10 per 1,000 but a higher age-adjusted prevalence of 1.99 per 1,000 in the population 65 years and above.⁶ The small population size

Table 3 Characteristics of stroke prevalence studies compared with this study

Study	Period of data collection	Location	Rural/urban	Type of study	Population denominator	Mean age for stroke (years)	Sex M:F ratio	Age-adjusted prevalence (n/1,000)	Age range studied (years)
Osuntokun et al ⁹	1982	Aiyete	Rural	Community-based	903	??	??	4.43	All
Osuntokun et al ⁴	1982	Igbo-ora, Nigeria	Rural	Community-based	18,954	??	??	0.53	>7
Longe and Osuntokun ⁸	1987	Udo, Nigeria	Rural	Community-based	2,925	??	??	0.68	All
Ogun et al ¹⁴	1993–2003	Sagamu, Nigeria	Urban	Hospital-based	??	61.5	1.0:1.2	??	??
Connor et al ¹¹	2001	Agincourt, South Africa	Rural	Community-based	42,378	??	1.0:1.8	3.3	≥15
Danesi et al ⁶	2005–2006	Surulere, Nigeria	Urban	Community-based	13,127	63.4	5.5:2.0	1.14	All
Owolabi et al ¹³	2009	Ibadan, Nigeria	Urban	Hospital-based	??	59.4	2:3	??	??
This study	2011	Ukpo, Nigeria	Rural	Community-based	6,150	60.7	3:2	1.63	All

Abbreviations: F, female; M, male; ??, data not found.

of our study, as is common in many Sub-Saharan African studies, makes it difficult to give statistical interpretation to these differences with certainty.

The mean age for stroke obtained in this study is comparable to hospital-based data reported by Owolabi et al¹³ and Ogun et al¹⁴ in Southwestern Nigeria (60.7 versus 59.4 and 61.5 years, respectively). However, our result was lower than the mean age for stroke reported by Danesi et al⁶ of 63.4 years, although their peak age-adjusted prevalence for stroke was also higher than ours at 1.11 (75–84 years age group) versus 1.00 (55–64 years age group) per 1,000 population, respectively. The low mean age for stroke has been variously attributed to the lower life expectancy and high prevalence of fatal stroke in developing countries.^{14,15} The mean ages and peak prevalence for stroke in developed countries have been reported to be higher than those of developing countries.¹³ Also, the male preponderance of stroke noted in our study is comparable to data obtained in Sub-Saharan African studies.^{3,6,11,16}

Conclusion

The interpretative challenges experienced with variations in prevalence data from different communities notwithstanding, the prevalence of stroke is higher than previously reported in rural Nigeria. However, the study site is rural, with inhabitants predominantly of low socioeconomic status and a relatively higher proportion of the population being at risk of stroke. These are factors that may potentially affect the prevalence of stroke in the community.

We acknowledge the limitations of our study, as a larger population size would have been more representative; however, this is a modest attempt in our predominantly low income setting to assess the current prevalence of stroke in Nigeria. A longitudinal register is being kept to ascertain the current incidence of stroke in our study population. More studies to define the risk factors for stroke are recommended.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Ashrafian H. Familial stroke 2700 years ago. *Stroke* 2010;41(4):e187.
2. National Institute of Neurological Diseases and Stroke; Stroke: Hope Through Research. 2004 NIH Publication No 99-2222. Available at http://www.ninds.nih.gov/disorders/stroke/detail_stroke.htm. Accessed May 8, 2014.
3. Feigin VL, Lawes CM, Bennett DA, Anderson CS. Stroke epidemiology: a review of population-based studies of incidence, prevalence and case-fatality in the late 20th century. *Lancet Neurol*. 2003;2:43–53.
4. Osuntokun BO, Adeuja AO, Schoenberg BS, et al. Neurological disorders in Nigerian Africans: a community based study. *Acta Neurol Scand*. 1987;75:13–21.
5. Osuntokun BO, Bademosi O, Akinkugbe OO, Oyediran AB, Carlisle R. Incidence of stroke in an African city: results from the stroke registry at Ibadan, Nigeria, 1973–1975. *Stroke*. 1979;10:205–207.
6. Danesi M, Okubadejo N, Ojini F. Prevalence of stroke in an urban, mixed-income community in Lagos, Nigeria. *Neuroepidemiology*. 2007;28:216–223.
7. Ahmad OE, Boschi-Pinto C, Lopez AD, Murray CJL, Lozano R, Innoue M. *Age Standardization of Rates: a New WHO Standard*. GPE Discussion Paper Series No 31. Geneva: World Health Organization; 2001. Available from: <http://www.who.int/healthinfo/paper31.pdf>. Accessed April 7, 2014.
8. Longe AC, Osuntokun BO. Prevalence of neurological disorders in Udo, a rural community in Southern Nigeria. *Trop Geogr Med*. 1989;41:36–40.
9. Osuntokun BO, Schoenberg BS, Nottidge VA, et al. Research protocol for measuring the prevalence of neurologic disorders in developing countries. *Neuroepidemiology*. 1982;1(3):143–153.
10. Adebo GM, Sekumade AB. Socio-economic influence of retiree's migration on rural development in Ekiti State, Nigeria. *Int J Humanit Soc Sci*. 2012;2(12):164–172.
11. Connor MD, Thorogood M, Casserly B, Dobson C, Warlow CP; SASPI Project Team. Prevalence of stroke survivors in rural South Africa: results from the Southern Africa Stroke Prevention Initiative (SASPI) Agincourt field site. *Stroke*. 2004;35:627–632.
12. Venketasubramanian N, Tan LC, Sahadevan S, Chin JJ, et al. Prevalence of stroke among Chinese, Malay and Indian Singaporeans: a community-based tri-racial cross-sectional survey. *Stroke*. 2005;36:551–556.
13. Owolabi MO, Ugoya S, Platz T. Racial disparity in stroke risk factors: the Berlin-Ibadan experience: a retrospective study. *Acta Neurol Scand*. 2009;119:81–87.
14. Ogun SA, Ojini FI, Ogungbo B, Kolapo KO, Danesi M. stroke in South West Nigeria: a 10-year review. *Stroke*. 2005;36(6):112–122.
15. Onwuekwe IO, Ezeala-Adikaibe BA, Ohaegbulam SC, Chikani MC, Amuta J, Uloh HN. Stroke mimics – a study of CT images in Nigerian African stroke patients. *J Neurol Sci Turk*. 2008;25(3):148–154.
16. Amu E, Ogunrin O, Danesi M. Reappraisal of risk factors for stroke in Nigerian Africans – a prospective case-control study. *Afr J Neurol Sci*. 2005;2:20–27.

Supplementary material

Study No:..... Initials:.....

Sex: Male ☐ Female ☐

Age:

Residence: Were you resident in this area before August, 2010?

(a) No ☐ (b) Yes, all my life ☐ (c) Yes, not all my life ☐ (d) Don't know ☐ (e) No response

Occupation:

(a) Unknown ☐ (b) Preschool ☐ (c) School ☐ (d) Unemployed ☐ (e) Complete housewife
(f) Trader ☐ (g) Artisan ☐ (h) Teacher ☐ (i) Farmer ☐ (j) Civil servant ☐ (k) Retiree ☐
(l) Other specify

Religion:

(a) Christianity ☐ (b) Islam ☐ (c) African traditional religion ☐ (d) Atheism ☐ (e) No response ☐

Marital status:

(a) Married ☐ (b) Single ☐ (c) Divorced ☐ (d) Widow/widower ☐ (e) No response ☐

Place of origin:

Town: L.G.A: State:

Relationship to respondent:

(a) Self ☐ (b) Spouse ☐ (c) Parent ☐ (d) Offspring ☐ (e) Other specify:

Q1 Do you suffer from headache even when you don't have fever?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

Q2 Do you suffer from severe headaches, chiefly on one side of the head, which come on from time to time?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (c) No response ☐ (d) Not applicable ☐

Q3 In association with headache, do you suffer from nausea or vomiting?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐ (e) Not applicable ☐

Q4 Have you ever noticed that your face or mouth was shifted to one side (paralysed) for more than 24h?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

Q5 Have you ever had episodes where you lose contact with your surroundings for some minutes?

(a) Yes ☐ (b) Possible ☐ (c) Never ☐ (d) Don't know ☐ (e) No response ☐

Q6 Have you ever had any episode of shaking of your hands or legs for a brief period and which you could not control?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

Q7 Have you ever had loss of sensation or abnormal sensation affecting one side of your body (arm or leg) lasting for more than 24h?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

Q8 Have you ever had paralysis of one side of your body (arm or leg) lasting for more than 24h?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

Q9 Have you ever had convulsions that occurred when you had a fever?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

Q10 Have you ever had convulsions that occurred when you did not have a fever?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

Q11 Have you ever had any weakness or paralysis in both legs?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

To be screened by stroke questionnaire (for office use only)

(a) Yes ☐ (b) No ☐

Figure S1 Modified World Health Organization questionnaire for detecting neurological diseases.

SECTION A**Personal data:**

Study No:..... Initials:.....

Date of birth: (day)..... (month)..... (year).....

Sex: (a) Male ☐ (b) Female ☐

Marital status:

(a) Single ☐ (b) Married ☐ (c) Divorced ☐ (d) Widow/widower ☐ (e) No response ☐

Religion:

(a) Christianity ☐ (b) Islam ☐ (c) African traditional religion ☐ (d) Atheism ☐ (e) No response ☐

Education:

(a) Primary ☐ (b) Post-primary ☐ (c) Tertiary ☐ (d) No formal education ☐

Occupation:

(a) Unemployed ☐ (b) Student ☐ (c) Housewife ☐ (d) Artisan ☐ (e) Professional ☐ (f) Clergy ☐
(g) Farmer ☐ (h) Civil servant ☐ (i) Entertainer ☐ (j) Retiree ☐

Date of initial survey:.....2011

SECTION B

(1) Have you ever noticed that your face or mouth suddenly shifted to one side (paralysed) for more than 24h?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

(2) Have you ever had sudden loss of sensation or abnormal sensation affecting your arms and legs, lasting for more than 24h?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

(3) Have you ever had paralysis of one side of the body (arm or leg) lasting for more than 24h?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐ (d) No response ☐

(4) When did you first suffer from paralysis (weakness) or altered sensation affecting your arm or leg?

(day) (month) (year)

(5) How did the paralysis or altered sensation start?

(a) Suddenly ☐ (b) Gradually (over a period of 1 week or more) ☐

(6) If sudden, what were you doing when the paralysis occurred firstly?

(a) Resting ☐ (b) Intense activity (eg. arguing) ☐ (c) Others (specify)

(7) What other symptoms did you have when the paralysis started?

(a) None ☐ (b) Vomiting ☐ (c) Convulsion ☐ (d) Headache ☐ (e) Loss of consciousness ☐ (f) Loss of speech ☐ (g) Others specify:

(8) What time did the paralysis occur (approximate time)? AM: PM:.....

(9) Were you treated?

(a) Yes ☐ (b) No ☐ (c) Don't know ☐

- (10) If yes, where were you treated?
(a) Hospital ☐ (b) Home ☐ (c) Traditional healing home ☐ (d) Church ☐
(e) Others specify:
- (11) Were you admitted?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐
- (12) If yes, how long were you admitted?
(a) 1 week ☐ (b) Within 2 weeks ☐ (c) Within 1 month ☐ (d) >1 month ☐
- (13) What type of treatment did you receive?
(a) Oral drugs ☐ (b) IV drugs ☐ (c) Infusions ☐ (d) Herbs ☐ (e) Physiotherapy ☐ (f) Counselling ☐
(g) Prayers ☐ (h) Divination ☐ (i) Don't know ☐
- (14) When did you start receiving treatment?
(a) Same day ☐ (b) Within 1 week ☐ (c) After 1 week ☐ (d) No treatment ☐
- (15) How much did you recover by discharge?
(a) No recovery ☐ (b) Partial recovery ☐ (c) Full recovery ☐
- (16) Did you receive follow-up?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐
- (17) What side of your body was first affected by the paralysis?
(a) Right ☐ (b) Left ☐
- (18) What hand do you normally use to write or work?
(a) Right ☐ (b) Left ☐
- (19) Current neurological state (from observation or questioning)
(a) Paralysis still present ☐ (b) Paralysis recovered partially ☐ (c) Paralysis recovered fully ☐ (d) Speech difficulties still present ☐ (e) Speech difficulties recovered partially ☐ (f) Speech difficulty recovered fully ☐

SECTION C

- (20) Have you had this experience more than once?
(a) Yes ☐ (b) No ☐
- (21) If yes, how many times have you had it?
(a) Once ☐ (b) Twice ☐ (c) Thrice ☐ (d) More than thrice ☐
- (22) How long ago was the last experience?
(a) Within 1 month ☐ (b) Within 6 months ☐ (c) Within 1 year ☐ (d) More than 1 year ☐
- (23) How long after the first did you suffer the second stroke?
(a) Within 1 month ☐ (b) Within 6 months ☐ (c) Within 1 year ☐ (d) After 1 year ☐
- (24) What other symptoms did you have when the paralysis recurred?
(a) None ☐ (b) Vomiting ☐ (c) Convulsion ☐ (d) Headache ☐ (e) Loss of consciousness ☐ (f) Loss of speech ☐ (g) Others specify:

- (25) What time did the paralysis occur (approximate time)? AM:..... PM:.....
- (26) What were you doing when the paralysis recurred?
(a) Resting ☐ (b) Intense activity (eg. arguing) ☐ (c) Others specify:.....
- (27) Were you treated?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐
- (28) If yes, where were you treated?
(a) Hospital ☐ (b) Home ☐ (c) Traditional healing home ☐ (d) Church ☐ (e) Others specify:.....
- (29) Were you admitted?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐
- (30) If yes, how long were you admitted?
(a) <1 week ☐ (b) Within 2 weeks ☐ (c) Within 1 month ☐ (d) >1 month ☐
- (31) What type of treatment did you receive?
(a) Oral drugs ☐ (b) IV drugs ☐ (c) Infusions ☐ (d) Herbs ☐ (e) Physiotherapy ☐ (f) Counselling ☐
(g) Prayers ☐ (h) Divination ☐ (i) Don't know ☐
- (32) When did you start receiving treatment?
(a) Same day ☐ (b) Within 1 week ☐ (c) After 1 week ☐ (d) No treatment ☐
- (33) How much did you recover by discharge?
(a) No recovery ☐ (b) Partial recovery ☐ (c) Full recovery ☐
- (34) Did you receive follow-up?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐
- (35) What side of your body was affected by the repeat paralysis?
(a) Same side ☐ (b) Other side ☐
- (36) Current neurological state (from observation or questioning)
(a) Paralysis still present ☐ (b) Paralysis recovered partially ☐ (c) Paralysis recovered fully ☐ (d) Speech difficulties still present ☐ (e) Speech difficulties recovered partially ☐ (f) Speech difficulties recovered fully ☐
- (37) How long after the second did you suffer the third stroke?
(a) Within 1 month ☐ (b) Within 6 months ☐ (c) Within 1 year ☐ (d) After 1 year ☐
- (38) What other symptoms did you have when the paralysis recurred?
(a) None ☐ (b) Vomiting ☐ (c) Convulsion ☐ (d) Headache ☐ (e) Loss of consciousness ☐ (f) Loss of speech ☐ (g) Others specify:.....
- (39) What were you doing when the paralysis recurred?
(a) Resting ☐ (b) Intense activity (e.g. arguing) ☐ (c) Others specify:.....
- (40) What time did the paralysis occur? AM:..... PM:.....
- (41) Were you treated?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐

- (42) If yes where were you treated?
(a) Hospital ☐ (b) Home ☐ (c) Traditional healing home ☐ (d) Church ☐ (e) Others specify:.....
- (43) Were you admitted?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐
- (44) If yes, how long were you admitted?
(a) <1 week ☐ (b) Within 2 weeks ☐ (c) Within 1 month ☐ (d) >1 month ☐
- (45) What type of treatment did you receive?
(a) Oral drugs ☐ (b) IV drugs ☐ (c) Infusions ☐ (d) Herbs ☐ (e) Physiotherapy ☐ (f) Counselling ☐
(g) Prayers ☐ (h) Divination ☐ (i) Don't know ☐
- (46) When did you start receiving treatment?
(a) Same day ☐ (b) Within 1 week ☐ (c) After 1 week ☐ (d) No treatment ☐
- (47) How much did you recover by discharge?
(a) No recovery ☐ (b) Partial recovery ☐ (c) Full recovery ☐
- (48) Did you receive follow-up?
(a) Yes ☐ (b) No ☐ (c) Don't know ☐
- (49) What side of your body was affected by the recurred stroke?
(a) Same side ☐ (b) Other side ☐
- (50) Current neurological state (from observation or questioning)
(a) Paralysis still present ☐ (b) Paralysis recovered partially ☐ (c) Paralysis fully recovered ☐ (d) Speech difficulties still present ☐ (e) Speech difficulties recovered partially ☐ (f) Speech difficulties recovered fully ☐

Figure S2 Stroke-specific questionnaire used in Stage II of the study.

Personal data:

Study No:..... Initials:.....

Date of birth: (day) (month) (year)

Sex: (a) Male ☐ (b) Female ☐

Marital status:

(b) Single ☐ (b) Married ☐ (c) Divorced ☐ (d) Widow/widower ☐ (e) No response ☐

Religion:

(b) Christianity ☐ (b) Islam ☐ (c) African traditional religion ☐ (d) Atheism ☐ (e) No response ☐

Education:

(b) Primary ☐ (b) Post-primary ☐ (c) Tertiary ☐ (d) No formal education ☐

Occupation:

(b) Unemployed ☐ (b) Student ☐ (c) Housewife ☐ (d) Artisan ☐ (e) Professional ☐ (f) Clergy ☐
(g) Farmer ☐ (h) Civil servant ☐ (i) Entertainer ☐ (j) Retiree ☐

Date of initial survey:.....2011

Questionnaire

(1) Do you have any of the following conditions?

(a) Hypertension (high blood pressure) ☐ (b) Diabetes ☐ (c) Heart disease ☐
(d) Sickle cell disease ☐ (e) Epilepsy ☐ (f) Migraine ☐ (g) Previous stroke ☐
(h) Bleeding tendency ☐ (i) Others specify:

(2) Are you using any of these medications?

(a) Blood pressure medicine ☐ (b) Diabetes medicine ☐ (c) Others specify:.....

(3) Do you have a family member who has suffered from stroke in the past?

(a) Yes ☐ (b) No ☐If yes, specify who. (a) Father ☐ (b) Mother ☐ (c) Brother ☐ (d) Sister ☐ (e) Other relatives, specify relationship:.....

(4) Do you have a family member with hypertension (high blood pressure)?

(a) Yes ☐ (b) No ☐If yes, specify relationship. (a) Father ☐ (b) Mother ☐ (c) Brother ☐ (d) Sister ☐ (e) Other relative, specify relationship:.....

(5) Do you have a family member with diabetes?

(a) Yes ☐ (b) No ☐

(6) What is your level of education?

(a) No formal education ☐ (b) Primary ☐ (c) Secondary ☐ (d) Tertiary ☐

(7) What is your approximate monthly income?

(a) <^ 50,000 ☐ (b) ^ 50,000—^ 500,000 ☐ (c) >^ 500,000 ☐

- (8) Do you take alcohol (at present or in the past)?
- (a) Yes ☐ (b) No ☐
- If yes, specify:
- (i) Type of alcoholic drink: (a) Beer ☐ (b) Spirit ☐ (c) Stout ☐
- (ii) Number of bottles per week:.....
- (iii) Number of years:.....
- (9) Do you smoke (at present or in the past)?
- (a) Yes ☐ (b) No ☐
- If yes, specify:
- (i) Number of sticks per day:.....
- (ii) Duration of smoking (in years):.....
- (10) Do you use tobacco in any form (at present or in the past)?
- (a) Yes ☐ (b) No ☐
- If yes, specify: (a) Chewing ☐ (b) Snuffing ☐ (c) Cigar ☐

Figure S3 Risk factors for stroke questionnaire used in Stage II of the study.

Cardiovascular examination:

Pulse:

Blood pressure:

Apex:

Heart sounds:

Carotid bruit:

Nervous system examination:

Facial asymmetry:

Gait:

Power: right upper limb..... left upper limb.....

right lower limb..... left lower limb.....

Tone: right upper limb..... left upper limb.....

right lower limb..... left lower limb.....

Deep tendon reflex: biceps.....triceps.....

patellar.....achilles.....

Plantar response: right.....left.....

Laboratory investigation results:

Fasting blood sugar:

Total cholesterol:

LDL cholesterol:

Triglycerides:

HDL cholesterol:

Hb:

Total WBC:

Figure S4 Clinical examination findings (Proforma).**Abbreviations:** LDL, low density lipoprotein; HDL, high density lipoprotein; Hb, hemoglobin; WBC, white blood cells.**Vascular Health and Risk Management**

Dovepress

Publish your work in this journal

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement of

metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/vascular-health-and-risk-management-journal>