

# Cardiac arrest teams and time of day: effects on surviving in-hospital resuscitation

Martin Christ  
Wolfgang Dierschke  
Katharina Isabel von  
Auenmueller  
Marc van Bracht  
Martin Grett  
Hans-Joachim Trappe

Department of Cardiology and  
Angiology, Marienhospital Herne,  
Ruhr – University Bochum, Herne,  
Germany

**Objectives:** Little is known about the factors that influence survival following in-hospital resuscitation, but previous investigations have suggested that in-hospital resuscitations outside of regular working hours are associated with worse survival rates.

**Material and methods:** In-hospital cardiac arrest teams at our hospital were instructed to complete a questionnaire following every emergency call between July 2011 and June 2013. Data on all resuscitation attempts were collected and analyzed.

**Results:** A total of 65 in-hospital resuscitations were recorded in 42 males (64.6%) and 23 females (35.4%) (mean age 72.0±14.3 years). A total of 54 (83.1%) cardiac arrests were witnessed; seven (10.8%) showed a shockable rhythm at the time of the first ECG. Resuscitation attempts lasted 29.3±41.3 minutes, and 4.1±3.1 mg epinephrine was given. Return of spontaneous circulation could be achieved in 38 patients (58.5%); 29 (44.6%) survived the first day, 23 (35.4%) the seventh day, and 15 patients (23.1%) were discharged alive. Significantly more in-hospital resuscitations were obtained for those performed during non-regular working hours ( $P<0.001$ ), with higher neuron-specific enolase levels at 72 hours after resuscitation during nonregular working hours ( $P=0.04$ ). Patients who were discharged alive were significantly younger ( $P=0.01$ ), presented more often with an initial shockable rhythm ( $P=0.04$ ), and had a shorter duration of resuscitation ( $P<0.001$ ) with the need of a lower dose of epinephrine ( $P<0.001$ ).

**Discussion:** Survival rates following in-hospital resuscitation were poor at any time, but appear to depend less on time-dependent effects of the quality of resuscitation and more on time-dependent effects of recognition of cardiac arrests.

**Keywords:** sudden cardiac death, emergency medicine, time of day

## Objectives

Previous studies have estimated that a large number of in-hospital cardiac arrests occur; for example, more than 200,000 in-hospital cardiac arrests occur annually in the United States.<sup>1</sup> Despite this substantial number of patients, only a few studies have focused on this theme. Factors that have been described as being relevant for surviving an in-hospital cardiac arrest include an initial shockable rhythm, younger age, shorter duration of arrest, and time of arrest.<sup>2</sup> Time-dependent effects were also reported in another large investigation that described lower survival rates of in-hospital cardiac arrest during nights and weekends.<sup>3</sup>

Since the implementation of in-hospital cardiac arrest teams during the past few years, we are of the opinion that the time of an arrest can no longer automatically be associated with differences in the quality of resuscitation and the outcome following an in-hospital cardiac arrest. For this reason, we initiated this study to test our

Correspondence: Martin Christ  
Department of Cardiology and Angiology,  
Marienhospital Herne, Ruhr – University  
Bochum, Hoelkeskampring 40,  
44625 Herne, Germany  
Tel +49 2323 499 5620  
Fax +49 2323 499 301  
Email martin.christ@marienhospital-  
herne.de

hypothesis that, due to specialized emergency teams being available 24 hours a day, there are no time-dependent differences in survival of in-hospital cardiac arrest in a good neurological state.

## Material and methods

### In-hospital cardiac arrest team

In clinically unstable inpatients, early intervention by a medical emergency team significantly reduces the incidence of and mortality from unexpected in-hospital cardiac arrest.<sup>4</sup> Consequently, in our hospital, a central in-hospital cardiac arrest team was implemented a few years ago. This team comprises one emergency physician and an experienced intensive care nurse who are available 24 hours a day by emergency call and can reach every in-hospital emergency within 4 minutes.

### Data collection

In our hospital, all in-hospital emergency calls initially reach the coordination center, who then alert the emergency physician and the intensive care nurse. All emergency calls, therefore, can be registered in a central database, and our in-hospital cardiac arrest teams could be instructed to fill out a questionnaire following every emergency call due to a cardiac arrest between July 2011 and June 2013. The questionnaires contained data regarding sex, age, place of emergency, initial rhythm, and duration of resuscitation, as well as the number of defibrillations and applied medication.

Regular working hours were defined as the time between 8 am and 5 pm Monday through Friday, except for holidays. This threshold was chosen in accordance with the regular working hours of the physicians in our hospital.

The cardiac arrest was classified as “in-hospital” if it occurred in a hospitalized patient who had a pulse at the time of admission. Resuscitation attempts following in-hospital cardiac arrests were recorded and stored on a central database. Data were completed by additional review of the patient records. The primary endpoint of our study was survival following in-hospital cardiac arrest. The secondary endpoint was survival in a good neurological state (cerebral performance category [CPC] 1 or 2). Statistical analyses were performed using Student’s *t*-tests and Pearson’s chi-squared tests with the Statistical Package for the Social Sciences (SPSS 22.0; IBM, Armonk, NY, USA).

The study adhered to all criteria of the WMA Declaration of Helsinki – Ethical Principles for Medical Research Involving Human Subjects.

## Results

### Patient data

A total of 65 in-hospital resuscitations were recorded between July 1, 2011, and June 30, 2013, in 42 males (64.6%) and 23 females (35.4%) with a mean age of  $72.0 \pm 14.3$  years. A total of 54 (83.1%) in-hospital resuscitations followed witnessed arrests (either directly or indirectly via ECG monitoring); seven patients (10.8%) presented with an initial shockable rhythm.

Resuscitation attempts lasted  $29.3 \pm 41.3$  minutes, and  $4.1 \pm 3.1$  mg epinephrine was given. Initial return of spontaneous circulation (ROSC) could be achieved in 38 patients (58.5%); 29 (44.6%) survived the first day, 23 (35.4%) the seventh day, and 15 patients (23.1%) were discharged alive. Among those discharged, eleven patients (16.9%) showed a good cerebral performance category (CPC 1), two patients (3.1%) had moderate disabilities (CPC 2), and two patients (3.1%) remained in a vegetative state (CPC 4).<sup>5</sup>

The cause of death was cardiac-related in 36 patients (55.4%), hypoxemia in 16 patients (24.6%), and other reasons in 13 patients (20.0%). Neuron-specific enolase (NSE) level was  $57.5 \pm 108.8$  ng/mL at 24 hours after resuscitation, and  $35.7 \pm 19.3$  ng/mL at 72 hours after resuscitation.

The first electrocardiogram (ECG) after resuscitation showed heart rates of  $90.2 \pm 27.6$  beats per minute (bpm) and a frequency-related QT time of  $462.1 \pm 42.5$  ms; maximal creatinine kinase was  $2197.9 \pm 1960.0$  U/L.

After ROSC, the following were observed: first pH =  $7.25 \pm 0.15$ , pCO<sub>2</sub> =  $48.2 \pm 26.6$  mmHg, pO<sub>2</sub> =  $130.3 \pm 104.8$  mmHg, lactate =  $5.6 \pm 4.8$  mmol/L, and potassium =  $4.2 \pm 0.8$  mmol/L (Table 1).

### Resuscitation during regular and nonregular working hours

According to the classification of regular and nonregular working hours as described in the Data collection section, significantly more in-hospital resuscitations were observed during nonregular working hours ( $P < 0.001$ ). No differences were observed between patients resuscitated during regular working hours and those resuscitated during nonregular working hours, except for a significantly higher NSE level at 72 hours after resuscitation during nonregular working hours ( $P = 0.04$ ; Table 1).

### Post hoc analyses depending on patient follow-up

A comparison of patients who were discharged alive following in-hospital resuscitation and those who died during follow-up showed several significant differences (Table 2). Patients who

**Table 1** Patient characteristics

	All patients (n=65)	Regular working hours (n=25)	Nonregular working hours (n=40)	P-value
Male, n (%)	42 (64.6)	14 (56.0)	28 (70.0)	0.25
Age (years)	72.0±14.3	69.5±13.6	73.6±14.6	0.26
Witnessed event, n (%)	54 (83.1)	21 (84.0)	33 (82.5)	0.88
Shockable rhythm, n (%)	7 (10.8)	4 (16.0)	3 (7.5)	0.28
Duration CPR (minutes)	29.3±41.3	35.7±62.1	25.2±19.6	0.42
Epinephrine (mg)	4.1±3.1	4.41±3.57	3.88±2.88	0.57
Return of spontaneous circulation, n (%)	38 (58.5)	17 (68.0)	21 (52.5)	0.22
Expected cause, n (%)				
Cardiac	36 (55.4)	17 (68.0)	19 (47.5)	0.36
Hypoxemia	16 (24.6)	5 (20.0)	11 (27.5)	
Others	13 (20.0)	3 (12.0)	10 (25.0)	
Mild therapeutic hypothermia (33°C), n (%)	3 (4.6)	0 (0.0)	3 (7.5)	0.28
Survival at first day, n (%)	29 (44.6)	12 (48.0)	17 (42.5)	0.66
Survival at seventh day, n (%)	23 (35.4)	11 (44.0)	12 (30.0)	0.25
Discharged alive, n (%)	15 (23.1)	7 (28.0)	8 (20.0)	0.46
NSE after 24 hours (ng/mL)	57.5±108.8	31.3±24.5	70.7±132.5	0.38
NSE after 72 hours (ng/mL)	35.7±19.3	18.4±3.9	42.6±18.5	<b>0.04</b>
Cerebral Performance Category, n (%)				
1	11 (16.9)	6 (24.0)	5 (12.5)	0.36
2	2 (3.1)	1 (4.0)	1 (2.5)	
3	0 (0.0)	0 (0.0)	0 (0.0)	
4	2 (3.1)	0 (0.0)	2 (5.0)	
Heart rate (beats/minute)	90.2±27.6	85.9±26.7	94.3±28.8	0.42
QTc duration (ms)	462.1±42.5	453.0±49.4	470.6±34.4	0.28
Creatinine kinase maximum (U/L)	2,197.9±1,960.0	1,276.5±1,418.4	3,119.3±2,098.1	0.11
First pH	7.25±0.15	7.26±0.19	7.25±0.10	0.86
pCO <sub>2</sub> (mmHg)	48.2±26.6	48.3±33.9	48.1±20.3	0.98
pO <sub>2</sub> (mmHg)	130.3±104.8	145.3±131.5	118.3±80.5	0.54
Lactate (mmol/L)	5.6±4.8	6.6±5.4	4.7±4.2	0.33
Potassium (mmol/L)	4.2±0.8	4.2±0.8	4.3±0.8	0.79

**Notes:** Continuous variables are expressed as mean ± standard deviation. Bold P-values were considered to be significant.

**Abbreviations:** CPR, cardiopulmonary resuscitation; NSE, neuron-specific enolase; QTc, frequency-related QT time.

were discharged alive were significantly younger ( $P=0.01$ ), presented more often with an initial shockable rhythm ( $P=0.04$ ), and had a shorter duration of resuscitation ( $P<0.001$ ) with the need for a lower dose of epinephrine ( $P<0.001$ ).

Patient characteristics, including first blood gas analysis and ECG criteria following ROSC, showed no significant differences.

## Discussion

Duration of cardiopulmonary resuscitation, age, initial rhythm, and witnessed events are factors that have been described to influence survival of any kind of sudden cardiac arrest; ie, in-hospital cardiac arrest<sup>6</sup> as well as out-of-hospital cardiac arrest.<sup>7</sup> However, as patients of in-hospital cardiac arrest are hospitalized due to preexisting illness, they cannot be directly compared with patients of out-of-hospital cardiac arrests; for example, acute apoplectic stroke, preexisting severe sepsis, and preexisting poor CPC scores have been

associated with a very poor prognosis for surviving in-hospital cardiac arrest in a good neurological state.<sup>8</sup>

For this reason, further studies were demanded to specifically focus on in-hospital cardiac arrests, and we initiated our study to investigate whether the time of arrest still influences survival from in-hospital cardiac arrest despite the implementation of in-hospital cardiac arrest teams.

Unfortunately, despite the implementation of in-hospital cardiac arrest teams, survival rates following in-hospital resuscitation are poor even with acceptable rates of initial ROSC.<sup>2,9–11</sup> In our study, only 23.1% of all resuscitated patients were discharged alive, and only 20.0% of all resuscitated patients left the hospital in a good neurological state (CPC 1 or 2).

## Effect of time on survival rates following in-hospital cardiac arrest

It has been assumed that the time of an arrest may influence survival rates following in-hospital cardiac arrest, with

**Table 2** Post hoc analyses depending on patient's survival during follow-up


	Discharged alive (n=15)	Died during follow-up (n=50)	P-value
Male, n (%)	9 (60.0)	33 (66.0)	0.76
Age (years)	62.9±13.6	74.7±13.4	<b>0.01</b>
Witnessed event, n (%)	14 (93.3)	40 (80.0)	0.43
Shockable rhythm, n (%)	4 (26.7)	3 (6.0)	<b>0.04</b>
Duration CPR (minutes)	6.7±7.4	36.0±44.9	<b>&lt;0.001</b>
Epinephrine (mg)	1.6±1.2	4.5±3.2	<b>&lt;0.001</b>
Return of spontaneous circulation, n (%)	15 (100.0)	23 (46.0)	<b>&lt;0.001</b>
Expected cause, n (%)			
Cardiac	9 (60.0)	27 (54.0)	0.27
Hypoxemia	5 (33.3)	11 (22.0)	
Others	1 (6.7)	12 (24.0)	
Mild therapeutic hypothermia (33°C), n (%)	2 (13.3)	1 (2.0)	0.13
Survival at first day, n (%)	15 (100.0)	14 (28.0)	<b>&lt;0.001</b>
Survival at seventh day, n (%)	15 (100.0)	8 (16.0)	<b>&lt;0.001</b>
Discharged alive, n (%)	15 (100.0)	0 (0.0)	<b>&lt;0.001</b>
NSE after 24 hours (ng/mL)	92.3±157.2	27.1±14.8	0.32
NSE after 72 hours (ng/mL)	41.9±21.3	27.3±15.8	0.35
Heart rate (bpm)	88.8±28.5	91.1±27.9	0.84
QTc duration (ms)	445.2±41.1	472.4±41.0	0.10
Creatinine kinase maximum (U/L)	2,485.8±1,963.5	1,992.3±2,086.7	0.69
First pH	7.30±0.09	7.23±0.17	0.21
pCO <sub>2</sub> (mmHg)	43.9±14.3	50.3±31.2	0.47
pO <sub>2</sub> (mmHg)	134.1±62.4	128.3±122.3	0.87
Lactate (mmol/L)	4.4±3.4	6.1±5.3	0.31
Potassium (mmol/L)	3.97±0.51	4.32±0.86	0.06

**Notes:** Continuous variables are expressed as mean ± standard deviation. Bold P-values were considered to be significant.

**Abbreviations:** CPR, cardiopulmonary resuscitation; NSE, neuron-specific enolase; QTc, frequency-related QT time.

poorer results obtained for resuscitation performed during nonregular working hours.<sup>2,3,10</sup> Nevertheless, previous studies did not consider that 24/7 in-hospital resuscitation teams might cause comparable conditions for victims of in-hospital cardiac arrests independent from the time of an arrest.

In our study, time-dependent effects were minimal; the only observed difference was a higher NSE level at 72 hours after in-hospital resuscitation during nonregular working hours, which affected neither the survival rates nor the neurological outcomes among survivors (Table 1 and Figure 1). We therefore doubt that the time of resuscitation has a relevant effect on the efficiency of the resuscitation procedure itself; at least in hospitals with 24/7 in-hospital cardiac arrest teams. Following this idea, we think that



**Figure 1** Patient survival following in-hospital cardiac arrest, depending on the time of resuscitation (regular versus nonregular working hours).

**Abbreviation:** ROSC, return of spontaneous circulation.

lower survival rates from in-hospital cardiac arrest during nights and weekends may be more a problem of a later detection of the cardiac arrest. This theory is supported by the following two findings. First, the nurse-to-bed ratio has been described as a relevant modifiable factor for survival following an in-hospital cardiac arrest.<sup>12</sup> Second, the median hospital cardiac arrest incidence rate has been described as 4.02 per 1,000 admissions,<sup>12</sup> but during our study period, approximately 35,000 admissions were observed in our hospital, which would imply that approximately 140 expected in-hospital cardiac arrests were observed during the same time. Nevertheless, only 65 resuscitation attempts were identified in our study, suggesting a significant percentage of in-hospital cardiac arrests that occurred without resuscitation attempts. Furthermore, 83.1% of cardiac arrests were witnessed in our study, which indicates that resuscitation attempts were mainly initiated following a witnessed event. Considering these findings, we think those previously described effects of time of arrest on survival rates might be more a problem of delayed recognition rather than poor resuscitation.

## Limitations

The main limitation of this study is the small number of patients. We therefore think that further studies should be performed to verify our results.

## Conclusion

Survival rates following in-hospital resuscitation might depend less on time-dependent effects of the quality of resuscitation and more on time-dependent effects of recognition. Nevertheless, our single-center results should be confirmed by further investigations and larger trials.

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Merchant RM, Yang L, Becker LB, et al; American Heart Association Get with the Guidelines-Resuscitation Investigators. Incidence of treated cardiac arrest in hospitalized patients in the United States. *Crit Care Med.* 2011;39(11):2401–2406.
2. Cooper S, Janghorbani M, Cooper G. A decade of in-hospital resuscitation: outcomes and prediction of survival? *Resuscitation.* 2006;68(2):231–237.
3. Peberdy MA, Ornato JP, Larkin GL, et al; National Registry of Cardiopulmonary Resuscitation Investigators. Survival from in-hospital cardiac arrest during nights and weekends. *JAMA.* 2008;299(7):785–792.
4. Buist MD, Moore GE, Bernard SA, Waxman BP, Anderson JN, Nguyen TV. Effects of a medical emergency team on reduction of incidence of and mortality from unexpected cardiac arrests in hospital: preliminary study. *BMJ.* 2002;324(7334):387–390.
5. Ajam K, Gold LS, Beck SS, Damon S, Phelps R, Rea TD. Reliability of the Cerebral Performance Category to classify neurological status among survivors of ventricular fibrillation arrest: a cohort study. *Scand J Trauma Resusc Emerg Med.* 2011;19:38.
6. Kantamineni P, Emani V, Saini A, Rai H, Duggal A. Cardiopulmonary Resuscitation in the Hospitalized Patient: Impact of System-Based Variables on Outcomes in Cardiac Arrest. *Am J Med Sci.* Epub 2014 Apr 23.
7. McNally B, Robb R, Mehta M, et al; Centers for Disease Control and Prevention. Out-of-hospital cardiac arrest surveillance – Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005–December 31, 2010. *MMWR Surveill Summ.* 2011; 60(8):1–19.
8. Ebell MH, Afonso AM, Geocadin RG; American Heart Association's Get With the Guidelines-Resuscitation (formerly National Registry of Cardiopulmonary Resuscitation) Investigators. Prediction of survival to discharge following cardiopulmonary resuscitation using classification and regression trees. *Crit Care Med.* 2013;41(12): 2688–2697.
9. Bloom HL, Shukrullah I, Cuellar JR, Lloyd MS, Dudley SC, Zafari AM. Long-term survival after successful in-hospital cardiac arrest resuscitation. *Am Heart J.* 2007;153(5):831–836.
10. Ružman T, Tot OK, Ivić D, Gulam D, Ružman N, Burazin J. In-hospital cardiac arrest: can we change something? *Wien Klin Wochenschr.* 2013; 125(17–18):516–523.
11. Kaernsted B, Indridason OS, Baldursson J, Arnar DO. [In-hospital cardiopulmonary resuscitation at Landspítali University Hospital in Reykjavik]. *Laeknabladid.* 2009;95(7–8):509–514. Icelandic.
12. Chen LM, Nallamothu BK, Spertus JA, Li Y, Chan PS; American Heart Association's Get With the Guidelines-Resuscitation (formerly the National Registry of Cardiopulmonary Resuscitation) Investigators. Association between a hospital's rate of cardiac arrest incidence and cardiac arrest survival. *JAMA Intern Med.* 2013;173(13): 1186–1195.

International Journal of General Medicine

**Publish your work in this journal**

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-general-medicine-journal>

Dovepress

A key focus is the elucidation of disease processes and management protocols resulting in improved outcomes for the patient. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.