Nutritional management of enterocutaneous fistula: a retrospective study at a Malaysian university medical center

Manal MH Badrasawi¹

Suzana Shahar¹

Ismail Sagap²

¹Dietetics Program, School of Health Care Sciences, Faculty of Health Sciences, ²Department of Surgery, Faculty of Medicine, UKM Medical Center, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia

Abstract: Enterocutaneous fistula is a challenging clinical condition with serious complications and considerable morbidity and mortality. Early nutritional support has been found to decrease these complications and to improve the clinical outcome. Location of the fistula and physiological status affect the nutrition management plan in terms of feeding route, calories, and protein requirements. This study investigated the nutritional management procedures at the Universiti Kebangsaan Malaysia Medical Center, and attempted to determine factors that affect the clinical outcome. Nutritional management was evaluated retrospectively in 22 patients with enterocutaneous fistula seen over a 5-year period. Medical records were reviewed to obtain data on nutritional status, biochemical indices, and route and tolerance of feeding. Calories and protein requirements are reported and categorized. The results show that surgery was the predominant etiology and low output fistula was the major physiological category; anatomically, the majority were ileocutaneous. The spontaneous healing rate was 14%, the total healing rate was 45%, and the mortality rate was 22%, with 14% due to fistula-associated complications. There was a significant relationship between body mass index/serum albumin levels and fistula healing; these parameters also had a significant relationship with mortality. Glutamine was used in 50% of cases; however, there was no significant relationship with fistula healing or mortality rate. The nutritional status of the patient has an important impact on the clinical outcome. Conservative management that includes nutrition support is very important in order to improve nutritional status before surgical repair of the fistula.

Keywords: enterocutaneous fistula, nutritional management, retrospective study

Background

Management of enterocutaneous fistula is a challenge for surgeons, gastroenterologists, and allied health professionals¹⁻⁴ because of the serious complications that can occur, ie, malnutrition, sepsis, and fluid and electrolyte disturbances,⁵ known as the “fistula triad”.⁶ The main goal in management of enterocutaneous fistula is to restore the integrity of the patient’s gastrointestinal tract to allow normal feeding while minimizing morbidity and mortality.³ The management plan is usually divided into three phases. The first phase, ie, recognition and stabilization, includes resuscitation of the patient, control of sepsis, electrolyte repletion, control of fistula drainage, and supply of nutrition. In the second phase, a fistulogram is performed to identify the anatomy and characteristics of the fistula. In the third phase, a decision is taken regarding the duration of conservative management and the operative plan based on an evaluation of the patient and the fistula.⁷

Nutritional support involves assessment, diagnosis, ordering, preparation, distribution, administration, and monitoring of nutrition. It includes total parenteral

Correspondence: Suzana Shahar
Dietetics Programme Centre of Health
Care Sciences, Faculty of Health Sciences,
Universiti Kebangsaan Malaysia,
Jalan Raja Muda Abd Aziz, Kampung Baru,
Kuala Lumpur 53100, Malaysia
Email suzana.shahar@gmail.com

nutrition (TPN), enteral feeding, or oral feeding, and plays a major role in management of enterocutaneous fistula.^{8–10} It begins with calculating calories and protein requirements, assessing tolerance of the feeding regimen, modifying feeding methods, adjusting nutritional requirements according to changes in clinical status, and finally, observing for feeding complications.¹¹ Classification of the fistula affects the route of feeding and caloric and protein requirements;¹² ie, patients with high output fistula should receive more calories and protein than those with low output fistula who mostly require TPN. Oral and enteral feeding increase the fistula output if the fistula occurs in the proximal part of the gut.² Therefore, a multidisciplinary team should be involved in designing and performing the management plan.¹³ Despite attention to nutritional support, malnutrition remains a major clinical problem in 50%–90% of patients with enterocutaneous fistula.² Earlier researchers have noted that there are no experimental data for optimal routes of feeding in patients with enterocutaneous fistula,¹⁴ and any relevant reports are based on observational studies.

This study reports on the nutritional management of patients with enterocutaneous fistula admitted to the National University of Malaysia Medical Center and factors affecting the outcome of treatment.

Materials and methods

This retrospective observational study identified 28 cases admitted to the National University of Malaysia Medical Center between 2007 and 2012 based on manual screening of the census books held in the general surgery ward. Twenty-six files available at the medical records unit were reviewed to extract the relevant data. Four cases were excluded because of incomplete information regarding nutritional management or lack of basic anthropometric measurements.

The data extracted included a detailed medical history, clinical status, and anatomic, etiologic, and physiologic classification of fistula. Physiologic classification was done by calculating the mean fistula output for the first 3 days, starting from the day the fistula was first recognized, using the following output cutoff points: low <200 mL/day; moderate 200–500 mL/day; and high >500 mL/day.² Each patient's nutritional status on admission and discharge was assessed using the available data for weight, body mass index, subjective global assessment, and serum albumin level.

The nutritional management plan and monitoring were investigated for each case individually, and categorized according to the route of feeding, duration of feeding, caloric and protein requirements, and type of formula, ie,

enteral or parenteral. Refeeding syndrome was deemed to have occurred if a team member recommended cessation or a decrease in feeding in the presence of hypokalemia, hyponatremia, and hypophosphatemia.¹⁵ The medical notes for nutritional management recorded by all health professionals, ie, surgeons, nurses, dietitians, clinical pharmacists, psychologists, and physiotherapists, were reviewed for all cases.

Enterocutaneous fistula-related complications, including sepsis, electrolyte disturbances, dehydration, and malnutrition documented in the medical notes were also recorded. Patients who had multiple admissions were considered as single cases. Nutritional management was planned individually, depending on the patient's general status, because guidelines for nutritional support in patients with enterocutaneous fistula were not available.

Fisher's exact test was used to determine relationships between categorical variables, ie, nutritional status classification, and mortality, spontaneous healing, or occurrence of complications. The relationship between nutrition parameters and fistula healing was determined using Spearman's correlation coefficient test. Further analysis using multiple linear regression was applied to determine the predictive variables. Cox regression analysis was performed to identify factors affecting mortality. Descriptive statistics were used for type of fistula, mortality, fistula healing, and clinical complication rates.

Results

Only 22 of the 28 cases with enterocutaneous fistula identified met our inclusion criteria. These included 16 males and six females of mean age 50 ± 17 (range 16–72) years. The majority of cases were Malay ($n=17$), with a minority being Chinese ($n=5$). Comorbidities included hypertension ($n=8$), stage 4 chronic kidney disease ($n=2$), diabetes ($n=2$), hepatic disease ($n=1$), and cerebrovascular accident ($n=1$). A history or presence of malignancy was reported in ten cases. Single or multiple laparotomy had been performed before formation of the fistula in all but two cases. The mean duration of hospital stay was 27.4 ± 16 (range 8–65) days.

Nutritional status

Mean body mass index was 20.1 ± 3.6 (14.0–32.0) kg/m². Five cases were underweight, eleven were normal weight, two were overweight, and four were obese. Subjective global assessment was reported for nine cases, seven of whom were severely malnourished and the other two were mildly to moderately malnourished. The mean albumin

level on admission was 27.9 ± 8.7 (range 15–44) mmol/L and at discharge (or the last reading before discharge) was 29.2 ± 8.6 (13–43) mmol/L.

Fistula classification and sequelae

The main etiology of fistula was surgical (in 20 cases), and the remaining cases were spontaneous, with one case due to inflammatory bowel disease and the other being an iatrogenic fistula. Most of the cases involved single fistula, with only one case of multiple fistulae. Low output was the major physiologic classification (15 cases), followed by high output fistula in six cases and medium output in one case. With regard to anatomic classification, the majority (15 cases) were iliac fistulae, while six cases involved the colon and two cases involved the jejunum. There was only one case of duodenal fistula (see Table 1).

The mortality rate was 23% (five cases). Three deaths were due to complications, eg, sepsis. Spontaneous healing of the fistula was achieved in only three cases by conservative treatment alone, while seven cases required conservative

management followed by surgical repair. Two cases were repaired surgically immediately after occurrence of the fistula. The overall healing rate with surgical repair and/or conservative management was 45%.

Sepsis occurred in eleven cases, electrolyte disturbances occurred in eight, and dehydration in three. Three cases developed a combination of sepsis, electrolyte disturbance, and dehydration. Four cases developed sepsis and electrolyte disturbance. Elevated liver enzymes were reported in two cases during TPN, and five cases were admitted to the intensive care unit. Conservative management was the first choice for management in 76% of cases.

Nutritional management regimen

The medical records showed that the following parameters affected the nutritional management plan: fistula output, functional status of the gastrointestinal tract, location of fistula (proximal or distal), feeding tolerance, presence of complications such as sepsis, TPN (liver)-related complications, and need for patient to be nil per mouth before surgery.

Table 1 Summary of the cases according to selected variables

Number	Age, years	Sex	Anatomic position	Etiology	Physiology	Start feeding*	Days of TPN	Days of EF	Oral	Healing	LOS	Mortality
1	46	Male	Jejunum	Postoperative	LOP	5	–	14	–	Not	19	–
2	71	Female	Jejunum	Postoperative	LOP	6	12	–	3	Not	45	–
3	67	Female	Ileum	Postoperative	LOP	5	3	14	–	Not	32	–
4	57	Female	Ileum	Postoperative	LOP	4	3	–	9	Healed	15	–
5	16	Male	Terminal ileum	IBD	HOP	7	25	–	5	Healed	38	–
6	72	Female	Duodenum	Iatrogenic	MOP	6	–	9	–	Not	23	+
7	72	Male	Ileum	Postoperative	HOP	7	8	20	–	Not	37	+
8	61	Male	Ileum	Postoperative	LOP	1	–	14	–	Not	17	–
9	32	Male	Ileum	Postoperative	HOP	6	9	–	5	Not	48	+
10	41	Male	Colon	Postoperative	LOP	4	–	14	–	Healed	11	–
11	27	Male	Colon	Postoperative	LOP	8	5	–	–	Healed	17	–
12	40	Male	Ascending colon	Postoperative	LOP	7	–	–	4	Not	8	–
13	52	Male	Ileum	Postoperative	LOP	6	–	9	–	Healed	8	–
14	41	Male	Ileum	Postoperative	LOP	6	–	20	4	Healed	9	–
15	41	Male	Ileum	Postoperative	LOP	7	14	14	11	Not	30	–
16	53	Male	Ileum	Postoperative	LOP	3	23	–	4	Healed	45	–
17	18	Male	Ileum	Postoperative	LOP	5	29	–	32	Healed	64	+
18	57	Female	Descending colon	Postoperative	LOP	4	–	–	14	Not	17	–
19	67	Female	Ileum	Postoperative	LOP	3	12	–	4	Not	18	–
20	57	Female	Descending colon	Postoperative	LOP	4	–	–	14	Not	17	–
21	53	Male	Ileum	Postoperative	LOP	3	23	–	4	Healed	45	–
22	41	Male	Colon	Postoperative	LOP	4	–	14	–	Healed	11	–

Notes: *From the day of admission; + mortality occurred and was reported; – mortality did not occur/was not reported.

Abbreviations: LOP, low output; MOP, medium output; HOP, high output; EF, enteral feeding; TPN, total parenteral nutrition; Oral, days of oral feeding using nutritional formula; IBD, inflammatory bowel disease; LOS, length of stay.

The nutritional management regimens were as follows: TPN with enteral feeding followed by oral nutrition or sip feeding; TPN followed by oral feeding; enteral feeding followed by oral feeding; TPN only; enteral feeding only; and oral feeding only. The duration and route of nutritional management is summarized in Table 1.

Glutamine was given to nine cases, either by intravenous infusion, or by adding it to the TPN bag. Indications for oral glutamine included occurrence of or risk of sepsis (ie, uncontrolled effluent from the fistula).⁷ However, no significant relationship was found between fistula healing and mortality with intravenous glutamine.

Ready-mixed TPN bags were given to 14 (50%) of the patients, while the rest were given compound TPN bags. Standard formulae were used in most patients, but a modular formula (myotein) was added to the standard formula in one case (4.5%). Three cases (14%) were temporally given elemental formula after converting from TPN to enteral feeding. Patient caloric and protein requirements were calculated using a rapid method. Calories were given as 20–35 kcal/kg/day and protein was given as 1.2–2 g/kg/day.

The relationship between anatomic fistula classification and spontaneous healing was analyzed by Fisher's exact test, and the associations were not significant at P -value <0.05 . The relationship between days of nil per mouth and the final albumin result was not statistically significant ($P=0.77$, Pearson's correlation coefficient test; Table 2). No significant predictor was found for mortality using Cox regression analysis (Table 3).

The relationship between fistula healing and mortality as dependent variables and etiology of the fistula, physiological status, and patient nutritional status showed different levels of significance (Table 4). The relationship between nutritional management regimen with serum albumin level at discharge was not statistically significant, but the relationship was significant with the length of hospital stay ($P<0.05$, Kruskal–Wallis test).

Table 2 Correlation between fistula healing and nutritional parameters

	Nutritional parameters				
	BMI	Albumin (on admission)	Total protein (on admission)	Days of NPO	Days of TPN
Fistula healing	0.344	0.330	0.343	−0.150	0.342
P -value*	0.069	0.081	0.070	0.451	0.153

Note: *Using Spearman's correlation test.

Abbreviations: BMI, body mass index; NPO, nil per mouth; TPN, total parenteral nutrition.

Table 3 Relationship between fistula healing, mortality, and nutritional parameters

Nutrition parameters	P-value	Exp (B)	95% CI	
			Lower bound	Higher bound
Fistula healing ^a				
BMI	0.008	1.328	0.959	1.839
Albumin (upon admission)	0.121	1.096	0.976	1.23
Days of NPO	0.484	0.809	0.447	1.46
Days of TPN	0.117	1.127	0.97	1.30
Use of glutamine	0.645	1.07	0.19	5.91
Mortality ^b				
BMI	0.343	0.847	0.60	1.194
Albumin (upon admission)	0.412	0.919	0.752	1.12
Days of NPO	0.264	2.934	0.443	10.42
Days of TPN	0.606	0.822	0.822	1.12
Use of glutamine	0.200	5.197	0.418	64.5

Notes: ^aLogistic regression test; ^bCox regression test.

Abbreviations: BMI, body mass index; CI, confidence interval; NPO, nil per mouth; TPN, total parenteral nutrition; Exp (B), exponential B (odds ratio).

Discussion

The results of this study show that the most common etiology of fistula was surgery and sepsis was the main reason for the considerable mortality rate. Fistula healing was significantly related to certain aspects of nutritional status. There was no significant difference in nutritional status according to the nutrition support regimen used, but there was a significant difference according to period of hospitalization. The spontaneous healing rate after conservative management was 14%, and the total healing rate was 45%. Spontaneous healing was associated with low output, a surgical etiology, and absence of sepsis. Several factors have already been reported to affect the rate of spontaneous healing, including surgical etiology and low output, with no associated abscess or comorbidity.¹⁴ However, a few studies have shown a higher fistula healing rate.^{16–18} The highest healing rate of 81% was reported by

Table 4 Relationship between fistula healing and mortality versus fistula classification, patient factors, and management plan

Independent variables	Fistula healing	Mortality
Etiologic classification	0.361 ^a	0.151
Anatomic	0.125 ^a	0.406
Physiologic	0.67 ^a	0.017*
Patient BMI category	0.05 ^a	0.017*
Nutrition support regimen	0.149 ^a	0.48
Glutamine	0.429 ^a	0.303
Days nil per mouth	0.123 ^b	0.503
Serum albumin at admission	0.004 ^{a,b}	0.274
BMI at admission	0.003 ^{a,b}	0.001*
Age	0.827 ^b	0.375

Notes: ^aUsing Fisher's exact test; ^busing independent t -test; *significant at $P<0.05$.

Abbreviation: BMI, body mass index.

Gupta et al,¹⁶ 67% was reported by Harriman et al,¹⁷ and 53.3% was reported by Nursal Tarik et al.¹⁸ This difference in reported rates is likely related to the experience and skill of the surgeon or the patient's condition.¹⁴

The mortality rate due to fistula complications was 14%, and included sepsis, electrolyte disturbance, dehydration, and malnutrition. This is consistent with a report by Martinez et al,¹⁹ who found that these complications had a significant relationship with mortality. Gupta et al¹⁶ also reported a rate of 14%, whereas a rate of 24.4% was reported by Nursal Tarik et al.¹⁸

Malnutrition was significantly related to mortality, as evidenced by the significant association of body mass index with mortality, even though it was not significantly associated with serum albumin level. Mawdsley et al²⁰ found a significant difference in serum albumin levels between surviving and nonsurviving groups ($P=0.01$). A large sample size could explain this significant relationship; however, their study did not investigate body mass index and its relationship with mortality.

We found no significant relationship between patient age and healing or mortality rates. This finding is not consistent with those of other studies;²⁰ in clinical practice, mortality rate is generally affected by age and nutritional status.²¹

Nutritional management and clinical outcome

The cases were managed conservatively, as recommended by the published studies on management of fistula,^{4,7,10,22} with the exception of one case that was treated surgically. Conservative treatment was used to build up nutritional status, stabilize the patient before surgery, improve the surgical outcome, and decrease the risk of postoperative complications.¹¹ A subjective global assessment was available for nine cases; seven were malnourished, indicating that this assessment was performed only for such cases. Due to the limited data on nutritional status in the patient files, assessment was based on body mass index and serum albumin level at admission or fistula recognition and the last albumin level before discharge. The significant relationship between serum albumin and spontaneous healing was in accordance with other similar studies. Nursal Tarik et al¹⁸ found a significant relationship between serum albumin and fistula healing ($P=0.014$). Patients with healed fistula had higher serum albumin levels than those with non-healed fistulae. Harriman et al found a significant relationship between fistula healing and serum albumin levels, and fistula healing and prealbumin levels ($P=0.018$ and $P=0.00$, respectively).¹⁷

Caloric and protein requirements have been shown to be affected by patient status, nutritional status, feeding tolerance, fistula output, and presence of complications such as refeeding syndrome and sepsis.²³ Individualizing the nutritional management plan in terms of caloric and protein requirements and feeding route has been recommended in several studies.^{6,11,24}

Relationship between glutamine, fistula healing, and mortality

Intravenous glutamine was administered to 41% of the patients. At our study center, glutamine is commonly given to patients with or at high risk of sepsis, including those with enterocutaneous fistula. However, administration of glutamine did not significantly improve the clinical outcome. Similar studies assessing the effects of glutamine or an immune-enhancing diet on the clinical outcome in patients with fistula were not found.

Conclusion

Nutritional status has a significant impact on both healing and clinical outcome of fistula. Building up nutritional status in patients with fistula is very important in conservative management and in preparing patients for surgical repair.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Draus JM, Huss SA, Harty NJ, Cheadle WG, Larson GM. Enterocutaneous fistula: are treatments improving? *Surgery*. 2006; 140(4):570–578.
2. Dudrick S, Panait L. Metabolic consequences of patients with gastrointestinal fistulas. *Eur J Trauma Emerg Surg*. 2011;37(3):215–225.
3. Evenson AR, Fischer JE. Current management of enterocutaneous fistula. *J Gastrointest Surg*. 2006;10(3):455–464.
4. Taggarshe D, Bakston D, Jacobs M, McKendrick A, Mittal VK. Management of enterocutaneous fistulae: a 10 years experience. *World J Gastrointest Surg*. 2010;2(7):242–246.
5. Pfeifer J, Tomasch G, Uranues S. The surgical anatomy and etiology of gastrointestinal fistulas. *Eur J Trauma Emerg Surg*. 2011;37(3): 209–213.
6. Schecter WP, Hirshberg A, Chang DS, et al. Enteric fistulas: principles of management. *J Am Coll Surg*. 2009;209(4):484–491.
7. Polk TM, Schwab CW. Metabolic and nutritional support of the enterocutaneous fistula patient: a three-phase approach. *World J Surg*. 2012;36(3):524–533.
8. Visschers RG, Damink SWO, Winkens B, Soeters PB, van Gemert WG. Treatment strategies in 135 consecutive patients with enterocutaneous fistulas. *World J Surg*. 2008;32(3):445–453.
9. Tong WM, Hope W, Overby DW, Hultman CS. Comparison of outcome after mesh-only repair, laparoscopic component separation, and open component separation. *Ann Plast Surg*. 2011;66(5):551–556.
10. Whelan J, Ivatury R. Enterocutaneous fistulas: an overview. *Eur J Trauma Emerg Surg*. 2011;37(3):251–258.

11. Yanar F, Yanar H. Nutritional support in patients with gastrointestinal fistula. *Eur J Trauma Emerg Surg*. 2011;37(3):227–231.
12. Shenkin A. Serum prealbumin: is it a marker of nutritional status or of risk of malnutrition? *Clin Chem*. 2006;52(12):2177–2179.
13. McNaughton V. Summary of best practice recommendations for management of enterocutaneous fistulae from the Canadian Association for Enterostomal Therapy ECF Best Practice Recommendations Panel. *J Wound Ostomy Continence Nurs*. 2010;37(2):173–184.
14. Lloyd D, Gabe S, Windsor A. Nutrition and management of enterocutaneous fistula. *Br J Surg*. 2006;93(9):1045–1055.
15. Khan L, Ahmed J, Khan S, MacFie J. Refeeding syndrome: a literature review. *Gastroenterol Res Pract*. 2010;2011:410971.
16. Gupta M, Sonar P, Kakodkar R. Small bowel enterocutaneous fistulae: the merits of early surgery. *Indian J Surg*. 2008;70(6):303–307.
17. Harriman S, Rodych N, Hayes P, Moser MAJ. The C-reactive protein-to-prealbumin ratio predicts fistula closure. *Am J Surg*. 2011;202(2):175–178.
18. Nursal Tarık Z, Ataç B, Hasan A, Ömer A. Factors affecting healing of enterocutaneous fistulas. *Turk J Gastroenterol*. 2000;11(3):222–226.
19. Martinez JL, Luque-de-León E, Ballinas-Oseguera G, Mendez JD, Juárez-Oropeza MA, Román-Ramos R. Factors predictive of recurrence and mortality after surgical repair of enterocutaneous fistula. *J Gastrointest Surg*. 2012;16(1):156–163.
20. Mawdsley J, Hollington P, Bassett P, Windsor A, Forbes A, Gabe S. An analysis of predictive factors for healing and mortality in patients with enterocutaneous fistulas. *Aliment Pharmacol Ther*. 2008;28(9):1111–1121.
21. Dempsey D, Mullen J, Buzby G. The link between nutritional status and clinical outcome: can nutritional intervention modify it? *Am J Clin Nutr*. 1988;47(2):352–356.
22. Schecter WP. Management of enterocutaneous fistulas. *Surg Clin North Am*. 2011;91(3):481–491.
23. Tong CY, Lim LL, Brody RA. High output enterocutaneous fistula: a literature review and a case study. *Asia Pac J Clin Nutr*. 2012;21(3):464–469.
24. Yuan Y, Ren J, Gu G, Chen J, Li J. Early enteral nutrition improves outcomes of open abdomen in gastrointestinal fistula patients complicated with severe sepsis. *Nutr Clin Pract*. 2011;26(6):688–694.

Journal of Multidisciplinary Healthcare

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or

Submit your manuscript here: <http://www.dovepress.com/journal-of-multidisciplinary-healthcare-journal>

healthcare processes in general. The journal covers a wide range of areas and welcomes submission from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress