ER- α 36: a novel biomarker and potential therapeutic target in breast cancer

Xingyun Su¹Xin Xu¹Guangliang Li^{1,4}Bingyi Lin²Jiang Cao³Lisong Teng¹

¹Department of Surgical Oncology, First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, Zhejiang, People's Republic of China; ²Key Laboratory of Combined Multi-Organ Transplantation, Ministry of Public Health, Hangzhou, Zhejiang, People's Republic of China; ³Clinical Research Center, The 2nd Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, Zhejiang, People's Republic of China;

⁴Department of Medical Oncology, Zhejiang Cancer Hospital, Hangzhou, Zhejiang Province, People's Republic of China

Correspondence: Lisong Teng
Department of Surgical Oncology,
First Affiliated Hospital, College of
Medicine, Zhejiang University, 79
Qingchun Road, Hangzhou 310003,
Zhejiang, People's Republic of China
Tel +86 571 8706 8873
Fax +86 571 8723 6628
Email lsteng@zju.edu.cn

Jiang Cao
Clinical Research Center, The
2nd Affiliated Hospital, College
of Medicine, Zhejiang University,
Hangzhou, Zhejiang Province,
People's Republic of China
Tel +86 571 8731 5201
Fax +86 571 8731 5201
Email caoj@zju.edu.cn

Abstract: Estrogen receptor-alpha36 (ER- α 36) is a 36-kDa variant of estrogen receptor-alpha (ER- α) firstly identified and cloned by Wang et al in 2005. It lacks both transactivation domains (activation function 1 and activation function 2) and has different biological characteristics compared to traditional ER- α (ER- α 66). ER- α 36 primarily locates on plasma membrane and cytoplasm and functions as a mediator in the rapid membrane-initiated non-genomic signaling pathway. Additionally, it inhibits the traditional genomic signaling pathway mediated by ER- α 66 in a dominant-negative pattern. Accumulating evidence has demonstrated that ER- α 36 regulates the physiological function of various tissues. Thus, dysregulation of ER- α 36 is closely associated with plenty of diseases including cancers. ER- α 36 is recognized as a molecular abnormality which solidly correlates to carcinogenesis, aggressiveness, and therapeutic response of breast cancer. Additionally, special attention has been paid to the role of ER- α 36 in endocrine therapy resistance. Therefore, ER- α 36 provides a novel biomarker of great value for diagnosis, prognosis, and treatment of breast cancer. It may also be a potential therapeutic target for breast cancer patients, especially for those who are resistant to endocrine therapy. In this review, we will overview and update the biological characteristics, underlying mechanism, and function of ER- α 36, focusing on its biological function in breast cancer and endocrine therapy resistance. We will evaluate its application value in clinical practice.

Keywords: ER, ER- α 36, breast cancer, endocrine therapy resistance

Introduction

Estrogen is essential to the development and physiology of target organs, such as uterus, ovary, and breast. Dysregulation of estrogen results in pathological processes including osteoporosis, breast cancer, and endometrial cancer.¹ These pleiotropic effects of estrogen are related to estrogen receptors (ERs) which are generally regarded as ligand-activated transcription factors belonging to the nuclear receptor superfamily.¹⁻³ Two different forms of ER have been discovered, ER- α and ER- β .^{3,4} They share similar architecture composed of three independent but interactional function domains.^{3,5} The N-terminal A/B domains possess a ligand-independent transactivation domain (activation function 1 [AF-1]), interacting with co-activators and activating transcription of target genes.⁶ The C or DNA-binding domain contains two zinc finger-like structures, allowing dimerization of receptors and then binding to target DNA sequences.⁶ The C-terminal D/E/F domains or ligand-binding domains mediate ligand binding, receptor dimerization, and nuclear localization and have a ligand-dependent activation domain (activation function 2 [AF-2]) (Figure 1B).⁶

ER- α 36 is a 36-kDa variant of ER- α generating from full length ER- α by alternative splicing or different usage of promoters.⁶⁻⁹ Compared to ER- α 66, ER- α 36 retains

Figure 1 Structure of ER-α66 and ER-α36.

Notes: (A) Transcription of ER-α36 initiates from a previously unidentified promoter in the first intron of ER-α66. We labeled the first exon of ER-α36 as “1” to distinguish it. Besides, ER-α36 has an extra, unique 27-amino-acid sequence at C-terminus which may broaden ligand-binding spectrum of ER-α36. (B) Compared to the protein structure of ER-α66, ER-α36 lacks both transactivation domains of AF-1 and AF-2 and retains DBD, Hinge, and LBD of ER-α66. Therefore, ER-α36 functions as a powerful competitor of ER-α66.

Abbreviations: AF-1, activation function 1; AF-2, activation function 2; DBD, DNA binding domain; ER-α36, estrogen receptor-alpha36; ER-α66, estrogen receptor-alpha66; LBD, ligand binding domain.

DNA-binding, partial dimerization, and ligand-binding domains but lacks both transactivation domains (AF-1 and AF-2).^{3,14,15} ER-α36 mainly locates on plasma membrane and cytoplasm mediating non-genomic estrogen signaling.^{6,10,11} Additionally, it inhibits traditional genomic estrogen signaling mediated by ER-α66 in a dominant-negative pattern.¹⁰ Accumulating evidence has demonstrated that ER-α36 participates in the development and physiological function of many tissues. Therefore, it is related to diseases such as postmenopausal osteoporosis,¹² airway hyperresponsiveness,¹³ and even malignancies including gastric,^{14–16} hepatic,¹⁷ endometrial,¹⁸ and breast cancers.¹⁹ ER-α36 is crucial in the carcinogenesis and aggressiveness of breast cancer; in addition, it explains why antiestrogen therapy loses efficacy in ER-positive breast cancer patients.^{20–22} These achievements imply that ER-α36 is a promising biomarker for diagnosis, prognosis, and treatment of various diseases, especially breast cancer.^{23–25} What's more, it provides a novel pharmaceutical approach to individualized treatment of breast cancer.

This review will overview and update the biological characteristics, molecular mechanism, and biological role of ER-α36, focusing on its function in carcinogenesis, progression, and endocrine therapy resistance of breast cancer and its potential application value in clinical practice.

Basic characteristics of ER-α36

Transcription of ER-α36 initiates from a previously unidentified promoter in the first intron of ER-α66, which continues

from exon 2 to exon 6 and skips the final exon 7 and exon 8 of ER-α66 (Figure 1A).^{3,6,10,14,15} As a result, ER-α36 lacks both transactivation domains (AF-1 and AF-2) and retains DNA-binding, partial dimerization, and ligand-binding domains of ER-α66 (Figure 1B). The last 138 amino acids encoded by the final exon 7 and exon 8 are replaced by an extra, unique 27-amino-acid sequence at C-terminus. This extra sequence may broaden the ligand-binding spectrum of ER-α36 so that it is able to interact with more factors other than estrogen.¹⁰ ER-α36 primarily locates on plasma membrane (50%) and cytoplasm (40%) rather than nucleus.^{10,19,26–28} Because it has three potential myristoylation sites near N-terminus instead of the nuclear-location signals of ER-α66, ER-α36 may be modified by palmitoylation posttranslationally and then locate on plasma membrane and cytoplasm just like ER-α46, another variant of ER-α66.^{6,10,11} Therefore, ER-α36 is a potential regulator of membrane-initiated mitogenic signaling pathways responding to E2α, E2β, E3, E4, and even tamoxifen, an estrogen antagonist.^{6,10,24}

Though ER-α36 exists in both ER-α66-positive and ER-α66-negative breast cancer cells,^{10,28} it is highly expressed in the majority of ER-α66-negative breast cancer.^{19,29} Furthermore, overexpression of ER-α36 is usually accompanied with a decrease of ER-α66,^{10,24,28,30} indicating that ER-α66 and ER-α36 are mutually regulated and inversely associated.^{10,20,24,27–30} Wilms' tumor suppressor WT1 is one of the coordinators which activates promoters of ER-α66 but suppresses ER-α36 promoter activity.³¹ Another coordinator

is synuclein γ (SNCG), which binds to ER- α 66 or ER- α 36 depending on E2 concentration.^{32,33} In addition, ER- α 66 can negatively regulate the promoter activity of ER- α 36.³⁰ Deng et al found that proteasome inhibitors dramatically increased the steady level of ER- α 66, indicating that the proteasome system took part in the expression regulation of ER- α 36 and ER- α 66.³⁴ With the dynamic regulation of ER- α 66 and ER- α 36, genomic and non-genomic estrogen signaling pathways are also coordinated to maintain a balance.³¹ Although the explicit mechanism needs further investigation, it is obvious that an imbalance of ER- α 66 and ER- α 36 may result in abnormal proliferation and differentiation, finally leading to diseases including breast cancer.¹

Molecular mechanism underlying ER- α 36 action

Under normal conditions, ERs bind to estrogen and then transfer to the nucleus, subsequently binding to specific estrogen response elements and regulating transcription of downstream DNA (Figure 2).³⁵ Other than the traditional genomic pathway, ERs mediate the non-genomic pathway by membrane-associated receptors.^{36–39} The non-genomic pathway controls more genes than the genomic pathway, regulating cellular growth, survival, motility, invasion, and apoptosis. What's more, non-physiologic estrogen also activates the non-genomic pathway, which is important to

so-called “nontarget” tissue such as the cardiovascular system.^{40,41} The unique structure and intracellular location of ER- α 36 decide its molecular mechanism. Lacking both transcriptional activation domains, ER- α 36 binds to the same target DNA sequence as ER- α 66 does, but possesses no intrinsic transcriptional activity. Thus, ER- α 36 functions as a powerful competitor of the genomic signaling pathway mediated by ER- α 66.^{6,10} Additionally, ER- α 36 mediates rapid membrane-initiated estrogen signaling cascades, which regulate various biological functions and are associated with tumorigenesis, aggressive phenotype, and treatment sensitivity of carcinomas (Figure 2).^{16,23,24,42–44}

ER- α 36 and downstream kinases

The mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/ERK) pathway is the most popular mitogenic signaling in cancers, which can be rapidly activated by E2.^{45,46} Research has demonstrated that membrane-initiated signaling pathways mediated by ER- α 36 could stimulate the transcription of ELK1 in the nucleus.^{10,26,42,47,48} Tong et al found that ER- α 36 also activated ERK1/2 through the protein kinase C delta signaling pathway and then elevated expression of cyclin D1/cyclin-dependent kinase 4, a transcriptional coregulator that regulates cell cycle progression.⁴⁸ The MAPK/ERK signaling pathway mediated by ER- α 36 contributes to the potential invasion and metastasis of cancer

Figure 2 Genomic and non-genomic activities mediated by ERs.

Notes: ER- α 66 binds to estrogen and then transfers to the nucleus to interact with specific EREs and induce transcription of target genes. ER- α 36 has a broader ligand-binding spectrum than ER- α 66 so that it can respond to E2 α , E2 β , E3, E4, and even tamoxifen. ER- α 36 can mediate rapid MIES such as MAPK/ERK, PI3K/Akt, and PKC δ to regulate biological functions of cells.

Abbreviations: Akt, protein kinase B; BCSCs, breast cancer stem cells; cdk 4, cyclin-dependent kinase 4; ER- α 36, estrogen receptor- α 36; ER- α 66, estrogen receptor- α 66; EREs, estrogen response elements; ERK, extracellular signal-regulated kinase; ERs, estrogen receptors; GSK3 β , glycogen synthase kinase 3 beta; JNKs, c-Jun N-terminal kinases; MAPK, mitogen-activated protein kinases; MIES, membrane-initiated estrogen signaling; PKC δ , protein kinase C delta; PI3K, phosphatidylinositol 3-kinase.

cells.^{23,24,44} What's more, it induces paclitaxel resistance through c-Jun N-terminal kinases (JNKs), an important component of the MAPK family mediating paclitaxel-induced apoptosis and death (Figure 2).^{23,24,44}

Phosphatidylinositol 3-kinase/protein kinase B (PI3K/Akt) is another classical pathway associated with malignant cancers.^{49–51} Rapid phosphorylation of Akt, which is induced by E2, testosterone, and tamoxifen, can be blocked by a PI3K inhibitor or knockdown of ER- α 36, suggesting that the PI3K/Akt pathway is one of the non-genomic pathways mediated by ER- α 36 (Figure 2).^{26,47}

ER- α 36 associated proteins

Several proteins are involved in the stability and function of ER- α 36, thus promoting mitogenic estrogen signaling mediated by ER- α 36. Glucose-regulated protein 94, an endoplasmic reticulum chaperone, is a potential downstream effector of the ER- α 36-mediated pathway in gastric cancer.¹⁵ SNCG, previously regarded as breast cancer-specific gene *BCSG1*,⁵² binds to ER- α 36 without heat shock protein 90 to prevent degradation of ER- α 36 and significantly strengthen ER- α 36-mediated mitogenic estrogen signaling pathways.³³

Many investigators discovered a significant co-expression of ER- α 36 and epidermal growth factor receptor (EGFR) in primary breast cancers, indicating that ER- α 36 took part in EGFR-related carcinogenesis.¹⁹ Further studies elucidated

that epidermal growth factor (EGF) induced phosphorylation of ERK1/2 via ER- α 36 in a time- and dose-dependent pattern.¹⁸ A positive feedback loop was confirmed that EGFR signaling activated transcription of ER- α 36 through an activator-protein-1-binding site in the promoter of ER- α 36. In turn, ER- α 36 interacted with the EGFR/Src/Shc complex to strengthen the EGFR signaling pathway and stabilize EGFR protein (Figure 3).²⁰ A similar feedback loop between ER- α 36 and human epidermal growth factor receptor 2 (HER-2), a member of the EGFR family, was reported.⁵³ Interestingly, our previous study found that HER-2 expression didn't increase in tamoxifen-resistant cells, which overexpressed ER- α 36 and EGFR.²³ In the Src/EGFR/signal transducer and activator of transcription 5 (STAT5) pathway mediated by ER- α 36, Src functions as a switch to adjust phosphorylation of EGFR and then recruits STAT5 as a downstream effector, which regulates activation of the MAPK/ERK signaling pathway and expression of cyclin D1 (Figure 3).^{16,20,54} Therefore, the positive feedback loops between ER- α 36 and growth factor receptors are important for cellular function, although the definite mechanism has not been clarified.

Biological roles of ER- α 36

ER- α 36 participates in the development and function of target tissues. For example, expression and subcellular localization of ER- α 36 are necessary for folliculogenesis,

Figure 3 Positive feedback loop between EGFR and ER- α 36.

Notes: A positive feedback loop has been confirmed that EGFR signaling activates transcription of ER- α 36 through an AP-1-binding site in the promoter region of ER- α 36. In turn, ER- α 36 is able to stabilize EGFR protein and mediate MAPK/ERK and Src/EGFR/STAT5 pathways. In the Src/EGFR/STAT5 pathway, Src functions as a switch by phosphorylation to enhance proliferation and malignant properties of cancer.

Abbreviations: AP-1, activator protein 1; EGFR, epidermal growth factor receptor; ER- α 36, estrogen receptor- α 36; ERK, extracellular signal-regulated kinase; MAPK, mitogen-activated protein kinases; STAT5, signal transducer and activator of transcription 5.

oocyte meiotic maturation, ovulation, and luteinization of the postnatal ovary, thus facilitating maturation of the female reproductive system.⁵⁵ Additionally, ER- α 36 maintains the bone density of postmenopausal women.¹² However, dysregulation of ER- α 36 leads to various dysfunctions and diseases, such as osteoporosis,¹² airway hyperresponsiveness,¹³ and even cancers.^{14,17–19} Since the mammary gland is estrogen-dependent, ER- α 36 plays a particularly vital role in breast cancer. Therefore, we will focus on its role in breast cancer and endocrine therapy resistance and its potential application value in clinical practice.

Biological function of ER- α 36 in breast cancer

In order to investigate the biological function of ER- α 36 in breast cancer, Zhang et al established ER- α 36-knockdown MDA-MB-231 cell lines in which expression of ER- α 36 decreased obviously. They found knockdown of ER- α 36 suppressed proliferation, migration, and invasion and increased sensitivity to paclitaxel in MDA-MB-231 cells.²⁴ Chaudhri et al got similar results that ER- α 36 was able to enhance cellular proliferation and metastasis and protect against apoptosis.⁴³ Recently, Yu et al studied the relationship between ER- α 36 and chemotherapy response in 120 breast cancer patients and found that ER- α 36-negative patients had better response to anthracycline and/or paclitaxel than ER- α 36-positive patients.²⁵

ER- α 36 and breast cancer stem cells (BCSCs)

BCSCs, a subpopulation of stem/progenitor cells that possesses the ability to self-renew, play a key role in occurrence, recurrence, metastasis, and chemotherapy resistance of breast cancer.^{56,57} CD44⁺/CD24^{–/low} and aldehyde dehydrogenase 1 (ALDH1) were identified as markers of BCSCs.^{56,58} Kang et al found that ER- α 36 was highly expressed in ALDH1-positive SK-BR-3 cells. Additionally, the depletion of ER- α 36 reduced the growth rate and proportion of ALDH1^{high} cells, indicating that ER- α 36 contributed to the proliferation and maintenance of stem-like cells.⁵³ Deng et al observed that estrogen significantly increased the population of CD44⁺/CD24^{–/low} cells in MCF-7 and T47D breast cancer cells, but failed to do so after knocking down ER- α 36, suggesting that estrogen positively regulated BCSCs through the ER- α 36-mediated signaling pathway.³⁴ Furthermore, they found that overexpression of ER- α 36 tended to promote self-renewal and tumor-seeding efficiency of BCSCs through the PI3K/

AKT/glycogen synthase kinase 3 β / β -catenin signaling pathway (Figure 2).³⁴

ER- α 36 and endocrine therapy resistance

Endocrine therapy is a dominant anticancer method for ER-positive breast cancer patients. Although aromatase inhibitors are increasingly applied in postmenopausal women,⁵⁹ antiestrogens are still used as first-line therapy, especially when aromatase inhibitors are ineffective. However, less or no response after long-term treatment is the major obstacle in antiestrogen treatment.^{60,61}

Tamoxifen

Tamoxifen is a competitive antagonist of ER- α , effectively blocking traditional genomic signaling pathway mediated by ER- α 66.^{61,62} However, it can activate different kinase cascades and produce second messengers, thus acting as an agonist that is equal to estrogen.⁶³ For example, tamoxifen strongly activates the MAPK/ERK signaling pathway to stimulate cell growth; additionally, it activates the p38/MAPK and SAPK/JNK pathways to induce apoptosis.^{10,64} In recent years, a large number of explorations have concentrated on the association between ER- α 36 and tamoxifen resistance. Due to the unique 27-amino-acid domain, ER- α 36 has a broader ligand-binding spectrum than ER- α 66 so that it can respond to tamoxifen.^{10,24} Usually, ER- α 36 is highly expressed in ER- α 66-negative breast cancer, which is insensitive to endocrine therapy.^{10,28,29} Clinical research demonstrated that high expression of ER- α 36 was associated with decreased tamoxifen sensitivity and poorer survival in ER- α 66-positive breast cancer patients who received tamoxifen as adjuvant therapy.²⁹ Additionally, laboratory studies revealed that the tamoxifen-resistant MCF-7 cell line (MCF-7/TAM) highly expressed ER- α 36.^{22,23,65} However, knocking down ER- α 36 in MCF-7/TAM cells could not restore their sensitivity to tamoxifen completely.²³ This may be because the expression level of ER- α 36 in MCF-7/TAM cells is too high to be completely blocked; besides, biological reprogramming during development of tamoxifen resistance is much more complicated than upregulation of ER- α 36. Furthermore, proteins maintaining the stability and function of ER- α 36 such as SNCG can strengthen tamoxifen resistance mediated by ER- α 36.³³ Therefore, ER- α 36 is an underlying cause of tamoxifen resistance in breast cancer patients.

Previous explorations have demonstrated that ER- α 36 is a potent mediator of membrane-initiated signaling pathways associated with agonist effects of tamoxifen, which leads to

tamoxifen resistance.^{10,24} For example, the MAPK/ERK and PI3K/AKT signaling pathways that are mediated by ER- α 36 induce the expression of protooncogene c-Myc, which has profound mitogenic effects (Figure 4).^{10,44} ER- α 36 also mediates the Src/EGFR/STAT5 pathway regulating activity of the MAPK/ERK signaling pathway and expression of the growth-promoting cyclin D1 (Figure 4).⁶⁶ However, it is still unknown whether other non-genomic pathways, such as protein kinase C, protein kinase A, and calcium channels are involved in tamoxifen resistance.

Further research has revealed that overexpression of EGFR with downregulation of ER- α 66 is responsible for tamoxifen resistance.^{67–69} Li et al established a tamoxifen-resistant MCF-7 cell line (MCF-7/TAM) that had an accelerated proliferation rate as well as enhanced migratory and invasive ability. In MCF-7/TAM cells, ER- α 36 upregulated EGFR expression and downregulated ER- α 66 expression, switching growth status from estrogen-dependent to growth-factor-dependent, which was a critical step in the development of tamoxifen resistance (Figure 4).²³

ICI 182,780 (Fulvestrant, Faslodex®)

ICI 182,780 (Fulvestrant, Faslodex), a ‘pure’ antiestrogen, not only accelerates degradation but also impairs dimerization and nuclear localization of ER- α 66.^{70,71} However, several studies reported that ICI 182,780 failed to induce

degradation of ER- α 36,^{6,72} presumably because ER- α 36 has a truncated ligand-binding domain lacking the last four helices (helix 9–12) of ER- α 66, which are essential for protein degradation induced by ICI 182,780.^{6,73} Therefore, failure of ER- α 36 degradation is a highly possible reason for ICI 182,780 resistance.

Application value of ER- α 36

Metastasis and recurrence are the most intractable issues in the management of cancer; thus, using molecular alteration to assess tumor degree and prognosis is of great significance. A large number of studies have confirmed that ER- α 36-positive breast cancers tend to have more lymph node metastasis, advanced degree, less sensitivity to chemotherapy or endocrine therapy, and poorer outcome than ER- α 36-negative breast cancer. Therefore, ER- α 36 is a promising biomarker for malignant behavior, therapeutic response, and outcome of breast cancer, guiding clinical decisions in early diagnosis, prognosis estimation, and personalized treatment. For example, we can combine the expression of ER- α 66 and ER- α 36 to predict whether endocrine therapy is suitable and effective for breast cancer patients.

Given its important role in carcinogenesis, progression, and therapeutic response of breast cancer, ER- α 36 is emerging as a potential therapeutic target for breast cancer. The surface localization of ER- α 36 makes it possible to generate

Figure 4 ER- α 36 and tamoxifen resistance.

Notes: ER- α 36 is an important cause of tamoxifen resistance. On the one hand, it activates membrane-initiated signaling pathways, such as MAPK/ERK, PI3K/AKT, and Src/EGFR/STAT5 pathways, causing the agonist effect of tamoxifen. On the other hand, ER- α 36 upregulates EGFR and downregulates ER- α 66 switching growth status from estrogen-dependent to growth-factor-dependent. Therefore, ER- α 36 leads to tamoxifen resistance.

Abbreviations: Akt, protein kinase B; EGFR, epidermal growth factor receptor; ER- α 36, estrogen receptor- α 36; ER- α 66, estrogen receptor- α 66; ERK, extracellular signal-regulated kinase; MAPK, mitogen-activated protein kinases; MCF-7/TAM, tamoxifen-resistant MCF-7 cell line; PI3K, phosphatidylinositol 3-kinase; STAT5, signal transducer and activator of transcription 5.

specific surface epitopes for treatment intervention with small molecules or biologics which directly down-modulate the ER- α 36 protein or indirectly block its downstream signaling pathways, such as biological antagonists, monoclonal antibodies, and small kinase inhibitors. Inhibition of ER- α 36 can suppress tumor growth and progression, benefitting breast cancer patients, especially those who are resistant to endocrine therapy. Though few studies have focused on the therapeutic application of ER- α 36 and no clinical trials currently, a targeted agent suitable for clinical testing will be available in the near future. Broussonet flavonoid B, a flavonoid purified from the bark of *Broussonetia papyrifera* (Moraceae), exhibits a more potent growth inhibition effect than tamoxifen in ER-negative breast cancer.^{74,75} It decreases the steady level of ER- α 36 and EGFR and restricts growth of stem-like cells in MDA-MB-231 cells.⁷⁵

Conclusion

ER- α 36 is a variant of ER- α with different characteristics and functions than ER- α 66. It locates on plasma membrane and cytoplasm. Therefore, ER- α 36 mediates the non-genomic signaling pathway and suppresses the traditional genomic signaling pathway. ER- α 36 plays a key role in breast cancer, leading to an accelerated proliferation rate together with enhanced migratory and invasive ability. Additionally, it mediates the agonist action of tamoxifen and switches the growth status of breast cancer cells from estrogen-dependent to growth-factor-dependent, both of which can be the molecular mechanisms of tamoxifen resistance. Furthermore, failure of ER- α 36 degradation is a possible reason for Fulvestrant resistance. Therefore, ER- α 36 is a promising biomarker for tumor behavior and therapeutic response, being a new indicator to select optimal candidates for therapeutic strategies. Additionally, it is a potential therapeutic target to improve survival of breast cancer patients.

These achievements make the heterogeneity of the estrogen effect more clear and put a new insight into the properties of ER- α 36. Additionally, studies of ER- α 36 deepen our knowledge of the carcinogenesis and progression of breast cancer and provide an alternative explanation for endocrine therapy resistance. However, there are still many challenges in the future. More research is needed to clarify the biological function and deep mechanism of ER- α 36. Additionally, developing an effective and specific inhibitor of ER- α 36 is a difficult work. Before the inhibitor is widely used, the inhibitory consequences should be validated regarding proliferation, invasion, migration, and survival of tumor cells and clinical trials with large sample sizes and long-term observation are also needed.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No 81272676), the National Science and Technology Major Project of the Ministry of Science and Technology of China (No 2013ZX09506015), and the Medical Science and Technology Project of Zhejiang Province (No 2011ZDA009).

Disclosure

The authors declare no conflicts of interest in this work.

References

1. Nilsson S, Mäkelä S, Treuter E, et al. Mechanisms of estrogen action. *Physiol Rev*. 2001;81(4):1535–1565.
2. Weihua Z, Andersson S, Cheng G, Simpson ER, Warner M, Gustafsson JA. Update on estrogen signaling. *FEBS Lett*. 2003;546(1):17–24.
3. Kong EH, Pike AC, Hubbard RE. Structure and mechanism of the oestrogen receptor. *Biochem Soc Trans*. 2003;31(Pt 1):56–59.
4. Gustafsson JA. Estrogen receptor beta – a new dimension in estrogen mechanism of action. *J Endocrinol*. 1999;163(3):379–383.
5. Greene GL, Gilna P, Waterfield M, Baker A, Hort Y, Shine J. Sequence and expression of human estrogen receptor complementary DNA. *Science*. 1986;231(4742):1150–1154.
6. Wang Z, Zhang X, Shen P, Loggie BW, Chang Y, Deuel TF. Identification, cloning, and expression of human estrogen receptor- α 36, a novel variant of human estrogen receptor- α 66. *Biochem Biophys Res Commun*. 2005;336(4):1023–1027.
7. Hirata S1, Shoda T, Kato J, Hoshi K. Isoform/variant mRNAs for sex steroid hormone receptors in humans. *Trends Endocrinol Metab*. 2003;14(3):124–129.
8. Murphy LC, Dotzlaw H, Leygue E, Douglas D, Coutts A, Watson PH. Estrogen receptor variants and mutations. *J Steroid Biochem Mol Biol*. 1997;62(5–6):363–372.
9. Zhang QX, Hilsenbeck SG, Fuqua SA, Borg A. Multiple splicing variants of the estrogen receptor are present in individual human breast tumors. *J Steroid Biochem Mol Biol*. 1996;59(3–4):251–260.
10. Wang Z, Zhang X, Shen P, Loggie BW, Chang Y, Deuel TF. A variant of estrogen receptor- α , hER- α 36: transduction of estrogen- and antiestrogen-dependent membrane-initiated mitogenic signaling. *Proc Natl Acad Sci U S A*. 2006;103(24):9063–9068.
11. Li L, Haynes MP, Bender JR. Plasma membrane localization and function of the estrogen receptor α variant (ER46) in human endothelial cells. *Proc Natl Acad Sci U S A*. 2003;100(8):4807–4812.
12. Xie H, Sun M, Liao XB, et al. Estrogen receptor α 36 mediates a bone-sparing effect of 17 β -estradiol in postmenopausal women. *J Bone Miner Res*. 2011;26(1):156–168.
13. Jia S, Zhang X, He DZ, et al. Expression and function of a novel variant of estrogen receptor- α 36 in murine airways. *Am J Respir Cell Mol Biol*. 2011;45(5):1084–1089.
14. Deng H, Huang X, Fan J, et al. A variant of estrogen receptor- α , ER- α 36 is expressed in human gastric cancer and is highly correlated with lymph node metastasis. *Oncol Rep*. 2010;24(1):171–176.
15. Fu Z, Deng H, Wang X, Yang X, Wang Z, Liu L. Involvement of ER- α 36 in the malignant growth of gastric carcinoma cells is associated with GRP94 overexpression. *Histopathology*. 2013;63(3):325–333.
16. Wang X, Deng H, Zou F, et al. ER- α 36-mediated gastric cancer cell proliferation via the c-Src pathway. *Oncol Lett*. 2013;6(2):329–335.
17. Miceli V, Cocciaferro L, Fregapane M, et al. Expression of wild-type and variant estrogen receptor α in liver carcinogenesis and tumor progression. *OMICS*. 2011;15(5):313–317.

18. Tu BB, Lin SL, Yan LY, Wang ZY, Sun QY, Qiao J. ER- α 36, a novel variant of estrogen receptor α , is involved in EGFR-related carcinogenesis in endometrial cancer. *Am J Obstet Gynecol*. 2011;205(3):227.e1–227.e6.
19. Vranic S, Gatalica Z, Deng H, et al. ER- α 36, a novel isoform of ER- α 66, is commonly over-expressed in apocrine and adenoid cystic carcinomas of the breast. *J Clin Pathol*. 2011;64(1):54–57.
20. Zhang XT, Kang LG, Ding L, Vranic S, Gatalica Z, Wang ZY. A positive feedback loop of ER- α 36/EGFR promotes malignant growth of ER-negative breast cancer cells. *Oncogene*. 2011;30(7):770–780.
21. Zheng Y, Zhang J, Xu ZZ, et al. Quantitative profiles of the mRNAs of ER- α and its novel variant ER- α 36 in breast cancers and matched normal tissues. *J Zhejiang Univ Sci B*. 2010;11(2):144–150.
22. Zhang X, Wang ZY. Estrogen receptor- α variant, ER- α 36, is involved in tamoxifen resistance and estrogen hypersensitivity. *Endocrinology*. 2013;154(6):1990–1998.
23. Li G, Zhang J, Jin K, et al. Estrogen receptor- α 36 is involved in development of acquired tamoxifen resistance via regulating the growth status switch in breast cancer cells. *Mol Oncol*. 2013;7(3):611–624.
24. Zhang J, Li G, Li Z, et al. Estrogen-independent effects of ER- α 36 in ER-negative breast cancer. *Steroids*. 2012;77(6):666–673.
25. Yu L, Ke W, Wang Y, et al. Predictive and prognostic value of ER- α 36 expression in breast cancer patients treated with chemotherapy. *Steroids*. 2014;84:11–16.
26. Lin SL, Yan LY, Liang XW, et al. A novel variant of ER- α , ER- α 36 mediates testosterone-stimulated ERK and Akt activation in endometrial cancer Hec1A cells. *Reprod Biol Endocrinol*. 2009;7:102.
27. Pelekanou V, Notas G, Kampa M, et al. ER α 36, a new variant of the ER α is expressed in triple negative breast carcinomas and has a specific transcriptomic signature in breast cancer cell lines. *Steroids*. 2012;77(10):928–934.
28. Lee LM, Cao J, Deng H, Chen P, Gatalica Z, Wang ZY. ER- α 36, a novel variant of ER- α , is expressed in ER-positive and -negative human breast carcinomas. *Anticancer Res*. 2008;28(1B):479–483.
29. Shi L, Dong B, Li Z, et al. Expression of ER- α 36, a novel variant of estrogen receptor α , and resistance to tamoxifen treatment in breast cancer. *J Clin Oncol*. 2009;27(21):3423–3429.
30. Zou Y, Ding L, Coleman M, Wang Z. Estrogen receptor- α (ER- α) suppresses expression of its variant ER- α 36. *FEBS Lett*. 2009;583(8):1368–1374.
31. Kang L, Wang L, Wang ZY. Opposite regulation of estrogen receptor- α and its variant ER- α 36 by the Wilms' tumor suppressor WT1. *Oncol Lett*. 2011;2(2):337–341.
32. Jiang Y, Liu YE, Goldberg ID, Shi YE. Gamma synuclein, a novel heat-shock protein-associated chaperone, stimulates ligand-dependent estrogen receptor α signaling and mammary tumorigenesis. *Cancer Res*. 2004;64(13):4539–4546.
33. Shi YE, Chen Y, Dackour R, et al. Synuclein gamma stimulates membrane-initiated estrogen signaling by chaperoning estrogen receptor (ER)- α 36, a variant of ER- α . *Am J Pathol*. 2010;177(2):964–973.
34. Deng H, Zhang XT, Wang ML, Zheng HY, Liu LJ, Wang ZY. ER- α 36-mediated rapid estrogen signaling positively regulates ER-positive breast cancer stem/progenitor cells. *PLoS One*. 2014;9(2):e88034.
35. Mangelsdorf DJ, Thummel C, Beato M, et al. The nuclear receptor superfamily: the second decade. *Cell*. 1995;83(6):835–839.
36. Wehling M. Specific, nongenomic actions of steroid hormones. *Annu Rev Physiol*. 1997;59:365–393.
37. Revankar CM, Cimino DF, Sklar LA, Arterburn JB, Prossnitz ER. A transmembrane intracellular estrogen receptor mediates rapid cell signaling. *Science*. 2005;307(5715):1625–1630.
38. Kelly MJ, Levin ER. Rapid actions of plasma membrane estrogen receptors. *Trends Endocrinol Metab*. 2001;12(4):152–156.
39. Driggers PH, Segars JH. Estrogen action and cytoplasmic signaling pathways. Part II: the role of growth factors and phosphorylation in estrogen signaling. *Trends Endocrinol Metab*. 2002;13(10):422–427.
40. Bulayeva NN, Wozniak AL, Lash LL, Watson CS. Mechanisms of membrane estrogen receptor- α -mediated rapid stimulation of Ca²⁺ levels and prolactin release in a pituitary cell line. *Am J Physiol Endocrinol Metab*. 2005;288(2):E388–E397.
41. Simoncini T, Mannella P, Genazzani AR. Rapid estrogen actions in the cardiovascular system. *Ann NY Acad Sci*. 2006;1089:424–430.
42. Kang L, Zhang X, Xie Y, et al. Involvement of estrogen receptor variant ER- α 36, not GPR30, in nongenomic estrogen signaling. *Mol Endocrinol*. 2010;24(4):709–721.
43. Chaudhri RA, Olivares-Navarrete R, Cuenca N, Hadadi A, Boyan BD, Schwartz Z. Membrane estrogen signaling enhances tumorigenesis and metastatic potential of breast cancer cells via estrogen receptor- α 36 (ER α 36). *J Biol Chem*. 2012;287(10):7169–7181.
44. Hiscox S, Morgan L, Barrow D, Dutkowskil C, Wakeling A, Nicholson RI. Tamoxifen resistance in breast cancer cells is accompanied by an enhanced motile and invasive phenotype: inhibition by gefitinib ('Iressa', ZD1839). *Clin Exp Metastasis*. 2004;21(3):201–212.
45. Watters JJ, Campbell JS, Cunningham MJ, Krebs EG, Dorsa DM. Rapid membrane effects of steroids in neuroblastoma cells: effects of estrogen on mitogen activated protein kinase signalling cascade and c-fos immediate early gene transcription. *Endocrinology*. 1997;138(9):4030–4033.
46. Márquez DC, Pietras RJ. Membrane-associated binding sites for estrogen contribute to growth regulation of human breast cancer cells. *Oncogene*. 2001;20(39):5420–5430.
47. Lin SL, Yan LY, Zhang XT, et al. ER- α 36, a variant of ER- α , promotes tamoxifen agonist action in endometrial cancer cells via the MAPK/ERK and PI3K/Akt pathways. *PLoS One*. 2010;5(2):e9013.
48. Tong JS, Zhang QH, Wang ZB, et al. ER- α 36, a novel variant of ER- α , mediates estrogen-stimulated proliferation of endometrial carcinoma cells via the PKC δ /ERK pathway. *PLoS One*. 2010;5(11):e15408.
49. Leever SJ, Vanhaesebroeck B, Waterfield MD. Signalling through phosphoinositide 3-kinases: the lipids take centre stage. *Curr Opin Cell Biol*. 1999;11(2):219–225.
50. Vivanco I, Sawyers CL. The phosphatidylinositol 3-Kinase AKT pathway in human cancer. *Nat Rev Cancer*. 2002;2(7):489–501.
51. Fresno Vara JA, Casado E, de Castro J, Cejas P, Belda-Iniesta C, González-Barón M. PI3K/Akt signalling pathway and cancer. *Cancer Treat Rev*. 2004;30(2):193–204.
52. Ji H, Liu YE, Jia T, et al. Identification of a breast cancer-specific gene, BCSG1, by direct differential cDNA sequencing. *Cancer Res*. 1997;57(4):759–764.
53. Kang L, Guo Y, Zhang X, Meng J, Wang ZY. A positive cross-regulation of HER2 and ER- α 36 controls ALDH1 positive breast cancer cells. *J Steroid Biochem Mol Biol*. 2011;127(3–5):262–268.
54. Zhang XT, Ding L, Kang LG, Wang ZY. Involvement of ER- α 36, Src, EGFR and STAT5 in the biphasic estrogen signaling of ER-negative breast cancer cells. *Oncol Rep*. 2012;27(6):2057–2065.
55. Xu BZ, Lin SL, Li M, et al. Changes in estrogen receptor- α variant (ER- α 36) expression during mouse ovary development and oocyte meiotic maturation. *Histochem Cell Biol*. 2009;131(3):347–354.
56. Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci U S A*. 2003;100(7):3983–3988.
57. Charafe-Jauffret E, Ginestier C, Iovino F, et al. Breast cancer cell lines contain functional cancer stem cells with metastatic capacity and a distinct molecular signature. *Cancer Res*. 2009;69(4):1302–1313.
58. Ginestier C, Hur MH, Charafe-Jauffret E, et al. ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell Stem Cell*. 2007;1(5):555–567.
59. Buzdar AU, Cuzick J. Anastrozole as an adjuvant endocrine treatment for postmenopausal patients with breast cancer: emerging data. *Clin Cancer Res*. 2006;12(3 Pt 2):1037s–1048s.
60. Urruticoechea A. The oestrogen-dependent biology of breast cancer. Sensitivity and resistance to aromatase inhibitors revisited: a molecular perspective. *Clin Transl Oncol*. 2007;9(12):752–759.

61. Clarke R, Liu MC, Bouker KB, et al. Antiestrogen resistance in breast cancer and the role of estrogen receptor signaling. *Oncogene*. 2003;22(47):7316–7339.
62. Dutertre M, Smith CL. Molecular mechanisms of selective estrogen receptor modulator (SERM) action. *J Pharmacol Exp Ther*. 2000; 295(2):431–437.
63. Vivacqua A, Bonfiglio D, Recchia AG, et al. The G protein-coupled receptor GPR30 mediates the proliferative effects induced by 17 β -estradiol and hydroxytamoxifen in endometrial cancer cells. *Mol Endocrinol*. 2006;20(3):631–646.
64. Zhang CC, Shapiro DJ. Activation of the p38 mitogen-activated protein kinase pathway by estrogen or by 4-hydroxytamoxifen is coupled to estrogen receptor-induced apoptosis. *J Biol Chem*. 2000;275(1):479–486.
65. Zhao Y, Deng C, Lu W, et al. let-7 microRNAs induce tamoxifen sensitivity by downregulation of estrogen receptor α signaling in breast cancer. *Mol Med*. 2011;17(11–12):1233–1241.
66. Zhang X, Ding L, Kang L, Wang ZY. Estrogen receptor- α 36 mediates mitogenic antiestrogen signaling in ER-negative breast cancer cells. *PLoS One*. 2012;7(1):e30174.
67. Gutierrez MC, Detre S, Johnston S, et al. Molecular changes in tamoxifen-resistant breast cancer: relationship between estrogen receptor, HER-2, and p38 mitogen-activated protein kinase. *J Clin Oncol*. 2005;23(11):2469–2476.
68. Johnston SR, Sacconi-Jotti G, Smith IE, et al. Changes in estrogen receptor, progesterone receptor, and pS2 expression in tamoxifen-resistant human breast cancer. *Cancer Res*. 1995;55(15):3331–3338.
69. Massarweh S, Osborne CK, Creighton CJ, et al. Tamoxifen resistance in breast tumors is driven by growth factor receptor signaling with repression of classic estrogen receptor genomic function. *Cancer Res*. 2008;68(3):826–833.
70. Fawell SE, White R, Hoare S, Sydenham M, Page M, Parker MG. Inhibition of estrogen receptor-DNA binding by the “pure” antiestrogen ICI 164,384 appears to be mediated by impaired receptor dimerization. *Proc Natl Acad Sci U S A*. 1990;87(17):6883–6887.
71. Dauvois S, Danielian PS, White R, Parker MG. Antiestrogen ICI 164,384 reduces cellular estrogen receptor content by increasing its turnover. *Proc Natl Acad Sci U S A*. 1992;89(9):4037–4041.
72. Kang L, Wang ZY. Breast cancer cell growth inhibition by phenethyl isothiocyanate is associated with down-regulation of oestrogen receptor- α 36. *J Cell Mol Med*. 2010;14(6B):1485–1493.
73. Mahfoudi A, Roulet E, Dauvois S, Parker MG, Wahli W. Specific mutations in the estrogen receptor change the properties of antiestrogens to full agonists. *Proc Natl Acad Sci U S A*. 1995;92(10):4206–4210.
74. Guo M, Wang M, Zhang X, Deng H, Wang ZY. Brousoflavonol B restricts growth of ER-negative breast cancer stem-like cells. *Anticancer Res*. 2013;33(5):1873–1879.
75. Guo M, Wang M, Deng H, Zhang X, Wang ZY. A novel anticancer agent Brousoflavonol B downregulates estrogen receptor (ER)- α 36 expression and inhibits growth of ER-negative breast cancer MDA-MB-231 cells. *Eur J Pharmacol*. 2013;714(1–3):56–64.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress