Effects of immunosuppressive treatment on protein expression in rat kidney

Karolina Kędzierska¹
Katarzyna Sporniak-Tutak²
Krzysztof Sindrewicz²
Joanna Bober³
Leszek Domański¹
Mirośław Parafiniuk⁴
Elżbieta Urańska⁵
Andrzej Ciechanowicz⁶
Maciej Domański¹
Tomasz Smektała²
Marek Masiuk⁵
Wiesław Skrzypczak⁶
Małgorzata Ożgo⁶
Joanna Kabat-Koperska¹
Kazimierz Ciechanowski¹

¹Department of Nephrology, Transplantology, and Internal Medicine, ²Department of Dental Surgery, ³Department of Medical Chemistry, ⁴Department of Forensic Medicine, ⁵Department of Pathomorphology, Pomeranian Medical University, ⁶Department of Physiology, Cytobiology, and Proteomics, West Pomeranian University of Technology, Szczecin, Poland

Correspondence: Karolina Kędzierska
Department of Nephrology,
Transplantology and Internal Medicine,
Powstańców Wlkp 72, 70-111 Szczecin,
Pomeranian Medical University, Poland
Tel +48 91 466 11 96
Fax +48 91 466 11 96
Email karolina.kedzierska@interia.pl

Abstract: The structural proteins of renal tubular epithelial cells may become a target for the toxic metabolites of immunosuppressants. These metabolites can modify the properties of the proteins, thereby affecting cell function, which is a possible explanation for the mechanism of immunosuppressive agents' toxicity. In our study, we evaluated the effect of two immunosuppressive strategies on protein expression in the kidneys of Wistar rats. Fragments of the rat kidneys were homogenized after cooling in liquid nitrogen and then dissolved in lysis buffer. The protein concentration in the samples was determined using a protein assay kit, and the proteins were separated by two-dimensional electrophoresis. The obtained gels were then stained with Coomassie Brilliant Blue, and their images were analyzed to evaluate differences in protein expression. Identification of selected proteins was then performed using mass spectrometry. We found that the immunosuppressive drugs used in popular regimens induce a series of changes in protein expression in target organs. The expression of proteins involved in drug, glucose, amino acid, and lipid metabolism was pronounced. However, to a lesser extent, we also observed changes in nuclear, structural, and transport proteins' synthesis. Very slight differences were observed between the group receiving cyclosporine, mycophenolate mofetil, and glucocorticoids (CMG) and the control group. In contrast, compared to the control group, animals receiving tacrolimus, mycophenolate mofetil, and glucocorticoids (TMG) exhibited higher expression of proteins responsible for renal drug metabolism and lower expression levels of cytoplasmic actin and the major urinary protein. In the TMG group, we observed higher expression of proteins responsible for drug metabolism and a decrease in the expression of respiratory chain enzymes (thioredoxin-2) and markers of distal renal tubular damage (heart fatty acid-binding protein) compared to expression in the CMG group. The consequences of the reported changes in protein expression require further study.

Keywords: Proteomics, drug effects, immunosuppression, rats

Introduction

Modern immunosuppressive regimens have contributed to a reduction in the incidence of acute rejection episodes, which has resulted in very good 1-year renal graft survival, reaching 90%–95%. Interestingly, the present average half-life of a transplanted kidney in the United States is only approximately 1 year longer than the half-life in 1988. Currently, based on data published by the Collaborative Transplant Study, the half-life reaches approximately 16 years in Europe, whereas in the United States, it is only 8.8 years.^{1,2} The data are a stimulus for seeking the causes of this phenomenon. Traditionally, the side effects of immunosuppressive drugs have been considered to be one of the main reasons for this situation.

The process of urine concentration in the renal tubules directly exposes cells to the harmful influence of toxins contained in the tubular fluid. Toxins permanently damage the cells of the renal tubules, especially in the proximal part, which shows the highest metabolic activity and the most advanced transport mechanisms. The S3 segment of the proximal tubules contains peroxisomes and amino acid-degrading enzymes. Moreover, several enzymes involved in the metabolism of drugs, such as oxidases and cytochrome p450, have high activity in the region.³ Extremely intense biochemical changes occur in the proximal tubules. Proximal convoluted tubule cells contain many mitochondria and an extensive Golgi system. Mitochondrial oxygen metabolism is the primary source of energy (concentrated in ATP) in this part of a nephron. Renal tubular epithelial cells' structural proteins may become a target for immunosuppressive drugs' toxic metabolites. The toxins can modify the properties of the proteins, thereby changing the cells' functions. This phenomenon likely explains the mechanism of immunosuppressive agents' toxicity in a graft.

Asif et al studied the effect of mycophenolate mofetil on protein expression in the kidneys of Wistar rats. They found that the treatment mainly affected proteins involved in drug metabolism, but there were also certain changes in proteins that are important in glucose, amino acid, and lipid metabolism and in redox reactions.⁴ In turn, Lamoureux et al compared the effects of cyclosporin A and tacrolimus on human embryonic kidney (HEK) cells and proved that cyclosporin A affected protein expression more significantly than tacrolimus did.⁵ In the same study, respiratory chain and cytoskeletal protein expression was also affected. Klawitter et al observed similar changes in protein expression in Wistar rats treated with cyclosporin A and tacrolimus.⁶ In the cyclosporine group, compared to the control group, higher expression of cytoskeletal proteins and renal aminoacylase (which is important in renal drug metabolism) was observed. The expression of proteins participating in redox reactions was decreased. No similar changes were found in rats treated with tacrolimus.

The aim of our study was to investigate the effect of drugs used in standard immunosuppressive regimens on protein expression in the rat kidney. We created a model of immunosuppressant action that may be compared to chronic immunosuppressive therapy in humans. We chose a physiological enteric route of drug administration, and the duration of the study was extended to 6 months. The immunosuppressive regimens most commonly used in clinical practice

were examined. The medication doses used during the study reached levels in the therapeutic range.

Materials and methods

Study design

The study was conducted on 18 male Wistar rats obtained from a licensed breeder (the Institute of Occupational Medicine, Lodz, Poland). At the start of the experiments, the rats were 14 weeks old. The animals had genetic and health certificates issued by a veterinarian. The experiments were approved by the Local Ethical Committee for Experiments on Animals in Szczecin (No 06/08, dated February 4, 2008, and No 24/08, dated November 24, 2008).

The animals were housed in cages, with six rats per cage, in the Animal Facility of Pomeranian Medical University. The room humidity was approximately 55%, and the air temperature was $22^{\circ}\text{C} \pm 2^{\circ}\text{C}$. The lighting, on a 12/12-hour cycle, was controlled by automatic timers. Before the study, all animals were weighed, and their mean weight was 305 g. During 2 weeks of adaptation, the animals were fed specialized laboratory diet LSM (1,474 kJ/100 g, 17.6% protein, Agro Pol, Motycz, Lublin, Poland) and given water to drink *ad libitum*. All animals survived the adaptation period.

The experiments were performed using the pharmaceutical form of each drug. The animals received a drug in a ball of bread orally. The drug doses were based on data available in the literature.^{7–11} The doses used in the study were as follows: tacrolimus (Prograf, Astellas Pharma, Warsaw, Poland): 4 mg/kg/day; mycophenolate mofetil (CellCept, Roche Registration Limited, Welwyn Garden City, Great Britain): 20 mg/kg/day; cyclosporin A (Sandimmun Neoral, Novartis Pharma GmbH, Nuremberg, Germany): 5 mg/kg/day, and prednisone (Encorton, Polfa, Pabianice, Poland): 4 mg/kg/day. The animals received medication every 24 hours for 6 months. After 3 months of study, the animals were weighed again, and each medication dose was adequately adjusted based on the changed weight. Six rats did not receive treatment, as they formed a control group. A diagram of the study is presented in Table 1.

All 18 rats completed the study. After 6 months of study, the animals were anesthetized by intraperitoneal ketamine hydrochloride injection (50 mg/kg) and blood samples were obtained to determine drug levels and urea and creatinine concentration. Their body weight was also measured. Subsequently, necropsies were performed, and the kidneys were harvested for histopathological examination. The right kidney was placed in a vat of liquid nitrogen, and the left one was placed in a formalin solution.

Table I Study design

Group	Glucocorticosteroids (G)	Tacrolimus (T)	Cyclosporin A (C)	Mycophenolate mofetil (M)
Control (n=6)	–	–	–	–
CMG (n=6)	+	–	+	+
TMG (n=6)	+	+	–	+

Note: The abbreviations of drugs used for naming groups are in brackets.

Determination of urea and creatinine concentrations

The concentration of urea and creatinine were determined in blood serum, and glomerular filtration rate (GFR) was calculated from obtained results. The concentration of urea and creatinine were determined by the colorimetric method two-point type “fix time” using the reagent kits (Aqua Med, Lodz, Poland); results were reported in mg/dL.

Determination of drug concentrations

Drug concentrations were determined in accordance with the literature after 4 hours of enteral administration.^{11,12}

The concentrations of tacrolimus and cyclosporin A were determined in the rats' whole blood. The test was performed in the Clinical Central Laboratory in Szczecin. An IMx assay, based on a microparticle enzyme immunoassay, was used to determine the tacrolimus level. The test was performed using an Abbott analyzer (Abbott Laboratories, Abbott Park, IL, USA).

The concentration of cyclosporin A was determined using an Abbott AxSYM assay, which is a fluorescence-based method (a fluorescence polarization immunoassay).

Sample preparation

Dissected tissue fragments were washed twice with 0.9% NaCl (0°C) and thereafter twice with 20 mM Krebs-HEPES buffer (Sigma-Aldrich Co., St Louis, MO, USA; 0°C, pH 7.4). Subsequently, the tissue fragments were placed in liquid nitrogen, homogenized, and dissolved in lysis buffer containing 5 M urea, 2 M thiourea, 4% CHAPS, 40 mM Tris, and 0.2% carrier ampholyte. The samples prepared in that manner were then stored (–80°C) until further analysis.

Two-dimensional electrophoresis mass spectrometry

Prior to analysis, the samples were thawed at 0°C, precipitated with four volumes of cold acetone (–20°C), and centrifuged (15,000 rpm/30 minutes). The protein pellets obtained in this way were later dissolved in lysis buffer (5 M urea, 2 M

thiourea, 4% w/v CHAPS, 40 mM Tris, 0.2% w/v 3–10 ampholytes, and 2 mM TBP; Sigma-Aldrich Co.). Isoelectric focusing was performed (Protean, IEF Cell, Bio-Rad, Hercules, CA, USA) using 3–10, 17 cm NL ReadyStrip IPG Strips (Bio-Rad) (400 µg protein/400 µL rehydration buffer) over a total of 92400 Vh. The focused IPG strips were reduced with dithiothreitol (DTT) in equilibration buffer (6 M urea, 0.5 M Tris/HCl, pH 6.8, 2% w/v SDS, 30% w/v glycerol, and 1% w/v DTT; Sigma-Aldrich Co.) for 15 minutes and then alkylated with iodoacetamide (2.5% w/v) for 20 minutes at ambient temperature. After the equilibration process, sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) was performed on large-format 12% SDS polyacrylamide gels at 40 V for 1 hour and subsequently at 120 V for 18 hours at 10°C. After two-dimensional electrophoresis (2-DE) separation, the gels were stained for 72 hours with colloidal Coomassie Brilliant Blue G-250. Gel images were acquired (GS-800 Calibrated Densitometer, Bio-Rad). The 2-D image computer analysis was performed using PDQuest Analysis software version 8.01 Advanced (Bio-Rad). Analytical procedures performed on each gel included spot background subtraction, spot detection, and matching. Normalization of each individual spot was performed using local regression model. Coefficient of variation was calculated for replicate groups. Significance of protein expression changes was measured by the aid of integrated with the software statistic tool based on Student's *t*-test.

Protein identification was performed with the aid of matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MS), as previously described by Skrzypczak et al.¹³

Histopathological evaluation

The study was blinded. A semiquantitative method was used for sample evaluation. In each sample, ten consecutive vision fields were assessed at 10×20 magnification while maintaining a margin of approximately 200 µm from the edge of the sample. A microscope (Carl Zeiss, Oberkochen, Germany) equipped with a camera was used. Images were recorded

using Aperio ImageScope v9 software (Aperio Technologies, Inc, Vista, CA, USA). Two experienced pathologists independently assessed the histological preparations stained with hematoxylin/eosin (HE) and periodic acid–Schiff (PAS). The material was evaluated in terms of potential acute or chronic inflammatory changes within the renal tubules, glomeruli, and vessels in accordance with a modified version of the Banff 2009 criteria (we omitted the criteria for graft rejection [Table S1]).¹⁴

Results

Rat body weight and laboratory blood test results of rats

Analyzing obtained results, we found lower body weight in rats receiving immunosuppressive therapy. Rats from the control group reached the highest weight (median 562.5 g); lower weight was observed in the cyclosporine, mycophenolate mofetil, and glucocorticoids (CMG) group (554 g), and the lowest in the group receiving tacrolimus, mycophenolate mofetil, and glucocorticoids (TMG) (median, 482.5 g). This difference was not statistically significant. Blood biochemical parameters of rats were also analyzed. We found significantly lower serum urea in the CMG group compared to control ($P < 0.01$). There was no significant difference in the concentration of creatinine and GFR between examined groups and control (Table 2). The results of drug concentration are shown in Table 3.

Table 2 Body weight and laboratory blood test results of rats in the treatment groups

Parameter/group	Control	CMG	TMG
Body mass (g)			
av $\pm$ SD	573.2 $\pm$ 85.6	544 $\pm$ 50.2	482.5 $\pm$ 66.7
Median	562.5	554	482.5
Min/max	474–700	481–600	405–580
Creatinine (mg/dL)			
av $\pm$ SD	0.78 $\pm$ 0.16	0.72 $\pm$ 0.06	0.68 $\pm$ 0.06
Median	0.78	0.71	0.69
Min/max	0.57–0.95	0.65–0.8	0.6–0.77
GFR (mL/min)			
av $\pm$ SD	2.99 $\pm$ 1.52	3.19 $\pm$ 0.64	3.66 $\pm$ 0.79
Median	2.68	3.2	3.47
Min/max	1.59–5.5	2.44–4.09	2.71–4.89
Urea (mg/dL)			
av $\pm$ SD	71.9 $\pm$ 24.6	24.9 $\pm$ 11.8	73.4 $\pm$ 17.8
Median	71.88	22.67*	74.41
Min/max	37.5–106.3	12.5–43.8	50–100

Note: * $P < 0.01$ vs control.

Abbreviations: av, average; CMG, cyclosporine, mycophenolate mofetil, and glucocorticoids; GFR, glomerular filtration rate; max, maximum; min, minimum; SD, standard deviation; TMG, tacrolimus, mycophenolate mofetil, and glucocorticoids.

Table 3 Medication trough level in examined rats

	Group CMG (n=6)	Group TMG (n=6)	Control (n=6)
Cyclosporin A (ng/mL)	785.2 $\pm$ 83.3	–	–
Tacrolimus (ng/mL)	–	14.1 $\pm$ 13.1	–

Note: Results are presented as mean $\pm$ standard deviation.

Abbreviations: CMG, cyclosporine, mycophenolate mofetil, and glucocorticoids; TMG, tacrolimus, mycophenolate mofetil, and glucocorticoids.

Proteomic analysis

After performing 2-DE, we separated 75 spots, which corresponded to protein expression differences between the two groups (Figure 1).

Using MS, we identified and described 56 of the 75 proteins. Their names, masses in Da, isoelectric points, and functions are presented in Table S2.

Most of the identified proteins (27%) belonged to a group of enzymes involved in glucose, amino acid, and lipid metabolism. Approximately 25% of the identified proteins were involved in redox reactions and the intracellular respiration process. Approximately 14% of the identified proteins played a role in drug metabolism. In total, 13% of the identified proteins were albumins, and 5% were cytoskeletal structural proteins. The remaining 16% of the identified proteins were nuclear and ribosomal proteins, among others.

We then performed a detailed analysis of all of the proteins whose expression was significantly different between the study groups. Proteins that could not be identified were characterized based on sample-specific point (SSP).

The lowest number of proteins distinguished the CMG group from the control group (Table S3). We observed the most significant differences between the control group and the TMG group (Table S4) and between the CMG and the TMG groups (Table S5).

In the CMG group, which received cyclosporin A, mycophenolate mofetil (MMF), and corticosteroids, there was a significant increase in the expression of albumin and the 5A subunit of mitochondrial cytochrome c oxidase (belonging to a group of respiratory chain proteins) in comparison with expression in the control group. In the CMG group, we observed a significant decrease in the expression of alpha enolase and 3-hydroxyisobutyrate dehydrogenase, which are both very important in glucose, amino acid, and lipid metabolism. A significant decrease in the expression of alcohol dehydrogenase was also observed in the CMG group. Proteins with spot numbers of 47, 48, and 49 were specifically expressed in the CMG group.

There were many significant differences in protein expression between the TMG group and the control group (Table S3).

Figure 1 Two-dimensional electrophoresis of rat kidney proteins.

Note: The arrows show the specific spot numbers of proteins.

In the former group, we observed increased expression of the following proteins:

1. A group of redox enzymes and components of respiratory chain proteins (mitochondrial transferase one succinyl-CoA, the pyruvate dehydrogenase complex, malate dehydrogenase)
2. Enzymes involved in glucose, lipid, and amino acid metabolism (mitochondrial aspartate aminotransferase, 17- β -hydroxysteroid dehydrogenase, triose-phosphate isomerase, glutamine synthetase).

In addition, we noticed increased expression of 60S acidic ribosomal protein and the unidentified proteins SSP 3404, SSP 7304, and SSP 7511. In the TMG group, we also found increased expression of the glutathione S-transferase alpha-3 subunit, whereas the expression of the P subunit stopped.

In the TMG group, compared to the control group, we observed significantly decreased expression of albumin; redox enzymes and components of the respiratory chain (NADH ubiquinone oxidoreductase, the F subunit of the V-type proton ATPase, the 5A subunit of mitochondrial cytochrome c oxidase); proteins involved in the metabolism of glucose, lipids, and amino acids (heart fatty acid-binding protein [H-FABP]), cytoskeletal proteins (myosin light polypeptide, cytoplasmic actin 1), nuclear and ribosomal

proteins, and others (the major urinary protein [MUP] complex, the target of rapamycin complex 2 subunit MAPKAP1, protein regulators of microtubule dynamics). Additionally, the expression of several unidentified proteins (SSP numbers 3815, 4807, 5801, 5804, 5805, 5807, 5808, 5811, 5901, 6405, 9107) was lower in this group.

We also compared the expression of proteins in the CMG and TMG groups. We found several significant differences, and the groups differed in the expression of 38 proteins overall (Table S4). Differences in the expression of the proteins from spots 51 to 75 were found only between the CMG and the TMG groups.

In the TMG group, compared to the CMG group, the expression of the following proteins increased significantly: proteins in the redox enzyme group; components of the respiratory chain (mitochondrial malate dehydrogenase, mitochondrial enoyl-CoA hydratase, a protein associated with phosphotriesterase, cytoplasmic isocitrate dehydrogenase); proteins involved in the metabolism of glucose, lipids, and amino acids (triose-phosphate isomerase, 17- β -hydroxysteroid dehydrogenase X, glutamine synthetase); a protein involved in the metabolism of drugs (aminoacylase 1A); and others (phosphatidylethanolamine-binding protein and the unidentified proteins SSP 3404 and SSP 3412).

Many more proteins in the TMG group had lower expression compared to levels in the CMG group. The most significant differences that we observed were in the expression of not only redox enzymes and components of the respiratory chain (thioredoxin, glucose-regulated protein 78 kDa, mitochondrial aconitase, the 5A subunit of mitochondrial cytochrome c oxidase, the 6A1 subunit of mitochondrial cytochrome c oxidase) but also proteins involved in the metabolism of glucose, lipids, and amino acids (pyridoxine kinase homologue, lambda crystallin, mitochondrial aspartate aminotransferase, H-FABP); proteins involved in the metabolism of drugs (selenium-binding protein); and other proteins (transthyretin, the MUP), including unidentified ones (SSP 3815, 4903, 5801, 5805, 5807, 3412, and 4205).

In the compared groups, the expression of two proteins was ambivalent: the mitochondrial aspartate aminotransferase spot 2 (SSP 507) was higher in the TMG than in the CMG group, but the animals in the TMG group did not demonstrate expression of mitochondrial aspartate aminotransferase spot 51 (SSP 1405). Similarly, the expression of aminoacylase 1A spot 69 (SSP 5503) was higher in the TMG group, whereas in the case of aminoacylase 1A spot 70 (SSP 5509), higher expression was observed in the CMG group.

In summary, in a proteomic analysis of protein expression in the rat kidney, the least significant differences were observed between the CMG and the control groups. Moreover, in comparison with the control group, we observed higher expression of proteins responsible for drug metabolism (the glutathione S-transferase alpha-3 subunit, triose-phosphate isomerase) and lower cytoplasmic actin and MUP expression in the TMG group. Higher expression of proteins responsible for renal drug metabolism (aminoacylase) was also observed in the TMG group in comparison with the CMG group. Lower expression of respiratory chain enzymes, thioredoxin-2 and markers of distal renal tubular damage (H-FABP) in the TMG group was another important difference between the two treatment groups.

Kidney preparation analysis by light microscopy

The analysis of the HE- and PAS- stained slides obtained from each group showed the following (Figure 2 and 3):

- Control group: Normal glomeruli, tubules, and vessel structures were observed. According to the Banff 2009 criteria, the features were assessed as ct0, Ci0, cg0, ah0, t0, i0.

Figure 2 HE staining of rat kidney.

Notes: TMG group (A); CMG group (B); control group (C). Magnification $\times 40$.

Abbreviations: CMG, cyclosporine, mycophenolate mofetil, and glucocorticoids; HE, hematoxylin/eosin; TMG, tacrolimus, mycophenolate mofetil, and glucocorticoids.

- CMG group: No differences from the control group were found; Banff 2009 criteria: ct0, Ci0, cg0, ah0, t0, i0.
- TMG group: Well-preserved proximal tubules with homogeneous cytoplasm and single cells in the interstitium ($<10\%$) were observed, which was similar to the control group; Banff 2009 criteria: ct0, Ci0, cg0, ah0, t0, i0.

Discussion

This study involved three groups of animals. Two treatment groups corresponded to the immunosuppressant schemes most commonly used in clinical practice. The drugs were dosed to achieve the therapeutic range of trough levels. Other investigators have used toxic doses of medicines.^{15–17} Drug concentrations were determined 4 hours after their enteral supply, which was based on published data.^{11,12} Due to differences from humans' metabolism of immunosuppressants, this timing was considered to be optimal for determining the concentrations of drugs in rats' blood.

In our experiments in rat kidneys, we observed certain changes in the expression of the same proteins as those that Lamoureux et al and Klawitter et al noticed in their studies.^{5,6} Importantly and interestingly, the direction of the fluctuations was often opposite to that found in the cited researchers' experiments.

We proved that the group receiving CMG did not differ a lot from the control group in protein expression. In the course of further analysis, it was observed that the most significant differences in protein expression were observed not only between the control group and the group receiving TMG but also between the two treatment groups (CMG and TMG).

There were many significant differences in protein expression between the TMG and the control groups. The TMG group had higher expression of proteins from the redox enzyme complex, components of the respiratory chain (malate dehydrogenase), and proteins involved in the metabolism of glucose, lipids, and amino acids (triose-phosphate isomerase, mitochondrial aspartate aminotransferase). In addition, the expression of 60S acidic ribosomal protein was higher than in the control group. Compared to the control group, the TMG group also had higher expression of the alpha-3 subunit of glutathione S-transferase, but the expression of the glutathione S-transferase P subunit was lower. Asif found similar changes in rats treated with MMF, including higher expression of malate dehydrogenase, triose-phosphate isomerase, mitochondrial aspartate aminotransferase, and the alpha-1 and alpha-4 subunits of glutathione S-transferase.⁴ Other researchers did not observe such changes.^{5,6}

In the TMG group, the expression of NADH oxidoreductase was significantly lower compared to expression in the control group. Klawitter et al found no change in the expression of this protein in a group receiving tacrolimus.⁶

We also observed a reduction in the expression of H-FABP compared to expression in the control group. H-FABP, in addition to its fatty-acid transport function, is a

Figure 3 PAS staining of rat kidney.

Notes: TMG group (A); CMG group (B); control group (C). Magnification $\times 40$.

Abbreviations: CMG, cyclosporine, mycophenolate mofetil, and glucocorticoids; PAS, periodic acid–Schiff; TMG, tacrolimus, mycophenolate mofetil, and glucocorticoids.

marker of distal renal tubular injury.¹⁸ Lamoureux et al found no effect of tacrolimus on the expression of H-FABP.⁵

In the TMG group, we observed lower expression of cytoplasmic actin and the MUP. Lamoureux et al found no effect of tacrolimus on the expression of actin.⁵ According to

Klawitter's study, tacrolimus does not contribute to altered expression of the MUP.⁶

The TMG group had higher expression of cytoplasmic isocitrate dehydrogenase, triose-phosphate isomerase, and aminoacylase compared to expression in the CMG group. Higher expression of aminoacylase may have a protective role against the toxic metabolites of MMF. Similar results have been presented by Asif et al, who observed higher expression of aminoacylase and triose-phosphate isomerase in the kidneys of rats treated with MMF.⁴ Klawitter did not find any effect of tacrolimus on the expression of NADH.⁶

Many more proteins in the TMG group had lower expression than in the CMG group. This phenomenon was most pronounced among redox enzymes and the components of the respiratory chain (thioredoxin, glucose-regulated protein 78 kDa, the 5A subunit of mitochondrial cytochrome c oxidase, the 6A1 subunit of mitochondrial cytochrome c oxidase). Pyridoxine kinase expression proved to be lower in the TMG group compared to expression in the other group. Lower expression of H-FABP, the MUP, and selenium-binding protein was also observed in this group. Asif et al showed higher expression of selenium-binding protein and pyridoxine kinase in the kidneys of rats treated only with MMF.⁴ Other researchers presented opposite results. Lamoureux et al found no influence of tacrolimus on the expression of thioredoxin and glucose-regulated protein 78 kDa or the 5A subunit of mitochondrial cytochrome c oxidase in HEK cells.⁵

In the present study, we also observed lower expression of the two subunits of the mitochondrial cytochrome c oxidase, which was not observed by other researchers.⁵

In the treatment groups, the expression of two proteins was ambivalent: the expression of mitochondrial aspartate aminotransferase was higher in the TMG group than in the CMG group, but the rats in the TMG group did not demonstrate the expression of mitochondrial aspartate aminotransferase spot 51 (SSP 1405). We observed a similar situation for the expression of aminoacylase 1A spot number 69 (SSP 5503): its activity was higher in the TMG group compared to the CMG group, whereas the pattern of aminoacylase 1A spot number 70 (SSP 5509) expression was the opposite.

Certainly, the results of the proteomic analysis were influenced by the application of three-drug regimens, including calcineurin inhibitors, MMF, and corticosteroids. Several of the changes observed in protein expression have been described previously.⁴⁻⁶ Most of the cited studies were based on single-drug treatment (cyclosporine, tacrolimus, or MMF). Additionally, the treatment groups were mainly compared to a control group, and the evaluation of protein expression

differences between treatment groups was not standard in those studies.⁶ In our study on rats treated with different regimens, the changes in protein expression might have been caused by interactions between immunosuppressants. Despite significant differences in the expression of many proteins, we found no impact of those changes on the histopathological picture. Similar results have been presented by Klawitter.⁶

Summary and conclusion

In the proteomic analysis of rat kidneys, less pronounced differences in protein expression were observed between the control group and the CMG group. In the TMG group, higher expression of proteins responsible for renal drug metabolism, such as the glutathione S-transferase alpha-3 subunit and triose-phosphate isomerase, and lower cytoplasmic actin and MUP expression were observed compared to expression in the control group. In the TMG group, higher expression of proteins responsible for renal drug metabolism (aminoacylase) and a decrease in the expression of respiratory chain enzymes (thioredoxin-2) and markers of distal renal tubular damage (H-FABP) were observed in comparison with expression in the CMG group. Medications directly or indirectly (eg, by the intensification of oxidative stress) cause an entire series of changes in the expression of proteins in the kidney. Treatment schemes based on cyclosporin A have a much lower impact on the expression of proteins within the kidney compared to tacrolimus-based regimens. MMF in combination with tacrolimus has its own additional effect on the expression of kidney proteins in rats. This regimen causes a very significant increase in the expression of proteins responsible for drug metabolism and a decrease in the expression of proteins that are part of the respiratory chain and of markers of renal tubular damage. Despite the substantial differences in the expression of many proteins, there were no histopathological or functional changes within the examined kidneys.

Acknowledgments

Work was funded by the statutory budget of the Pomeranian Medical University.

Disclosure

The authors declare no conflicts of interest in this work.

References

1. Meier-Kriesche HU, Schold JD, Kaplan B. Long-term renal allograft survival: have we made significant progress or is it time to rethink our analytic and therapeutic strategies? *Am J Transplant*. 2004;4(8): 1289–1295.

2. Brennan DC. Long-term trends in allograft survival. *Adv Chronic Kidney Dis*. 2006;13(1):11–17.
3. Myśliwiec M. *Nefrologia. Wielka Interna*. Warszawa: Medical Tribune Poland; 2009:458–459. Polish.
4. Asif AR, Armstrong VW, Volland A, Wieland E, Oellerich M, Shipkova M. Proteins identified as targets of the acyl glucuronide metabolite of mycophenolic acid in kidney tissue from mycophenolate mofetil treated rats. *Biochimie*. 2007;89(3):393–402.
5. Lamoureux F, Mestre E, Essig M, Sauvage FL, Marquet P, Gastinel LN. Quantitative proteomic analysis of cyclosporine-induced toxicity in a human kidney cell line and comparison with tacrolimus. *J Proteomics*. 2011;75(2):677–694.
6. Klawitter J, Klawitter J, Kushner E, et al. Association of immunosuppressant-induced protein changes in the rat kidney with changes in urine metabolite patterns: a proteo-metabonomic study. *J Proteome Res*. 2010;9(2):865–875.
7. van Westrhenen R, Aten J, Hajji N, et al. Cyclosporin A induces peritoneal fibrosis and angiogenesis during chronic peritoneal exposure to a glucose-based, lactate-buffered dialysis solution in the rat. *Blood Purif*. 2007;25(5–6):466–472.
8. Joffe I, Katz I, Sehgal S, et al. Lack of change of cancellous bone volume with short-term use of the new immunosuppressant rapamycin in rats. *Calcif Tissue Int*. 1993;53(1):45–52.
9. Jolicœur EM, Qi S, Xu D, Dumont L, Daloze P, Chen H. Combination therapy of mycophenolate mofetil and rapamycin in prevention of chronic renal allograft rejection in the rat. *Transplantation*. 2003;75(1):54–59.
10. Katz IA, Takizawa M, Jaffe II, Stein B, Fallon MD, Epstein S. Comparison of the effects of FK 506 and cyclosporine on bone mineral metabolism in the rat. *Transplantation*. 1991;52(3):571–574.
11. Ma Y, Kobayashi T, Kuzuya T, et al. Is absorption profile of cyclosporine really important for effective immunosuppression? *Biol Pharm Bull*. 2006;29(2):336–342.
12. Schmitz V, Klawitter J, Bendrick-Pearl J, et al. Metabolic profiles in urine reflect nephrotoxicity of sirolimus and cyclosporine following rat kidney transplantation. *Nephron Exp Nephrol*. 2009;111(4):e80–e91.
13. Skrzypczak WF, Ozgo M, Lepczynski A, Herosimczyk A. Defining the blood plasma protein repertoire of seven day old dairy calves – a preliminary study. *J Physiol Pharmacol*. 2011;62(3):313–319.
14. Perkowska-Ptasińska A. [Diagnosis of changes in the transplanted kidney]. *Pol J Pathol*. 2011;62(2 Suppl 1):s111–s149. Polish.
15. Shing CM, Fassett RG, Brown L, Coombes JS. The effects of immunosuppressants on vascular function, systemic oxidative stress and inflammation in rats. *Transpl Int*. 2012;25(3):337–346.
16. Rovira J, Marcelo Arellano E, Burke JT, et al. Effect of mTOR inhibitor on body weight: from an experimental rat model to human transplant patients. *Transpl Int*. 2008;21(10):992–998.
17. Sánchez-Pozos K, Lee-Montiel F, Pérez-Villalva R, et al. Polymerized type I collagen reduces chronic cyclosporine nephrotoxicity. *Nephrol Dial Transplant*. 2010;25(7):2150–2158.
18. Alter ML, Kretschmer A, Von Websky K, et al. Early urinary and plasma biomarkers for experimental diabetic nephropathy. *Clin Lab*. 2012;58(7–8):659–671.

Supplementary materials

Table SI Evaluation criteria

Score	ct (% of the area of cortical tubules)	ci (% of cortical area)	cg (double contours in % of peripheral capillary loops in the most affected of nonsclerotic glomeruli)	t (number of cells/tubular cross section or 10 tubular cells)	i (% of parenchyma inflamed)	ah
0	0	<5	<10	0	<10	No PAS-positive hyaline thickening
1	<25	6–25	10–25	1–4	10–25	Mild to moderate PAS-positive hyaline thickening in at least one arteriole
2	26–50	26–50	26–50	5–10	26–50	Moderate to severe PAS-positive hyaline thickening in more than one arteriole
3	>50	>50	>50	>10 or the presence of at least moderate interstitial inflammation	>50	–

Abbreviations: ah, arteriolar hyaline thickening; cg, allograft glomerulopathy; ci, interstitial fibrosis; ct, tubular atrophy; i, interstitial inflammation; PAS, periodic acid–Schiff; t, tubulitis.

Table S2 Proteins identified in the study groups of rats (control group, CMG, and TMG)

Spot	Protein name	Protein sequence coverage, %	Score	Nominal mass (Da)	Calculated pI
Redox and respiratory chain					
73	Thioredoxin	88	112	12,008	4.80
5	Malate dehydrogenase, mitochondria	69	177	36,117	8.93
44	V-type proton ATPase subunit F	69	80	13,362	5.52
71	N(G),N(G)-dimethylarginine dimethylaminohydrolase I	66	200	31,805	5.75
63	Cytochrome c oxidase subunit 6A1, mitochondrial	61	58	12,293	9.30
61	Isocitrate dehydrogenase cytoplasmic	56	146	47,047	6.53
72	78 kDa glucose-regulated protein	49	269	72,473	5.07
56	Aconitate hydratase, mitochondria	47	274	86,121	7.87
58	Enoyl-CoA hydratase, mitochondria	47	125	31,895	8.39
36	Pyruvate dehydrogenase E1 component subunit beta, mitochondrial	44	106	39,299	6.20
40	NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial	37	105	80,331	5.65
9	Succinyl-CoA:3-ketoacid coenzyme A transferase I, mitochondrial	33	57	56,624	8.70
43	Cytochrome c oxidase subunit 5A, mitochondrial	28	73	16,347	6.08
38	60 kDa heat shock protein, mitochondrial	27	64	61,088	5.91
Albumin					
29	Serum albumin	59	246	70,682	6.09
22		58	285	70,682	6.09
23		58	265	70,682	6.09
30		54	222	70,682	6.09
17		52	228	70,682	6.09
34		52	219	70,682	6.09
18		42	137	70,682	6.09
Glucose, amino acid, and lipid metabolism					
57	Triosephosphate isomerase	74	180	27,345	6.89
7		63	180	27,345	6.89
4	3-hydroxyacyl-CoA dehydrogenase type-2	62	114	27,343	8.91
51	Aspartate aminotransferase, mitochondria	61	199	47,683	9.13
68	Lambda-crystallin homolog	61	161	35,717	5.94
32	Fatty acid-binding protein, heart	60	105	14,766	5.90
48	Alpha-enolase	60	191	47,440	6.16
2	Aspartate aminotransferase, mitochondria	53	162	47,683	9.13
52	Fructose-bisphosphate aldolase B	50	139	40,049	8.66
15	Pyridoxal kinase	43	90	35,114	6.32
8	Fumarylacetoacetase	36	73	46,231	6.67
11	Glutamine synthetase	34	110	42,982	6.64
49	3-hydroxyisobutyrate dehydrogenase, mitochondria	33	88	35,679	8.73
53	Alanine aminotransferase 2, mitochondria	31	60	57,905	8.33
21	Malate dehydrogenase, cytoplasmic	30	59	36,631	6.16
Drug metabolism					
70	Aminoacylase-1A	70	284	46,060	6.03
69		67	303	46,060	6.03
1	Glutathione S-transferase alpha-3	66	138	25,360	8.78
59	Phosphotriesterase-related protein	65	177	39,462	6.40
47	Alcohol dehydrogenase [NADP(+)]	64	173	36,711	6.84
65	Selenium-binding protein I	55	131	53,069	6.10
6	Glutathione S-transferase P	40	58	23,652	6.89
55	Cytochrome P450 2J3	29	56	58,388	8.87

(Continued)

Table S2 (Continued)

Spot	Protein name	Protein sequence coverage, %	Score	Nominal mass (Da)	Calculated pI
Cytoskeleton protein					
46	Myosin light polypeptide 6	46	74	17,135	4.46
39	Actin, cytoplasmic I	35	77	42,052	5.29
13	Keratin, type I cytoskeletal 10	27	91	56,699	5.10
Nuclear, ribosomal and other protein					
74	Phosphatidylethanolamine-binding protein 1	72	146	20,902	5.48
67	Transthyretin	60	110	15,824	5.77
75	Deoxyuridine 5'-triphosphate nucleotidohydrolase	50	67	22,274	9.01
42	Major urinary protein	49	62	21,009	5.85
3	60S acidic ribosomal protein P0	35	72	34,365	5.91
50	Ribonuclease inhibitor	27	79	51,653	4.67
62	WD repeat-containing protein 78	20	53	90,825	5.30
16	Regulator of microtubule dynamics protein 3	19	56	52,565	4.94
12	Target of rapamycin complex 2 subunit MAPKAP1	15	63	59,519	7.23

Abbreviations: CMG, cyclosporine, mycophenolate mofetil, and glucocorticoids; pI, isoelectric point; TMG, tacrolimus, mycophenolate mofetil, and glucocorticoids.

Table S3 Characteristics of proteins where expression was different in the control group and CMG group

Spot number	SSP	Protein name	Control	CMG	Control vs CMG	P-value
30	5,812	Serum albumin	81.8	113.4	↑	0.05
34	6,801	Serum albumin	72.0	117.9	↑	0.01
43	8,108	Cytochrome c oxidase subunit 5A, mitochondrial	831.8	1,080.2	↑	0.05
47	2,411	Alcohol dehydrogenase [NADP(+)]	1,010.6	564.8	↓	0.01
48	4,518	Alpha-enolase	655.2	420.2	↓	0.05
49	5,301	3-hydroxyisobutyrate dehydrogenase	164.0	79.2	↓	0.05

Abbreviations: CMG, cyclosporine, mycophenolate mofetil, and glucocorticoids; SSP, sample-specific point.

Table S4 Characteristics of proteins where expression was different in the control group and TMG group

Spot number	SSP	Protein name	Control	TMG	Control vs TMG	P-value
1	218	Glutathione S-transferase alpha-3	149.4	204.3	↑	0.05
2	507	Aspartate aminotransferase, mitochondrial	586.6	1,067.6	↑	0.05
3	1,315	60S acidic ribosomal protein P0	48.9	103.6	↑	0.05
4	1,316	3-hydroxyacyl-CoA dehydrogenase type-2	239.5	417.6	↑	0.01
5	1,417	Malate dehydrogenase, mitochondrial	535.1	1,016.4	↑	0.01
6	2,203	Glutathione S-transferase P	98.9		↓	0.05
7	2,311	Triosephosphate isomerase	299.0	456.2	↑	0.05
9	2,719	Succinyl-CoA:3-ketoacid coenzyme A transferase I, mitochondrial	89.2	149.6	↑	0.05
10	3,404	Not identified	112.7	189.3	↑	0.01
11	3,511	Glutamine synthetase	160.0	342.7	↑	0.01
12	3,712	Target of rapamycin complex 2 subunit MAPKAP1	108.0	35.4	↓	0.01
14	3,815	Not identified	90.1	36.5	↓	0.01
16	4,702	Regulator of microtubule dynamics protein 3	98.6	42.1	↓	0.05
18	4,806	Serum albumin	55.6	20.9	↓	0.05
19	4,807	Not identified	37.7	7.7	↓	0.05
22	5,702	Serum albumin	436.1	216.2	↓	0.05
24	5,801	Not identified	37.0	17.3	↓	0.01
25	5,804	Not identified	41.9	15.1	↓	0.01
26	5,805	Not identified	29.0	11.2	↓	0.01
27	5,807	Not identified	33.9	13.4	↓	0.05
28	5,808	Not identified	32.9	10.5	↓	0.05
29	5,811	Not identified	96.0	43.2	↓	0.01
31	5,901	Not identified	44.3	9.6	↓	0.01
32	6,105	Fatty acid-binding protein, heart	653.9	441.6	↓	0.05
33	6,405	Not identified	231.1	152.6	↓	0.05
34	6,801	Serum albumin	72.0	45.4	↓	0.05
35	7,304	Not identified	37.2	67.2	↑	0.01
36	7,410	Pyruvate dehydrogenase E1 component subunit beta, mitochondrial	151.3	278.1	↑	0.05
37	7,511	Not identified	102.7	129.5	↑	0.05
38	7,703	60 kDa heat shock protein, mitochondrial	106.4		↓	0.01
39	7,804	Actin, cytoplasmic I	90.5	22.5	↓	0.05
40	7,806	NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial	162.2	80.6	↓	0.05
42	8,102	Major urinary protein	782.4	377.4	↓	0.05
43	8,108	Cytochrome c oxidase subunit 5A, mitochondrial	831.8	476.0	↓	0.05
44	8,112	V-type proton ATPase subunit F	216.4	107.8	↓	0.05
45	9,107	Not identified	690.4	357.7	↓	0.05
46	9,108	Myosin light polypeptide 6	384.0	196.0	↓	0.05

Abbreviations: SSP, sample-specific point; TMG, tacrolimus, mycophenolate mofetil, and glucocorticoids.

Table S5 Characteristics of proteins where expression was different between the groups CMG and TMG

Spot number	SSP	Protein name	CMG	TMG	CMG vs TMG	P-value
2	507	Aspartate aminotransferase, mitochondria	479.8	1,067.6	↑	0.01
4	1,316	3-hydroxyacyl-CoA dehydrogenase type-2	–	417.6	↑	0.01
5	1,417	Malate dehydrogenase, mitochondria	448.6	1,016.4	↑	0.01
10	3,404	Not identified	105.7	189.3	↑	0.05
11	3,511	Glutamine synthetase	128.2	342.7	↑	0.01
12	3,712	Target of rapamycin complex 2 subunit MAPKAP1	90.0	35.4	↓	0.01
14	3,815	Not identified	69.4	36.5	↓	0.05
15	4,413	Pyridoxal kinase	368.9	139.9	↓	0.01
18	4,806	Serum albumin	58.8	20.9	↓	0.01
20	4,903	Not identified	35.1	8.2	↓	0.01
24	5,801	Not identified	40.1	17.3	↓	0.05
26	5,805	Not identified	27.9	11.2	↓	0.05
27	5,807	Not identified	44.7	13.4	↓	0.05
32	6,105	Fatty acid-binding protein, heart	722.3	441.6	↓	0.05
34	6,801	Serum albumin	117.9	45.4	↓	0.05
42	8,102	Major urinary protein	931.2	377.4	↓	0.05
43	8,108	Cytochrome c oxidase subunit 5A, mitochondrial	1,080.2	476.0	↓	0.05
51	1,405	Aspartate aminotransferase, mitochondria	532.6	–	↓	0.01
53	2,601	Alanine–glyoxylate aminotransferase 2, mitochondria	153.0	–	↓	0.05
55	2,718	Cytochrome P450 2J3	64.8	33.7	↓	0.05
56	2,812	Aconitate hydratase, mitochondria	108.9	83.4	↓	0.05
57	3,206	Triosephosphate isomerase	186.1	340.6	↑	0.01
58	3,304	Enoyl-CoA hydratase, mitochondrial	298.5	563.5	↑	0.05
59	3,411	Phosphotriesterase-related protein	51.8	85.2	↑	0.05
60	3,412	Not identified	186.7	305.2	↑	0.05
61	3,507	Isocitrate dehydrogenase [NADP] cytoplasmic	179.7	501.1	↑	0.05
62	3,806	WD repeat-containing protein 78	100.4	48.8	↓	0.05
63	4,125	Cytochrome c oxidase subunit 6A1, mitochondrial	858.5	418.9	↓	0.01
64	4,205	Not identified	140.9	91.0	↓	0.05
65	4,607	Selenium-binding protein I	199.3	110.5	↓	0.05
67	5,114	Transthyretin	435.8	201.4	↓	0.01
68	5,411	Lambda-crystallin homolog	754.8	433.2	↓	0.01
69	5,503	Aminoacylase-1A	394.7	550.3	↑	0.05
70	5,509	Aminoacylase-1A	169.1	100.9	↓	0.05
72	7,812	78 kDa glucose-regulated protein	69.5	43.0	↓	0.05
73	8,111	Thioredoxin	858.3	262.7	↓	0.05
74	8,204	Phosphatidylethanolamine-binding protein I	181.4	264.7	↑	0.05
75	8,609	Deoxyuridine 5'-triphosphate nucleotidohydrolase	128.9	71.5	↓	0.05

Notes: – indicates no expression; ↓ indicates decreased expression; ↑ increased expression.

Abbreviations: CMG, cyclosporine, mycophenolate mofetil, and glucocorticoids; SSP, sample-specific point; TMG, tacrolimus, mycophenolate mofetil, and glucocorticoids; vs, versus.

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>