Progressive osseous heteroplasia: diagnosis, treatment, and prognosis

Robert J Pignolo¹⁻³
Girish Ramaswamy^{2,3}
John T Fong^{2,3}
Eileen M Shore²⁻⁴
Frederick S Kaplan¹⁻³

¹Department of Medicine,

²Department of Orthopaedic Surgery,

³The Center for Research in FOP and

Related Disorders, ⁴Department of

Genetics, University of Pennsylvania

Perelman School of Medicine,

Philadelphia, PA, USA

Abstract: Progressive osseous heteroplasia (POH) is an ultrarare genetic condition of progressive ectopic ossification. Most cases of POH are caused by heterozygous inactivating mutations of *GNAS*, the gene encoding the alpha subunit of the G-stimulatory protein of adenylyl cyclase. POH is part of a spectrum of related genetic disorders, including Albright hereditary osteodystrophy, pseudohypoparathyroidism, and primary osteoma cutis, that share common features of superficial ossification and association with inactivating mutations of *GNAS*. The genetics, diagnostic criteria, supporting clinical features, current management, and prognosis of POH are reviewed here, and emerging therapeutic strategies are discussed.

Keywords: progressive osseous heteroplasia, *GNAS*, heterotopic ossification

Introduction

Progressive osseous heteroplasia (POH) is an ultrarare genetic condition of progressive extraskeletal bone formation (Online Mendelian Inheritance in Man 166350).¹ POH is clinically suspected by cutaneous ossification, usually presenting in early life, that involves subcutaneous and then subsequently deep connective tissues, including muscle and fascia. Most cases of POH are caused by heterozygous inactivating mutations of *GNAS*, the gene encoding the alpha subunit of the G-stimulatory protein of adenylyl cyclase (Gsα).² POH is among a number of related genetic disorders, including Albright hereditary dystrophy (AHO), pseudohypoparathyroidism (PHP), and primary osteoma cutis (OC), that share common features of superficial heterotopic ossification (HO) in association with inactivating mutations of *GNAS*.³⁻⁵ Although there are similarities among these conditions, POH is distinguished clinically from these related disorders by the deep and progressive nature of the heterotopic bone that forms in POH.⁶ Clinically, POH overlap syndromes are recognized in which both POH-like HO and features of AHO or PHP are present. Most cases of POH, PHP1a (pseudohypoparathyroidism type 1a), and AHO are associated with heterozygous inactivating mutations of the *GNAS* gene, which is transcriptionally regulated through multiple promoters and the production of several transcripts, both protein-coding and noncoding RNAs.^{7,8} The major product of the locus is the G-protein subunit Gsα. Additional regulatory complexity of the *GNAS* locus results from genomic imprinting, which causes allele-specific regulation of transcript expression that influences the spectrum of clinical phenotypes of the *GNAS* inactivation disorders.^{2,9-31}

Correspondence: Robert J Pignolo
Department of Medicine and Department
of Orthopaedic Surgery, University
of Pennsylvania Perelman School of
Medicine, 424B Stemmler Hall,
36th Street and Hamilton Walk,
Philadelphia, PA 19104-6081, USA
Tel +1 215 746 8138
Fax +1 215 573 2133
Email pignolo@mail.med.upenn.edu

characterized of which encodes the ubiquitously expressed α -subunit of the stimulatory G protein ($G\alpha$). Other protein-coding transcripts produce XL α s, the extra-large variant of $G\alpha$ (*Gnasxl* in mice), and NESP55, a neuroendocrine secretory protein (mouse *Nesp*).^{3,40,41} Each of the *GNAS* transcripts are initiated at unique promoters and first exons but share common downstream exons (exons 2–13 in humans and 2–12 in mice) of the *GNAS* locus (Figure 1). Alternative splicing of exon 3 generates short and long forms of both $G\alpha$ and XL α s, and neuronal-specific splicing to include exon N1, which resides between exons 3 and 4, leads to the $G\alpha$ -N1 and XL α s-N1 transcripts that have a truncated C terminus. A second open reading frame of XL α s mRNA produces a protein called ALEX that is unrelated to G-proteins. In addition, the transcripts A/B (mouse exon 1A) and *GNAS* antisense (human *GNAS*-AS1 or mouse *Nespas*) appear to be non-protein-coding transcripts, although translation of A/B is predicted to start at an in-frame ATG start site within exon 2 and to produce a truncated $G\alpha$ isoform.^{3,40–43}

The *GNAS* locus also exhibits genomic imprinting, adding yet another level of regulatory complexity.^{3,40,41,44,45} Allele-specific expression of *GNAS* transcripts is dependent on parent of origin, resulting in transcript expression from only one allele. The effects of preferential expression of one of the two *GNAS* alleles are reflected in the different disease phenotypes that result from *GNAS* inactivation of paternally versus maternally inherited genes. For example, PHP1a is primarily caused by maternally inherited heterozygous mutations in *GNAS* locus, whereas POH is correlated with inactivating mutations in the paternally inherited allele.

XL α s and A/B transcripts are expressed only from the paternally inherited *GNAS* gene copy, whereas NESP55 is synthesized only from the maternally inherited allele. In contrast, $G\alpha$ is expressed biallelically in most tissues, including bone and white adipose tissue. However, $G\alpha$ transcription is regulated by tissue-specific imprinting and is restricted to expression from the maternal allele in tissues including renal proximal tubules, thyroid, pituitary, and gonads.^{3,40,41,44,45}

Various functions have been attributed to *GNAS* transcripts on the basis of phenotypes in mice with specific deletions. Mice that are null for $G\alpha$ are embryonic lethal, whereas heterozygotes show different metabolic and tissue-specific phenotypes according to parent of origin of the mutation. Similar to PHP patients, mice with maternal inheritance of exon 1 mutations that decrease $G\alpha$ and cAMP levels exhibit resistance to PTH and thyroid stimulating hormone.^{46,47} Turan et al found that paternal silencing of $G\alpha$ in renal proximal

tubules does not occur until after the early postnatal period, which could explain the development of PTH resistance and hypocalcemia only after infancy.⁴⁸ Maternal allele *GNAS* inactivation is also expected to affect NESP55 expression. Plagge et al generated mice that are deficient in *Nesp*, which showed enhanced reactivity to novel environments appropriate for the protein's predominant expression in the central nervous system.⁴⁹ XL α s, which shares sequence and functional similarity with $G\alpha$ at a protein level, forms heterotrimers with $G\beta\gamma$ subunits and activates adenylyl cyclase in specific cell types, similar to $G\alpha$.^{3,40} In addition, based on phenotypic observations from mouse models and data from newborn children, XL α s has been known to play important functions during perinatal and early postnatal development.⁵⁰ A summary of other mouse models that resemble human conditions of *GNAS* mutations has been provided elsewhere.⁸

GNAS regulation of cellular differentiation via cAMP/protein kinase A (PKA) activation

$G\alpha$ activation, through coupling to various receptors and ligands (PTH, adrenaline, glucagon, adenosine, etc) governs multiple important cellular processes to maintain physiologic functions and development in a variety of tissues. $G\alpha$ has been implicated in stem cell renewal⁵¹ and differentiation pathways, including osteogenesis,^{52,53} myogenesis,⁵⁴ adipogenesis,^{55,56} chondrogenesis,⁵⁷ and neurogenesis.⁵⁸ In the context of POH, in which ossifications occur within subcutaneous fat, a role of $G\alpha$ may be to maintain the balance in adipogenesis/osteogenesis in the mesenchymal stem cell (MSC) lineage. Although the molecular pathology of POH remains incompletely understood, substantial evidence shows that paternally inherited loss of $G\alpha$ function leads to subcutaneous HO and significant leaner phenotype in mice.^{55,59} Consistent with these observations, in vitro assays have shown that a paternally inherited $G\alpha$ -inactivating mutation impairs adipogenesis and enhances osteogenesis in MSCs.^{52,53,55}

In addition to the PTH receptor, other G protein-coupled receptors play roles in regulating the adipogenesis/osteogenesis balance in MSCs, including the adenosine, β -adrenergic, and purinergic receptors. Studies using mouse preosteoblasts and rat bone marrow MSCs demonstrate that adenosine A_{2A} receptors support adipogenic differentiation, whereas A_1 receptors play a role in the lipogenic activity of adipocytes. Conversely, A_{2B} receptors inhibit adipogenesis and activate osteogenesis, illustrating the differential effects of adenosine receptors on MSC differentiation.^{60,61} The roles of beta-adrenergic receptors (β -AR) in lipolysis and

thermogenesis are well documented in vitro and in vivo.⁶² Moreover, studies have described β 2- and β 3-AR as playing a part in regulating adipogenesis/osteogenesis partly via the cAMP/PKA pathway. Results from these studies indicate that antagonists of adrenergic receptors induce both adipogenesis and osteogenesis of mouse bone marrow MSCs. Similar to adenosine receptors, mRNA and protein expression of both β 2- and β 3-AR are elevated during adipogenesis and osteogenesis in bone marrow-derived MSCs. Interestingly, β 3-AR and β 2-AR were found to be the dominant receptors in adipogenesis and osteogenesis, respectively.^{63,64} The purinergic receptor P2Y family also stimulates adenylyl cyclase activation and cAMP production⁶⁵ and, additionally, have been implicated in white adipocyte physiology, including leptin secretion and lipolysis.⁶⁶

Signal transduction pathways downstream of *GNAS* and cAMP/PKA

In POH, HO initiates within subcutaneous fat before progressing to deep tissue, suggesting abnormal osteogenesis of mesenchymal precursor in adipose tissues. In fact, paternal inactivation of *Gs α* in adipose stromal cells (ASCs) enhances osteogenesis in vitro and promotes intramembranous HO in subcutaneous fat in vivo.⁵² Conversely, paternal-inactivation of *Gs α* in ASCs severely limits adipogenesis in vitro.⁵⁵ Importantly, this defect can be rescued by an adenylyl cyclase activator (forskolin),⁵⁵ indicating that *Gs α* -cAMP signals regulate the bipotential adipogenic-osteogenic lineage cell fate switch.

Gs α expression is tightly regulated under physiologic and developmental conditions. Although *Gs α* signaling is ubiquitous, various cell types are expected to respond to G-protein signaling and cAMP in cell and developmentally specific manners. *Gs α* appears to have crucial roles in maintaining balance in two key signaling pathways: Wnt/ β -catenin and Hedgehog (Hh).⁶⁷ Gain-of-function mutation of *Gs α* leads to overactive Wnt/ β -catenin signal and is associated with fibrous dysplasia,⁶⁸ whereas loss-of-function mutations of *Gs α* lead to increased Hh signaling and are associated with POH.⁶⁷

Hh signaling controls numerous aspects of development, including proliferation, patterning, and morphogenesis. Three Hh proteins have been identified in vertebrates: Sonic (SHH), Indian, and Desert. Desert Hh expression is limited to the male reproductive tract, Indian Hh regulates chondrogenesis and endochondral bone formation, and SHH plays a critical role in the formation of the skeleton and in cell differentiation.⁶⁹ SHH signaling was also found to have antiadipogenic and pro-osteogenic effects in mouse ASCs

(mASCs).⁷⁰ Hh signaling is activated when Hh protein binds to its receptor to relieve the protein smoothened (Smo) from its repressive state. Active Smo triggers a cascade of events that activates GLI transcription factors. Interestingly, SHH signaling enhances bone morphogenetic protein (BMP) 2 signaling (a known cytokine to be critical for osteogenesis) in mASCs. It is interesting to speculate that in POH, loss of function of *Gs α* leading to enhanced Hh signaling would increase BMP signaling and contribute to HO.

PKA (activated by cAMP) is known to negatively regulate Hh signaling by inhibiting GLI nuclear localization⁷¹ and targeting GLI for proteosomal degradation/truncation.^{72–74} Homozygous inactivation of *GNAS* in MSCs leads to overactivation of Hh signaling (observed in both in vitro and in vivo studies) and causes POH-like HO in mice.⁶⁷ Thus, elevation in Hh signaling in *Gs α* -deficient cells appears to be the upstream signal that contributes to HO in POH.

The BMP signal transduction pathway, a key regulator of osteoblast differentiation, serves to phosphorylate SMAD proteins and transcriptionally activate osteogenic genes.^{75,76} The relationship between *GNAS* and BMP signaling is not clear. However, using forskolin to stimulate cAMP production in mouse embryonic stem cells at the earliest stages of osteogenesis, Zhang et al demonstrated that BMP signaling and osteogenic markers (*Mx2*, *Osterix*) are significantly reduced, whereas adipogenic markers (*LPL*, *PPARG*, *EBP1*, *aP2*) are elevated.⁷⁷ These findings provide further evidence that cAMP serves as a key regulator of osteoblast and adipocyte lineage commitment upstream of BMP signaling.⁷⁷

GNAS mutation in POH patients

Heterozygous inactivating *GNAS* mutations occur in most patients with a definitive clinical diagnosis of POH (see following). All the POH-associated *GNAS* mutations identified in our cohort have mutations that cause a shift in the protein-coding reading frame: small deletions, insertions, duplications, or alteration of conserved splice site donor/acceptor dinucleotides (Table 1). Nonsense mutations leading to early protein termination have not been identified in this cohort. Although some patients diagnosed with AHO/PHP1a have mutations in exon 1,⁷⁸ as do patients diagnosed with PHP1a/POH, we have not identified exon 1 mutations in patients with a confirmed diagnosis of POH. Two POH patients with an exon 1 mutation have been reported;⁷⁹ however, the clinical descriptions are ambiguous, and thus these diagnoses are not confirmed (see following for further discussion). Any functional significance for the absence of exon 1 mutations is unclear and could reflect the small sample

Table 1 *GNAS* mutations in progressive osseous heteroplasia patients

GNAS location	Mutation site (cDNA)	Frameshift start (codon)
Exon 5	344–345insT	115
Exon 5	348delC;	116
Exon 5	348–349insC	117
Exon 5	355delC	119
Exon 7	565–68delGACT	189
Exon 9	679–80insC	227
Exon 10	725delC	242
Exon 10	835–39duplAACAG	280
Exon 11	860–61delTG	287
Exon 11	960insCT	321
Intron 12	IVS12+1G>C splice donor site	347
Intron 12	IVS12-1G>C splice acceptor site	347
Exon 13	1053–77dupl25n	360
Exon 13	1107–08delTG	369–370
Exon 13	1162delC	388

size of POH patients. *GNAS* mutations in POH patients reduce Gs α protein levels and decrease cAMP signaling (our unpublished data).

Our unpublished data and other reports⁸⁰ support that POH is preferentially caused by *GNAS* mutations occurring on the paternally inherited copy of the gene. This suggests that the paternal and maternal *GNAS* alleles function differently to regulate osteoblast differentiation (see earlier sections: *GNAS* regulation of cellular differentiation via cAMP/protein kinase A (PKA) activation and Signal transduction pathways downstream of *GNAS* and cAMP/PKA).

Classic mosaicism, or the presence of at least two genotypically different cell populations derived from a single zygote, may explain the pattern of lesion distribution in POH that distinguishes it from other *GNAS*-based conditions in which HO does not progress to deeper tissues. For example, a germline mutation in *GNAS* may be followed by a second mutation in the other allele during development and both be retained by a finite number of progenitor cells in the postnatal state. Possible mechanisms include the presence of somatic mutations or random inactivation of the second *GNAS* allele in progenitor cells, a de novo mutation in a gene that normally functions in a *GNAS*-interacting pathway, or epigenetic changes in somatic cells. Depending on the location of resident progenitor cells or their predisposition toward certain migration patterns, HO formation with progression to deeper tissues may be favored in POH. Revertant mosaicism in uninvolved dermomyotomes, or in patients with *GNAS* mutations and no apparent or very limited disease, cannot be excluded.⁸¹

Most POH mutations appear to be de novo mutations in a family, with a given specific mutation found in only a

single family.⁶ However, a four-nucleotide deletion in exon 7 that was found to be a mutational hot spot for AHO/PHP1a is found as a recurring mutation in POH as well. Within a family, carriers with the mutation but no clinical manifestations have been identified.

Clinical features and diagnosis

As a disorder of extraskeletal bone formation, POH must first be differentiated from nonhereditary as well as other genetic conditions of HO to diagnose the condition (Table 2).

Table 2 Differential diagnosis of extraskeletal bone formation

Genetic
Primary osteoma cutis
Fibrodysplasia ossificans progressiva
Progressive osseous heteroplasia
Albright hereditary osteodystrophy
Pseudohypoparathyroidism 1a/1c (PHP1a/1c)
Progressive osseous heteroplasia overlap syndromes (POH/PHP1a/1c)
Nonhereditary
Injury
Traumatic head injury
Paraplegia/quadruplegia (spinal cord injury)
Poliomyelitis
Guillain-Barré syndrome
Muscle hematoma
Joint dislocation
After hip and knee arthroplasty
Surgical scars
Severe burns
Secondary osteoma cutis
Endovascular injury (atherosclerosis, valvular heart disease, cerebrovascular accident)
Arthropathy
Ankylosing spondylitis
Psoriatic arthritis
Seronegative arthropathies
Diffuse idiopathic skeletal hyperostosis
Aging
Postarthroplasty
Atherosclerosis
Cerebrovascular accident
Valvular heart disease
Atherosclerosis
Miliary osteoma (of the face)*
Pressure ulcers ^{†‡}
Urinary tract infections [‡]
Other
Metastatic osteosarcoma
Fibrosing lung disorders
Pulmonary venous hypertension
Conditions that increase calcium–phosphate product levels

Notes: *Occurs predominantly in middle-aged and older females. [†]At the site of reactive soft tissue ossification. [‡]May be a predisposing factor or a secondary complication of nonhereditary heterotopic ossification.

Abbreviations: AHO, Albright hereditary osteo dystrophy; POH, progressive osseous heteroplasia; PHP, Pseudohypoparathyroidism.

Nonhereditary forms of HO are excluded on the basis of prior trauma or surgery, age, and known history or suspicion of arthropathy. POH is distinguished from fibrodysplasia ossificans progressiva (FOP), another rare autosomal dominant genetic condition of HO, by the presence of cutaneous ossification, the absence of congenital malformation of the first toes, and the absence of preosseous tumor-like inflammation or “flare-ups.”^{82,83} Other genetic causes of HO (Table 2) can be excluded on clinical grounds alone.

POH is among several related genetic disorders, including AHO, PHP, and OC, which share the common features of superficial ossification and association with inactivating mutations of *GNAS*.^{4–6} AHO is characterized by variable subsets of features, in addition to superficial HO, including short adult stature, obesity, round faces, brachydactyly, and neurobehavioral problems (including mental retardation). PHP, or end-organ resistance to PTH, is subdivided into types 1a, 1b, and 1c.⁴ Clinically, PHP1a and 1c are identical and can include AHO features, deficient responses to PTH, and multiple other hormone resistance. PHP1a is distinguished from PHP1c by the presence of inactivating *GNAS* mutations and/or reduced activity of Gs α , the major protein product encoded by the *GNAS* locus. Patients with PHP1b have hormone resistance, usually limited to PTH target tissues, but no AHO features or reduced Gs α activity. PHP1b is associated with a *GNAS* imprinting defect and caused by heterozygous deletions of a suspected imprinting control element in familial forms.^{4,14,45,84,85} Pseudopseudo-hypoparathyroidism (PPHP) refers to the condition in patients with AHO who do not have PTH resistance. OC describes superficial HO without any hormone resistance or AHO features. A familial form of primary OC has been described.⁸⁶

POH is diagnosed on the basis of three major criteria: superficial HO that progresses to deep connective tissue; two or fewer AHO features, excluding HO; and no PTH resistance (Table 3). Dermal involvement appears as hard maculopapular lesions (Figure 2A and B). Over time, these lesions coalesce into plaques with spread into deeper connective tissues including fascia, skeletal muscle, tendon, and ligament (Figure 2C). Small spicules of dermal bone may occasionally extrude through the epidermis, although bone formation does not originate in the epidermis. Extensive ossification of the deep connective tissues can result in ankylosis of affected joints and growth retardation of involved limbs (Figure 2C).^{1,87–90} In addition to HO, some patients exhibit one or two AHO features, but never obesity or multiple AHO features.⁶ Hormonal abnormalities are rarely associated with POH, and never PTH resistance.⁶

Table 3 Diagnostic criteria for progressive osseous heteroplasia

Criteria
Major criteria
Superficial and deep heterotopic ossification
Two or fewer features of Albright hereditary dystrophy, not including heterotopic ossification
No parathyroid hormone resistance
Supporting clinical findings
<i>GNAS</i> mutation
Evidence for paternal inheritance
Radiographic evidence for reticular pattern of ossification
Exclusive intramembraneous ossification or both intramembraneous and endochondral ossification
Lateralization in a dermatomyotomal pattern
History of intrauterine growth retardation
Leanness
Age of onset younger than 1 year

In addition to these key diagnostic criteria, there are several clinical findings that support the diagnosis of POH (Table 3).⁶ Although some individuals can present with a later age of onset, most POH patients have an average age of onset earlier than 1 year. Almost two-thirds of POH patients have mutations in *GNAS*. However, those without detectable mutations are clinically indistinguishable from those with mutations. Maternally inherited mutations in *GNAS* cause PHP1a, whereas paternally inherited mutations are associated with POH and are supportive of the diagnosis, especially with delayed onset of more extensive ossification. Although maternally inherited mutations are more often found with AHO, paternally inherited mutations can also be associated with AHO and lead to PPHP. POH-like HO associated with the overlap syndromes can be inherited through the maternal as well as the paternal allele. Although exon 1 mutations have been reported in individuals with subcutaneous ossifications (including those close to muscle),⁷⁹ it is unclear whether these individuals meet the POH criterion of deep HO. In a large series of POH patients, diagnosed on the basis of key criteria described here only, exon 1 mutations were not found.⁶

Birth weight tends to be very low in patients with POH, usually at or below the fifth percentile compared with sex-matched normative data.⁵⁵ In fact, heterozygous *GNAS* mutations on either parental allele were found to be associated with intrauterine growth retardation, and when these mutations were located on the paternal *GNAS* allele, intrauterine growth retardation was considerably more pronounced compared with mutations on the maternal allele.⁵⁰ At any age, POH patients with paternally inherited inactivating *GNAS* mutations were always found to have a lean phenotype.^{6,91} There is also a striking lateralization of lesions in a dermatomyotomal

Figure 2 Appearance of heterotopic ossification in POH.

Notes: (A and B) Early clinical appearance of heterotopic ossification in POH. Note the maculopapular lesions that correspond to extensive dermal and subcutaneous ossification. (C) Advanced heterotopic ossification in POH results in large coalesced bony plaques. Note the distinct lateralization of lesions that is observed in some patients, as well as contracture and ankyloses of the left lower extremity.

Abbreviation: POH, progressive osseous heteroplasia.

distribution (Figure 2C),⁸¹ but this may be difficult to assess early in the presentation. Radiographic appearance of HO in POH shows a distinctive reticular pattern of web-like ectopic bone involving soft connective tissues from the dermis down to skeletal muscle (Figure 3). Half of lesional biopsies taken in patients ultimately confirmed to have POH demonstrated intramembraneous ossification (Figure 4), with 20% showing endochondral ossification and 30% both types.⁶

POH occasionally presents as an overlap syndrome with additional features associated with other *GNAS*-based disorders of HO. Two cases in which patients exhibited progressive HO together with characteristics of AHO (short stature, round face, and brachydactyly) and reduced levels of Gs α protein were reported by Eddy et al;⁹² one of the two cases had a heterozygous *GNAS* mutation. Another patient with progressive HO had severe plate-like OC and also possessed a mutation in the *GNAS* gene.^{93,94} These cases are consistent with POH being part of a clinical spectrum of HO disorders caused by inactivating *GNAS* mutations.

POH, other disorders associated with inactivating mutations of *GNAS*, and POH overlap syndromes are distinguished solely by clinical criteria (Figure 5). *GNAS*-based disorders of HO can be divided into those presenting with stable superficial bony lesions and those in which superficial lesions progress into deep connective tissue. Among the nonprogressive forms are OC, AHO/PPHP, and PHP1a/c.

Those without AHO features have OC. Those with AHO features and no hormone resistance have AHO/PPHP, and those with hormone resistance have PHP (Figure 5). The progressive types are POH and the POH-related syndromes. Patients with POH present with superficial HO that progresses to deeper tissues in the absence of multiple other AHO features and without hormone resistance (Figure 5). A small subset of patients has progressive HO with more extensive AHO features (POH/AHO) or with both AHO features and hormone resistance (POH/PHP1a/1c). It is possible that individuals without progressive HO could be too young at the time of initial diagnosis to have yet developed progressive disease. Similarly, individuals with POH could be too young at the time of diagnosis to have yet developed other features of AHO. Nevertheless, POH and progressive HO syndromes can be distinguished from other *GNAS*-based disorders by one clinical parameter alone: the extension of HO from superficial to deep tissue. *GNAS* inactivating mutations, either by presence alone or by mutation pattern within *GNAS*, do not predict a specific disorder, variability of phenotype, or severity of progression within this spectrum.

Current management and prognosis

In POH, the degree of morbidity depends on the location and extent of HO, and in some cases, the condition

Figure 3 Radiographic appearance of severe heterotopic ossification in POH.
Notes: (A) Anterior view roentgenogram of the lower extremities of a child with POH shows progressive heterotopic ossification on the left side. Note the web-like pattern of ossification. (B) Cross-sectional, midfemur view by computerized tomography demonstrates extensive ossification of soft tissues in the superficial and deep posterior compartments of the leg.
Abbreviation: POH, progressive osseous heteroplasia.

results in severe disability.^{1,93–95} Growth retardation may be associated with limited movement of extremities caused by joint ankyloses and bone pain, and secondary osteoporosis may ensue.^{1,93–95}

Because of the ultrarare nature of POH, we have limited information about prognosis. There are no distinguishing forms of POH based on progression; however, we did observe that HO in the dermis shows a seemingly random distribution

Figure 4 Photomicrograph of a POH lesional section shows intramembranous ossification.
Notes: Note that deposits of bone are surrounded by adipose tissue (inset). Hematoxylin and eosin staining; original magnification, 200×.
Abbreviation: A, adipose tissue; B, bone; POH, progressive osseous heteroplasia.

of affected areas and that this mosaic distribution of lesions lateralized in a distinct dermomyotomal pattern present in very few conditions.⁸¹ In some patients, dermomyotomal distribution was partial, which may suggest that lesion progression was incomplete or delayed at the time of presentation. Often it is only later in the course of the condition that one can clinically determine areas of severe involvement.

At this time, there are no effective treatments or prevention for POH. Surgical resection of diffuse lesions usually leads to recurrences or complications;^{1,2,82,87} however, areas of well-circumscribed HO can often be removed, with successful long-term results.⁹⁴ Successful functional repositioning of a joint after the development of a contracture from HO was reported in the case of one child.⁸⁸ Unfortunately, amputations are sometimes needed in the setting of severe growth retardation and functional ankylosis.⁸⁹

A single case report on the use of the bisphosphonate pamidronate in POH suggested stabilization of the

Figure 5 Diagnostic algorithm for distinguishing among *GNAS*-based conditions of heterotopic ossification.
Abbreviations: HO, heterotopic ossification; OC, osteoma cutis; POH, progressive osseous heteroplasia; AHO, Albright hereditary dystrophy; HR, hormone resistance; PPHP, pseudopseudo-hypoparathyroidism; PHP, pseudohypoparathyroidism.

condition,⁹⁶ but it is unclear how generally applicable this finding may be to prevention of new skin lesions. Treatment with bisphosphonate is unlikely to resolve preexisting bone formation in POH.

Physical therapy and meticulous skin care are important conservative approaches to preserving movement and preventing cutaneous breakdown, respectively.^{90,95}

Emerging therapeutic strategies

Regard et al showed that Gs α restricts bone formation to the normotopic skeleton by inhibiting Hh signaling in mesenchymal progenitor cells, whereas genetically mediated exogenous Hh signaling is sufficient to induce POH-like HO.⁶⁷ Furthermore, inhibition of this signaling pathway by genetic or pharmacological methods reduced the severity of ectopic bone formation.⁶⁷ Therefore, Hh inhibitors currently used for other conditions, such as cancer, may be potential candidate drugs for treating HO caused by *GNAS* inactivation.⁹⁷

Endochondral ossification is present singly or in combination with intramembranous ossification in 50% of POH lesional biopsies,⁶ and so known inhibitors of endochondral ossification are potential therapies. For example, retinoic acid receptor γ agonists were shown to be highly effective at inhibiting HO in mouse models.⁹⁸ In fact, a marked increase in Gs α expression at the transcriptional level is induced by retinoic acid, suggesting that the same retinoic acid receptor γ agonists may be used to increase production of Gs α protein from the normal allele and minimize the effects of *GNAS* inactivation.⁹⁹

Acknowledgments

We acknowledge Meiqi Xu for her excellent work in conducting the mutational sequence analyses presented in this article. This work was supported by the National Institutes of Health (R01-AR046831), the Progressive Osseous Heteroplasia Association, the Italian POH Association, the International Fibrodysplasia Ossificans Progressiva Association, the University of Pennsylvania Center for Research in FOP and Related Disorders, the Isaac and Rose Nassau Professorship of Orthopaedic Molecular Medicine (FSK), and the Cali/Weldon Professorship (EMS).

Disclosure

The authors report no conflicts of interest in this work.

References

- Kaplan FS, Craver R, MacEwen GD, et al. Progressive osseous heteroplasia: a distinct developmental disorder of heterotopic ossification. Two new case reports and follow-up of three previously reported cases. *J Bone Joint Surg Am.* 1994;76(3):425–436.
- Shore EM, Ahn J, Jan de Beur S, et al. Paternally inherited inactivating mutations of the *GNAS1* gene in progressive osseous heteroplasia. *N Engl J Med.* 2002;346(2):99–106.
- Bastepe M. Relative functions of G α s and its extra-large variant XL α s in the endocrine system. *Horm Metab Res.* 2012;44(10):732–740.
- Bastepe M, Jüppner H. *GNAS* locus and pseudohypoparathyroidism. *Horm Res.* 2005;63(2):65–74.
- Weinstein LS, Liu J, Sakamoto A, Xie T, Chen M. Minireview: *GNAS*: normal and abnormal functions. *Endocrinology.* 2004;145(12):5459–5464.
- Adegbite NS, Xu M, Kaplan FS, Shore EM, Pignolo RJ. Diagnostic and mutational spectrum of progressive osseous heteroplasia (POH) and other forms of *GNAS*-based heterotopic ossification. *Am J Med Genet A.* 2008;146A(14):1788–1796.
- Thiele S, Werner R, Ahrens W, et al. Selective deficiency of Gs α and the possible role of alternative gene products of *GNAS* in Albright hereditary osteodystrophy and pseudohypoparathyroidism type Ia. *Exp Clin Endocrinol Diabetes.* 2010;118(2):127–132.
- Turan S, Bastepe M. The *GNAS* complex locus and human diseases associated with loss-of-function mutations or epimutations within this imprinted gene. *Horm Res Paediatr.* 2013;80(4):229–241.
- Ahmed SF, Dixon PH, Bonthron DT, et al. *GNAS1* mutational analysis in pseudohypoparathyroidism. *Clin Endocrinol (Oxf).* 1998;49(4):525–531.
- Ahrens W, Hiort O, Staedt P, Kirschner T, Marschke C, Kruse K. Analysis of the *GNAS1* gene in Albright's hereditary osteodystrophy. *J Clin Endocrinol Metab.* 2001;86(10):4630–4634.
- Aldred MA, Trembath RC. Activating and inactivating mutations in the human *GNAS1* gene. *Hum Mutat.* 2000;16(3):183–189.
- Farfel Z, Iiri T, Shapira H, Roitman A, Mouallem M, Bourne HR. Pseudohypoparathyroidism, a novel mutation in the betagamma-contact region of Gs α impairs receptor stimulation. *J Biol Chem.* 1996;271(33):19653–19655.
- Fischer JA, Egert F, Werder E, Born W. An inherited mutation associated with functional deficiency of the alpha-subunit of the guanine nucleotide-binding protein Gs in pseudo- and pseudopseudohypoparathyroidism. *J Clin Endocrinol Metab.* 1998;83(3):935–938.
- Jan De Beur SM, O'Connell JR, Peila R, et al. The pseudohypoparathyroidism type Ib locus is linked to a region including *GNAS1* at 20q13.3. *J Bone Miner Res.* 2003;18(3):424–433.
- Linglart A, Carel JC, Garabédian M, Lé T, Mallet E, Kottler ML. *GNAS1* lesions in pseudohypoparathyroidism Ia and Ic: genotype phenotype relationship and evidence of the maternal transmission of the hormonal resistance. *J Clin Endocrinol Metab.* 2002;87(1):189–197.
- Luttikhuis ME, Wilson LC, Leonard JV, Trembath RC. Characterization of a de novo 43-bp deletion of the Gs alpha gene (*GNAS1*) in Albright hereditary osteodystrophy. *Genomics.* 1994;21(2):455–457.
- Miric A, Vechio JD, Levine MA. Heterogeneous mutations in the gene encoding the alpha-subunit of the stimulatory G protein of adenylate cyclase in Albright hereditary osteodystrophy. *J Clin Endocrinol Metab.* 1993;76(6):1560–1568.
- Nakamoto JM, Sandstrom AT, Brickman AS, Christenson RA, Van Dop C. Pseudohypoparathyroidism type Ia from maternal but not paternal transmission of a Gs α gene mutation. *Am J Med Genet.* 1998;77(4):261–267.
- Nakamoto JM, Zimmerman D, Jones EA, et al. Concurrent hormone resistance (pseudohypoparathyroidism type Ia) and hormone independence (testotoxicosis) caused by a unique mutation in the G alpha s gene. *Biochem Mol Med.* 1996;58(1):18–24.
- Patten JL, Johns DR, Valle D, et al. Mutation in the gene encoding the stimulatory G protein of adenylate cyclase in Albright's hereditary osteodystrophy. *N Engl J Med.* 1990;322(20):1412–1419.
- Schwindinger WF, Miric A, Zimmerman D, Levine MA. A novel Gs alpha mutant in a patient with Albright hereditary osteodystrophy uncouples cell surface receptors from adenylate cyclase. *J Biol Chem.* 1994;269(41):25387–25391.

22. Shapira H, Mouallem M, Shapiro MS, Weisman Y, Farfel Z. Pseudohypoparathyroidism type 1a: two new heterozygous frameshift mutations in exons 5 and 10 of the Gs alpha gene. *Hum Genet.* 1996; 97(1):73–75.
23. Walden U, Weissörtel R, Corria Z, et al. Stimulatory guanine nucleotide binding protein subunit 1 mutation in two siblings with pseudohypoparathyroidism type 1a and mother with pseudopseudohypoparathyroidism. *Eur J Pediatr.* 1999;158(3):200–203.
24. Warner DR, Gejman PV, Collins RM, Weinstein LS. A novel mutation adjacent to the switch III domain of G(s)alpha in a patient with pseudohypoparathyroidism. *Mol Endocrinol.* 1997;11(11):1718–1727.
25. Warner DR, Weng G, Yu S, Matalon R, Weinstein LS. A novel mutation in the switch 3 region of G(s)alpha in a patient with Albright hereditary osteodystrophy impairs GDP binding and receptor activation. *J Biol Chem.* 1998;273(37):23976–23983.
26. Weinstein LS, Gejman PV, de Mazancourt P, American N, Spiegel AM. A heterozygous 4-bp deletion mutation in the Gs alpha gene (GNAS1) in a patient with Albright hereditary osteodystrophy. *Genomics.* 1992;13(4):1319–1321.
27. Weinstein LS, Gejman PV, Friedman E, et al. Mutations of the Gs alpha-subunit gene in Albright hereditary osteodystrophy detected by denaturing gradient gel electrophoresis. *Proc Natl Acad Sci U S A.* 1990;87(21):8287–8290.
28. Wilson LC, Oude Luttikhuis ME, Clayton PT, Fraser WD, Trembath RC. Parental origin of Gs alpha gene mutations in Albright's hereditary osteodystrophy. *J Med Genet.* 1994;31(11):835–839.
29. Yokoyama M, Takeda K, Iyota K, Okabayashi T, Hashimoto K. A 4-base pair deletion mutation of Gs alpha gene in a Japanese patient with pseudohypoparathyroidism. *J Endocrinol Invest.* 1996;19(4):236–241.
30. Yu D, Yu S, Schuster V, Kruse K, Clericuzio CL, Weinstein LS. Identification of two novel deletion mutations within the Gs alpha gene (GNAS1) in Albright hereditary osteodystrophy. *J Clin Endocrinol Metab.* 1999;84(9):3254–3259.
31. Yu S, Yu D, Hainline BE, et al. A deletion hot-spot in exon 7 of the Gs alpha gene (GNAS1) in patients with Albright hereditary osteodystrophy. *Hum Mol Genet.* 1995;4(10):2001–2002.
32. Baltoumas FA, Theodoropoulou MC, Hamodrakas SJ. Interactions of the α -subunits of heterotrimeric G-proteins with GPCRs, effectors and RGS proteins: a critical review and analysis of interacting surfaces, conformational shifts, structural diversity and electrostatic potentials. *J Struct Biol.* 2013;182(3):209–218.
33. Oldham WM, Hamm HE. Heterotrimeric G protein activation by G-protein-coupled receptors. *Nat Rev Mol Cell Biol.* 2008;9(1):60–71.
34. Wettschreck N, Offermanns S. Mammalian G proteins and their cell type specific functions. *Physiol Rev.* 2005;85(4):1159–1204.
35. Heydorn A, Ward RJ, Jorgensen R, et al. Identification of a novel site within G protein alpha subunits important for specificity of receptor-G protein interaction. *Mol Pharmacol.* 2004;66(2):250–259.
36. McCudden CR, Hains MD, Kimple RJ, Siderovski DP, Willard FS. G-protein signaling: back to the future. *Cell Mol Life Sci.* 2005;62(5):551–577.
37. Huang C, Hepler JR, Gilman AG, Mumby SM. Attenuation of Gi- and Gq-mediated signaling by expression of RGS4 or GAIP in mammalian cells. *Proc Natl Acad Sci U S A.* 1997;94(12):6159–6163.
38. Roy AA, Lemberg KE, Chidiac P. Recruitment of RGS2 and RGS4 to the plasma membrane by G proteins and receptors reflects functional interactions. *Mol Pharmacol.* 2003;64(3):587–593.
39. Zheng B, Ma YC, Ostrom RS, et al. RGS-PX1, a GAP for G(s)alpha and sorting nexin in vesicular trafficking. *Science.* 2001;294(5548):1939–1942.
40. Bastepe M. The GNAS Locus: Quintessential Complex Gene Encoding G(s)alpha, XLalphas, and other Imprinted Transcripts. *Curr Genomics.* 2007;8(6):398–414.
41. Plagge A, Kelsey G, Germain-Lee EL. Physiological functions of the imprinted Gnas locus and its protein variants G(s)alpha(s) and XLalpha(s) in human and mouse. *J Endocrinol.* 2008;196(2):193–214.
42. Ishikawa Y, Bianchi C, Nadal-Ginard B, Homcy CJ. Alternative promoter and 5' exon generate a novel Gs alpha mRNA. *J Biol Chem.* 1990;265(15):8458–8462.
43. Puzhko S, Goodyer CG, Kerachian MA, et al. Parathyroid hormone signaling via G(s) is selectively inhibited by an NH(2)-terminally truncated G(s): implications for pseudohypoparathyroidism. *J Bone Miner Res.* 2011;26(10):2473–2485.
44. Mantovani G, Bondioni S, Locatelli M, et al. Biallelic expression of the G(s)alpha gene in human bone and adipose tissue. *J Clin Endocrinol Metab.* 2004;89(12):6316–6319.
45. Weinstein LS, Yu S, Warner DR, Liu J. Endocrine manifestations of stimulatory G protein alpha-subunit mutations and the role of genomic imprinting. *Endocr Rev.* 2001;22(5):675–705.
46. Chen M, Gavrilova O, Liu J, et al. Alternative Gnas gene products have opposite effects on glucose and lipid metabolism. *Proc Natl Acad Sci U S A.* 2005;102(20):7386–7391.
47. Germain-Lee EL, Schwindinger W, Crane JL, et al. A mouse model of albright hereditary osteodystrophy generated by targeted disruption of exon 1 of the Gnas gene. *Endocrinology.* 2005;146(11):4697–4709.
48. Turan S, Fernandez-Rebollo E, Aydin C, et al. Postnatal establishment of allelic G(s) silencing as a plausible explanation for delayed onset of parathyroid hormone resistance owing to heterozygous G(s) disruption. *J Bone Miner Res.* 2014;29(3):749–760.
49. Plagge A, Isles AR, Gordon E, et al. Imprinted Nesp55 influences behavioral reactivity to novel environments. *Mol Cell Biol.* 2005;25(8):3019–3026.
50. Richard N, Molin A, Coudray N, Rault-Guillaume P, Jüppner H, Kottler ML. Paternal GNAS mutations lead to severe intrauterine growth retardation (IUGR) and provide evidence for a role of XL(s) in fetal development. *J Clin Endocrinol Metab.* 2013;98(9):E1549–E1556.
51. Kobayashi NR, Hawes SM, Crook JM, Pébay A. G-protein coupled receptors in stem cell self-renewal and differentiation. *Stem Cell Rev.* 2010;6(3):351–366.
52. Pignolo RJ, Xu M, Russell E, et al. Heterozygous inactivation of Gnas in adipose-derived mesenchymal progenitor cells enhances osteoblast differentiation and promotes heterotopic ossification. *J Bone Miner Res.* 2011;26(11):2647–2655.
53. Wu JY, Aarnisalo P, Bastepe M, et al. G(s)alpha enhances commitment of mesenchymal progenitors to the osteoblast lineage but restrains osteoblast differentiation in mice. *J Clin Invest.* 2011;121(9):3492–3504.
54. Berdeaux R, Stewart R. cAMP signaling in skeletal muscle adaptation: hypertrophy, metabolism, and regeneration. *Am J Physiol Endocrinol Metab.* 2012;303(1):E1–E17.
55. Liu JJ, Russell E, Zhang D, Kaplan FS, Pignolo RJ, Shore EM. Paternally inherited G(s)alpha mutation impairs adipogenesis and potentiates a lean phenotype in vivo. *Stem Cells.* 2012;30(7):1477–1485.
56. Sinha P, Aarnisalo P, Chubb R, et al. Loss of G(s)alpha early in the osteoblast lineage favors adipogenic differentiation of mesenchymal progenitors and committed osteoblast precursors. *J Bone Miner Res.* 2014;29(11):2414–2426.
57. Sakamoto A, Chen M, Kobayashi T, Kronenberg HM, Weinstein LS. Chondrocyte-specific knockout of the G protein G(s)alpha leads to epiphyseal and growth plate abnormalities and ectopic chondrocyte formation. *J Bone Miner Res.* 2005;20(4):663–671.
58. Doze VA, Perez DM. G-protein-coupled receptors in adult neurogenesis. *Pharmacol Rev.* 2012;64(3):645–675.
59. Huso DL, Edie S, Levine MA, et al. Heterotopic ossifications in a mouse model of albright hereditary osteodystrophy. *PLoS ONE.* 2011;6(6):e21755.
60. Gharibi B, Abraham AA, Ham J, Evans BA. Adenosine receptor subtype expression and activation influence the differentiation of mesenchymal stem cells to osteoblasts and adipocytes. *J Bone Miner Res.* 2011;26(9):2112–2124.
61. Gharibi B, Abraham AA, Ham J, Evans BA. Contrasting effects of A1 and A2b adenosine receptors on adipogenesis. *Int J Obes.* 2012;36(3):397–406.

62. Bachman ES, Dhillon H, Zhang CY, et al. betaAR signaling required for diet-induced thermogenesis and obesity resistance. *Science*. 2002;297(5582):843–845.
63. Li H, Fong C, Chen Y, Cai G, Yang M. Beta-adrenergic signals regulate adipogenesis of mouse mesenchymal stem cells via cAMP/PKA pathway. *Mol Cell Endocrinol*. 2010;323(2):201–207.
64. Li H, Fong C, Chen Y, Cai G, Yang M. beta2- and beta3-, but not beta1-adrenergic receptors are involved in osteogenesis of mouse mesenchymal stem cells via cAMP/PKA signaling. *Arch Biochem Biophys*. 2010;496(2):77–83.
65. Torres B, Zambon AC, Insel PA. P2Y11 receptors activate adenylyl cyclase and contribute to nucleotide-promoted cAMP formation in MDCK-D(1) cells. A mechanism for nucleotide-mediated autocrine-paracrine regulation. *J Biol Chem*. 2002;277(10):7761–7765.
66. Lee H, Jun DJ, Suh BC, et al. Dual roles of P2 purinergic receptors in insulin-stimulated leptin production and lipolysis in differentiated rat white adipocytes. *J Biol Chem*. 2005;280(31):28556–28563.
67. Regard JB, Malhotra D, Gvozdenovic-Jeremic J, et al. Activation of Hedgehog signaling by loss of GNAS causes heterotopic ossification. *Nat Med*. 2013;19(11):1505–1512.
68. Regard JB, Cherman N, Palmer D, et al. Wnt/ β -catenin signaling is differentially regulated by G α proteins and contributes to fibrous dysplasia. *Proc Natl Acad Sci U S A*. 2011;108(50):20101–20106.
69. James AW. Review of Signaling Pathways Governing MSC Osteogenic and Adipogenic Differentiation. *Scientifica (Cairo)*. 2013;2013:684736.
70. James AW, Leucht P, Levi B, et al. Sonic Hedgehog influences the balance of osteogenesis and adipogenesis in mouse adipose-derived stromal cells. *Tissue Eng Part A*. 2010;16(8):2605–2616.
71. Sheng T, Chi S, Zhang X, Xie J. Regulation of Gli1 localization by the cAMP/protein kinase A signaling axis through a site near the nuclear localization signal. *J Biol Chem*. 2006;281(1):9–12.
72. Epstein DJ, Marti E, Scott MP, McMahon AP. Antagonizing cAMP-dependent protein kinase A in the dorsal CNS activates a conserved Sonic hedgehog signaling pathway. *Development*. 1996;122(9):2885–2894.
73. Makinodan E, Marnaros AG. Protein kinase A activation inhibits oncogenic Sonic hedgehog signalling and suppresses basal cell carcinoma of the skin. *Exp Dermatol*. 2012;21(11):847–852.
74. Tuson M, He M, Anderson KV. Protein kinase A acts at the basal body of the primary cilium to prevent Gli2 activation and ventralization of the mouse neural tube. *Development*. 2011;138(22):4921–4930.
75. Lian JB, Stein GS, Javed A, et al. Networks and hubs for the transcriptional control of osteoblastogenesis. *Rev Endocr Metab Disord*. 2006;7(1–2):1–16.
76. Matsubara T, Kida K, Yamaguchi A, et al. BMP2 regulates Osterix through Msx2 and Runx2 during osteoblast differentiation. *J Biol Chem*. 2008;283(43):29119–29125.
77. Zhang S, Kaplan FS, Shore EM. Different roles of GNAS and cAMP signaling during early and late stages of osteogenic differentiation. *Horm Metab Res*. 2012;44(10):724–731.
78. Lemos MC, Thakker RV. GNAS mutations in pseudohypoparathyroidism type 1a and related disorders. *Hum Mutat*. Epub September 13, 2014.
79. Lebrun M, Richard N, Abeguilé G, et al. Progressive osseous heteroplasia: a model for the imprinting effects of GNAS inactivating mutations in humans. *J Clin Endocrinol Metab*. 2010;95(6):3028–3038.
80. Elli FM, Barbieri AM, Bordogna P, et al. Screening for GNAS genetic and epigenetic alterations in progressive osseous heteroplasia: first Italian series. *Bone*. 2013;56(2):276–280.
81. Cairns DM, Pignolo RJ, Uchimura T, et al. Somitic disruption of GNAS in chick embryos mimics progressive osseous heteroplasia. *J Clin Invest*. 2013;123(8):3624–3633.
82. Kaplan FS, Shore EM. Progressive osseous heteroplasia. *J Bone Miner Res*. 2000;15(11):2084–2094.
83. Shore EM, Xu M, Feldman GJ, et al. A recurrent mutation in the BMP type I receptor ACVR1 causes inherited and sporadic fibrodysplasia ossificans progressiva. *Nat Genet*. 2006;38(5):525–527.
84. Bastepe M, Fröhlich LF, Linglart A, et al. Deletion of the NESP55 differentially methylated region causes loss of maternal GNAS imprints and pseudohypoparathyroidism type Ib. *Nat Genet*. 2005;37(1):25–27.
85. Jüppner H, Schipani E, Bastepe M, et al. The gene responsible for pseudohypoparathyroidism type Ib is paternally imprinted and maps in four unrelated kindreds to chromosome 20q13.3. *Proc Natl Acad Sci U S A*. 1998;95(20):11798–11803.
86. Ruggieri M, Pavone V, Polizzi A, et al. Familial osteoma of the cranial vault. *Br J Radiol*. 1998;71(842):225–228.
87. Athanasou NA, Benson MK, Brenton BP, Smith R. Progressive osseous heteroplasia: a case report. *Bone*. 1994;15(5):471–475.
88. Rosenfeld SR, Kaplan FS. Progressive osseous heteroplasia in male patients. Two new case reports. *Clin Orthop Relat Res*. 1995;(317):243–245.
89. Schmidt AH, Vincent KA, Aiona MD. Hemimelic progressive osseous heteroplasia. A case report. *J Bone Joint Surg Am*. 1994;76(6):907–912.
90. Urtizberea JA, Testart H, Cartault F, Boccon-Gibod L, Le Merrer M, Kaplan FS. Progressive osseous heteroplasia. Report of a family. *J Bone Joint Surg Br*. 1998;80(5):768–771.
91. Long DN, McGuire S, Levine MA, Weinstein LS, Germain-Lee EL. Body mass index differences in pseudohypoparathyroidism type 1a versus pseudopseudohypoparathyroidism may implicate paternal imprinting of Galpha(s) in the development of human obesity. *J Clin Endocrinol Metab*. 2007;92(3):1073–1079.
92. Eddy MC, Jan De Beur SM, Yandow SM, et al. Deficiency of the alpha-subunit of the stimulatory G protein and severe extraskelatal ossification. *J Bone Miner Res*. 2000;15(11):2074–2083.
93. Tresserra L, Tresserra F, Grases PJ, Badosa J, Tresserra M. Congenital plate-like osteoma cutis of the forehead: an atypical presentation form. *J Craniomaxillofac Surg*. 1998;26(2):102–106.
94. Yeh GL, Mathur S, Wivel A, et al. GNAS1 mutation and Cbfa1 mis-expression in a child with severe congenital platelike osteoma cutis. *J Bone Miner Res*. 2000;15(11):2063–2073.
95. Aynaci O, Mijgan Aynaci F, Cobanoğlu U, Alpay K. Progressive osseous heteroplasia. A case report and review of the literature. *J Pediatr Orthop B*. 2002;11(4):339–342.
96. Hou JW. Progressive osseous heteroplasia controlled by intravenous administration of pamidronate. *Am J Med Genet A*. 2006;140(8):910–913.
97. Trinh TN, McLaughlin EA, Gordon CP, McCluskey A. Hedgehog signalling pathway inhibitors as cancer suppressing agents. *Med Chem Comm*. 2014;5(2):117–133.
98. Shimono K, Tung WE, Macolino C, et al. Potent inhibition of heterotopic ossification by nuclear retinoic acid receptor- γ agonists. *Nat Med*. 2011;17(4):454–460.
99. Chan SD, Strewler GJ, Nissenson RA. Transcriptional activation of Gs alpha expression by retinoic acid and parathyroid hormone-related protein in F9 teratocarcinoma cells. *J Biol Chem*. 1990;265(33):20081–20084.

The Application of Clinical Genetics

Dovepress

Publish your work in this journal

The Application of Clinical Genetics is an international, peer-reviewed open access journal that welcomes laboratory and clinical findings in the field of human genetics. Specific topics include: Population genetics; Functional genetics; Natural history of genetic disease; Management of genetic disease; Mechanisms of genetic disease; Counseling and ethical

issues; Animal models; Pharmacogenetics; Prenatal diagnosis; Dysmorphology. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/the-application-of-clinical-genetics-journal>