

The efficacy and safety of add-on 0.1% brimonidine tartrate preserved with sodium chlorite in on-treatment Japanese normal-tension glaucoma patients [Corrigendum]

Tsumura T, Yoshikawa K, Kimura T, et al. *Clin Ophthalmol*. 2014;8:1681–1687.

The authors would like to make corrections to Table 3, on page 1684, as some of the data were incorrect. The revised version of Table 3 is shown below.

Table 3 Time course of conjunctival hyperemia and superficial punctate keratitis (SPK) scores^a

	Baseline	4 weeks	8 weeks	12 weeks
Conjunctival hyperemia scores	0.6±0.7	0.5±0.7	0.5±0.7	0.5±0.7
P-value		0.4545	0.0768	0.3750
SPK scores	0.2±0.3	0.1±0.2	0.2±0.3	0.2±0.3
P-value		0.5309	0.4381	0.6245

Notes: ^aScores are presented as the mean ± standard deviation; analyzed by Wilcoxon signed rank test.

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on

Submit your manuscript here: <http://www.dovepress.com/clinical-ophthalmology-journal>

PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.