ORIGINAL RESEARCH

P-wave and QT dispersion in patients with conversion disorder

Therapeutics and Clinical Risk Management downloaded from https://www.dovepress.com/ For personal use only.

Filiz Izci¹ Hilal Hocagil² Servet Izci³ Vedat Izci⁴ Merve Iris Koc⁵ Rezzan Deniz Acar³

Department of Psychiatry, Istanbul Bilim University, Sısli Florence Nightingale Hospital; ²Department of Emergency, Faculty of Medicine Hospital Zonguldak Bulent Ecevit University, Zonguldak, Turkey; ³Department of Cardiology, Kosuyolu High Specialization Training and Research Hospital, Istanbul, Turkey; ⁴Department of Emergency, Kartal Training and Research Hospital, Istanbul, Turkey; ⁵Department of Psychiatry, Erenköy Training and Research Hospital for Psychiatry, Istanbul, Turkey

submit your manuscript | www.dovepress.com

Dovencess

Objective: The aim of this study was to investigate QT dispersion (QTd), which is the noninvasive marker of ventricular arrhythmia and sudden cardiac death, and P-wave dispersion, which is the noninvasive marker of atrial arrhythmia, in patients with conversion disorder (CD).

Patients and methods: A total of 60 patients with no known organic disease who were admitted to outpatient emergency clinic and were diagnosed with CD after psychiatric consultation were included in this study along with 60 healthy control subjects. Beck Anxiety Inventory and Beck Depression Scale were administered to patients and 12-lead electrocardiogram measurements were obtained. Pd and OTd were calculated by a single blinded cardiologist.

Results: There was no statistically significant difference in terms of age, sex, education level, socioeconomic status, weight, height, and body mass index between CD patients and controls. Beck Anxiety Inventory scores (25.2 ± 10.8 and 3.8 ± 3.2 , respectively, P < 0.001) and Beck Depression Scale scores (11.24 \pm 6.15 and 6.58 \pm 5.69, respectively, P<0.01) were significantly higher in CD patients. P-wave dispersion measurements did not show any significant differences between conversion patients and control group (46 ± 5.7 vs 44 ± 5.5 , respectively, P=0.156). Regarding QTc and QTd, there was a statistically significant increase in all intervals in conversion patients (416±10 vs 398±12, P<0.001, and 47±4.8 vs 20±6.1, P<0.001, respectively).

Conclusion: A similar relation to that in literature between QTd and anxiety and somatoform disorders was also observed in CD patients. QTc and QTd were significantly increased compared to the control group in patients with CD. These results suggest a possibility of increased risk of ventricular arrhythmia resulting from QTd in CD patients. Larger samples are needed to evaluate the clinical course and prognosis in terms of arrhythmia risk in CD patients. Keywords: ECG, arrhythmia, conversion disorder

Introduction

Some psychiatric disorders are known to be correlated with cardiological disorders. There are a number of studies in literature examining the relationship between anxiety disorders and P-wave dispersion (Pd),¹ which is considered as an indirect indicator of atrial fibrillation (AF), and QT dispersion (QTd),² which is considered as an indirect indicator of ventricular arrhythmia.³⁻⁵ As well as anxiety disorders, heart rhythm disturbances have been observed in patients with somatoform disorders and a correlation between heart rate and psychiatric conditions is thought to exist.^{6,7} Conversion disorder (CD) is a type of somatoform disorder according to Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR). CD is defined as loss of function or decreased or increased function in motor, sensory, and neurovegetative system organs without an organic cause.⁸ According to DSM-IV-TR, diagnosis of CD can be made only when functional impairment cannot be explained by neurological and other medical conditions or drug effects and is associated with psychological factors,

http://dx.doi.org/10.2147/TCRM.S81852

© 2015 Izci et al. This work is published by Dove Medical Press Limited, and Licensed under Creative Commons Attribution — Non Commercial (unported, v3.0) License. The full terms of the License are available at http://creativecommons.org/licenses/by-nc/3.0/. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. Permissions beyond the scope of the License are administered by Dove Medical Press Limited. Information on how to request permission may be found at: http://www.dovepress.com/permissions.php

Correspondence: Filiz Izci Department of Psychiatry, Istanbul Bilim University, Sisli Florence Nightingale Hospital, Abide-I Hurriytet Cad. No:164 Sisli, 34381 Istanbul, Turkey Tel +90 505 450 3013 Email drfiliz2008@hotmail.com

is not malingering or merely pain or sexual dysfunction, and causes significant distress and impairment of psychological and occupational functioning.⁹

In our country, 12%–47% of those admitted to psychiatric emergency clinics have CD.^{10,11} Although a common disease, it is often overlooked or incorrectly treated. Because CD can mimic many other disorders, medical errors may occur during diagnosis.^{12,13} It is difficult to diagnose CD because its symptoms may get confused with other psychiatric and somatic disorders.¹¹ Here we aimed to investigate the risk of ventricular arrhythmia and AF in patients with CD who often visit emergency rooms with intense anxiety and somatization symptoms.

Methods

A total of 60 patients admitted to general emergency outpatient clinic whose blood count, routine biochemistry, and thyroid hormone profile were at normal range were included and cardiology consultations were made with a preliminary diagnosis of CD. All examination findings were normal, no organic pathology was found, none of the patients were on medication, and the patients were diagnosed with CD by a psychiatrist using structured interview for DSM-IV psychiatric disorders (Structured Clinical Interview for DSM-IV Axis I Disorders – Clinical Version). Also 60 healthy control subjects were included, and written informed consent was taken from all subjects.

The selected population, according to inclusion criteria, included patients who were between 18 and 45 years of age, were literate, and had a diagnosis of CD according to DSM-IV-TR and agreed to participate in the study.

Exclusion criteria were having mental retardation; alcohol and substance use-related disorders; anxiety disorders; mood disorders; schizophrenia and other psychotic disorder diagnosis according to DSM-IV-TR or dementia and other cognitive disorders; neurological disorders leading to unique personality traits such as epilepsy, migraine, multiple sclerosis, and Parkinson's disease; systemic disorders leading to cognitive impairment; or general chronic medical conditions that may disrupt cardiac conduction system (especially thyroid disease, hypertension, heart valve disorders, myocardial infarction, known atherosclerotic heart disease history, congestive heart failure, cardiomyopathy, pheochromocytoma or diabetes mellitus history, etc). Also patients on continuous drugs for a chronic disease or smoking were excluded.

The control group included volunteers who agreed to participate and were physically and mentally healthy, were not using any drugs, alcohol, or psychoactive substances, and were not smoking.

Materials

A semistructured sociodemographic and clinical data form designed by us according to clinical experience and information obtained from resources considering the purpose of the study (Structured Clinical Interview for DSM-IV Axis I Disorders, Beck Anxiety Inventory, and Beck Depression Inventory) was administered. Also, 12 standard electrocardiogram (ECG) was obtained using a standardized recording device at 50 mm/s paper speed and 1 mV/cm.

Measures

Patient follow-up form (sociodemographic and clinical data collection form)

This form was completed by the researcher doctor and questions regarded patient's age, sex, weight, height, marital status, education level, employment status, income level, health insurance type, smoking, alcohol, drug use, and psychiatric and cardiac disease history of self and family were included.

Structured Clinical Interview for DSM-IV Axis I Disorders – Clinical Version

It is a structured interview made by First et al¹⁴ for DSM-IV Axis I Disorders. Validity and reliability for Turkish population were made.¹⁵

Beck Anxiety Inventory

It was developed by Beck et al in 1988 because of the need for a scale to distinguish anxiety from depression.¹⁶

It measures the severity of anxiety symptoms and questions subjective anxiety and somatic symptoms. It is a Likert-type (scored 0–3), self-rating scale consisting of 21 items, rating between 0 and 63. Higher total scores indicate the severity of the anxiety symptoms. Validity and reliability studies for Turkey were conducted by Ulusoy et al.¹⁷

Beck Depression Inventory

It is a 21-item self-report scale developed by Beck in 1961 in order to measure emotional, cognitive, somatic, and motivational components.¹⁸ The scale is composed of 21 items of which 2 of them belong to emotions, 11 of them belong to cognition, 2 of them belong to behaviors, 5 of them belong to somatic symptoms, and 1 of them belongs to interpersonal symptoms. It has a total of 21 questions. Each question scores between 0 and 3 points, and a total score ranging between 0 and 63 points is obtained. Scores between 0 and 9 indicate no/minimal, 10–18 mild, 19–29 moderate, and 30–63 severe depression. Turkish validity and reliability study of Beck Depression Inventory which was used to determine the intensity of depression was conducted by Hisli.¹⁹

Electrocardiographic evaluation

Standard ECGs were taken from all patients with a sweeping rate of 50 mm/s and amplitude of 1 mV/cm. P-wave durations and the QT intervals were measured manually in all simultaneously recorded 12 leads of the surface ECG. The mean P-wave duration for at least three complexes was calculated in each lead. The onset of the P-wave is defined as the point of first visible upward slope from baseline for positive waveforms, and as the point of first downward slope from baseline for negative waveforms. The return to the baseline was considered as the end of the P-wave. The Pmax measured in any of the 12 leads of the surface ECG was used as the longest atrial conduction time. The maximum and minimum durations of P-waves (Pmax and Pmin, respectively) were detected, and the difference between Pmax and Pmin was defined as P-wave dispersion (Pd = Pmax – Pmin).

The QT interval was measured from the beginning of the QRS complex to the end of the T-wave at the level of the TP isoelectric baseline. Biphasic T-waves were measured to the time of their final return to the TP isoelectric baseline. When U-waves were present, the QT interval was measured from the beginning of the QRS complex to the nadir of the curve between the T- and U-waves. Heart rate-corrected QT (QTc) was calculated using Bazett's formula (QTc-d, JTc-d).¹ The ventricular repolarization dispersions were determined by calculating the difference between maximum and minimum QT intervals on each ECG, which were termed QTd and heart rate-corrected QTd (QTcd). Extrasystolic and post-extrasystolic cycles were excluded from the measurement.

Statistical analyses

Statistical analysis was performed using the Statistical Package for Social Sciences version 15.0 software for Windows (IBM, Armonk, NY). Categorical data were expressed as frequencies. Continuous variables were presented as the mean \pm standard deviation. The variances between the groups were compared with paired-samples *t*-test. The correlation between QTd and P-wave dispersion was analyzed using Pearson correlation analysis. *P*-values <0.05 were regarded as significant.

Results

A total of 60 patients diagnosed with CD according to DSM-IV-TR diagnostic criteria and 60 age- and sex-matched healthy control subjects were enrolled. In the patients group, 68.3% (n=41) were male, and in the control group, 31.7%

Table	L	Sociodemographic	characteristics	of	the	CD	patients
and hea	lt	hy controls					

	CD patients (n=60) (%)	Control group (n=60) (%)	P-value
Age (mean ± SD)	35.70±12.1	32.00±9.3	0.06
Sex			
Male	41 (68.3)	40 (66.6)	0.67
Female	19 (31.7)	20 (33.3)	
Educational status			
Not educated	24 (40.0)	13 (21.0)	0.102
Primary school	22 (36.7)	24 (38.8)	
High school	12 (20.0)	23 (37.0)	
University	2 (3.3)	2 (3.2)	
Marital status			
Married	33 (55.0)	25 (42)	0.14
Single	27 (45.0)	35 (58)	
Occupational status			
Employed	10 (16.6)	10 (20.0)	0.06
Unemployed	42 (70)	30 (60.0)	
Student	8 (13.3)	10 (20.0)	
BMI (kg/m²) (mean ± SD)	23.5±8.6	24.9±9.6	0.66

Abbreviations: BMI, body mass index; CD, conversion disorder; SD, standard deviation.

(n=19) were female. The mean ages in the patient and the control group were 35.70 ± 12.1 and 32.00 ± 9.3 , respectively. There was no statistically significant difference in terms of sociodemographic characteristics between patient and control groups (P>0.05) (Table 1).

Sixty participants with conversion in sinus rhythm and 60 individuals as a control group were enrolled. The mean heart rate of the patients was 86 ± 13 per minute. P-wave dispersion measurements did not show any significant differences between the patient and the control group (46 ± 5.7 vs 44 ± 5.5 , P=0.156). As seen in Figure 1, regarding QTc and QTd there were statistically significant increases in all intervals in patients with CD (416 ± 10 vs 398 ± 12 , P<0.001 and 47 ± 4.8 vs 20 ± 6.1 , P<0.001, respectively). P-wave measurements and QT parameters are set out in Table 2. The dispersion of P-wave did not show a significant correlation with QTd (r=-0.059; P=0.655).

Discussion

CD is defined as a loss of function in sensory and motor organs without an organic cause.²⁰ CD may occur as part of a chronic pattern known as somatization disorder in the form of physical expression of psychological distress. As a reflection of all these developments, CD took place under the head-ing "somatoform disorders" in DSM-IV.²¹ This disease has different clinical manifestations, and a study regarding the frequency of these manifestations with 86 patients revealed



Figure I QTc and QTd measurements of conversion patients and controls as figure. Notes: (A) QTc (ms) measurements of the conversion patients and controls. (B) QTd (ms) measurements of conversion patienst and controls. Abbreviations: BAI, Beck Anxiety Inventory; BDI, Beck Depression Inventory; QTc, corrected QT; QTd, dispersion QT.

that 52.3% of the cases had fainting symptoms, 22.1% of the cases had short-of-breath symptoms, and 18.6% had paralysis symptoms.²² CD may be seen at any age, although it is known that it is most common between 15 and 35 years of age.²³ Usually, it is seen by 10% of the population in developing and 13% in underdeveloped countries.²⁰ In our country, 12%-47% of psychiatric emergency admissions and 25.1% of psychiatric consultations are CD.24,25 Studies showed that there are no sex differences during preadolescent period; however, after puberty it is 2-19 times more frequent in women.²⁶⁻²⁸ In our study, female and male rates in the patient group were 68.3% and 31.7%, respectively. Before somatoform disorder diagnosis is made, the physician should carefully exclude undiagnosed general medical conditions, drug-related conditions, and other psychiatric disorders that may cause similar symptoms or simulated disorders and factitious disorders in which the patient behaves as if existing symptoms were not present. However, medical disorders other than neurological or psychiatric disorders may occur later in 25%-50% of CD patients. Therefore, it is of utmost importance to make full medical and neurological evaluation of CD patients.²⁹ One of these that may occur in anxiety and somatoform disorders is heart conduction abnormalities.

Studies investigating the relationship between anxiety disorders, somatoform disorders, and P-wave and QTd which are considered as indicators of cardiac arrhythmia are very few. Some studies suggested QTd and Pd as indicators of atrial and ventricular arrhythmias in patients with social phobia and panic disorder.^{4,30} However, similar studies with CD patients were not found. Here we aimed to evaluate P-wave and QTd and identify possible cardiac conduction abnormalities in CD in which significant somatization and anxiety symptoms were seen.

We tried to measure heart rate, rhythm, and time required for electrical activity to reach to the other side of the heart by using our ECG device.³¹ P-waves indicate atrial depolarization³² and P-wave dispersion (Pd) indicates the difference between maximum and minimum P-wave duration.³³ P-wave

	-	-		
CD patients (n=60)	Control group (n=60)	P-value		
(mean \pm SD)	(mean \pm SD)			
99±7.6	101±8.0	0.051		
54±6.0	53±5.7	0.221		
46±5.7	44±5.5	0.156		
416±10	398±12	0.000*		
47±4.8	20±6.1	0.000*		
	CD patients (n=60) (mean ± SD) 99±7.6 54±6.0 46±5.7 416±10 47±4.8	CD patients (n=60) (mean ± SD) Control group (n=60) (mean ± SD) 99±7.6 101±8.0 54±6.0 53±5.7 46±5.7 44±5.5 416±10 398±12 47±4.8 20±6.1		

Table 2 P-wave dispersion (ms) and QT dispersion (ms) in CD patients and healthy controls

Note: *P<0.000.

Abbreviations: CD, conversion disorder; SD, standard deviation; ms, milliseconds; QTd, QT dispersion; Pmax, the maximum measurement of P wave; Pmin, the minimum measurements of P wave; Pd, Pmax-Pmin; QTc, corrected QT.

The measurements	CD patients (n=60)	Control group (n=60)	P-value
of scales	(mean \pm SD)	(mean \pm SD)	
BAI (mean + SD)	25.2±10.8	3.8±3.2	0.001*
BDI (mean + SD)	11.24±6.15	6.58±5.69	0.01**

Table 3 BAI and BDI scores in CD patients group and healthy group

Notes: *P<0.001, **P<0.01. According to BAI and BDI scores administered to patients and controls: BAI scores were 25.2±10.8 and 3.8±3.2 in the patient and control groups, respectively. BAI scores were statistically significantly higher in the patient group (P<0.001). BDI scores were 11.24±6.15 and 6.58±5.69 in the patient and control groups, respectively. BDI scores were also statistically significantly higher in the patient group (P<0.01).

Abbreviations: CD, conversion disorder; BAI, Beck Anxiety Index; BDI, Beck Depression Index; SD, standard deviation.

dispersion is considered as a sensitive and specific precursor of AF in various clinical situations,³⁴ and it is reported to be used as a predictor for paroxysmal AF.1 Significant variations in cardiac atrial conduction were associated with systemic autonomic symptoms seen during anxiety episodes.³⁰ In a study examining the relationship between the presence of arrhythmia in anxiety disorders and P-wave dispersion, state anxiety was found to effect Pd more than trait anxiety.35 In a study with 30 patients with hypochondriasis (a type of somatoform disorder) and 30 healthy control subjects, Pd durations which were considered to be associated with anxiety were significantly higher in the patient group.³⁶ In another study with patients with panic disorder (a type of anxiety disorder) who experience intense somatic symptoms, 40 patients and controls were examined, and similarly, P-wave dispersion was found to be prolonged in panic disorder patients.³⁰ In our study, anxiety levels were similarly higher; however, there was no statistically significant difference in Pd durations between patients and controls.

QTd, defined as the maximal interlead difference in QT intervals on 12 leads of the surface ECG, reflects the regional heterogeneity of ventricular repolarization and has been suggested as an important marker for risk of arrhythmia in addition to the QT interval. Some investigators proposed that it might be a predisposing factor for arrhythmic events and sudden death. Studies showed that in anxiety disorders like panic disorder and social phobia, QTd was significantly higher than controls. This correlation was considered as a result of prolonged anxiety.4,5 In a study conducted with patients with anxiety, when autonomic innervation and QTd were examined, sympathetic activity was found to be increased, QTd was found to be prolonged, and like hypertension, these were thought to increase the occupation or sitting in the left ventricle.³⁷ In hypertension patients with higher levels of anxiety, QTd and left ventricular hypertrophy were seen to increase, and it was considered to be associated with increased sympathetic and decreased parasympathetic modulation.³⁸ In another study, anxiety levels measured by Trait and State Anxiety Inventory were considered to be a possible marker of prolonged QTd.39

A relationship was found between some psychiatric disorders such as somatoform disorders and circadian heart rate.⁶ Also, in a similar study with psychiatric patients with eating disorder, it was seen that the group with high depression and anxiety scores had a more prolonged QTd.⁴⁰ In a meta-analysis of five studies, it was found that individuals with higher anxiety levels had higher QTd.⁴¹ QTd was found to be significantly increased in patients with Major Depressive Disorder.⁴² Similarly, in our study, depression, anxiety, and QTd levels were significantly higher in patients than in controls.

Our study has several limitations: the number of participants studied was quite small, there is a possibility of presence of higher than normal anxiety levels in patients with CD who were admitted to emergency clinic, and QTd and Pd levels were measured manually.

In conclusion, in psychiatric disorders like generalized anxiety disorder, social phobia, panic disorder, hypochondriasis, and somatoform disorders with heart conduction abnormalities were observed in literature. Similarly, in our study CD patients were also found to have increased QTd. As prolonged QTd may predict an increased risk of ventricular arrhythmias, we believe that CD patients may benefit from being evaluated in terms of cardiological risk of arrhythmia. However, larger samples are needed to clarify the relationship between CD and cardiac conduction disorders.

Disclosure

The authors report no conflicts of interest in this work.

References

- Aytemir K, Ozer N, Atalar E, et al. Dispersion on 12 lead electrocardiography in patients with paroxysmal atrial fibrillation. *PACE*. 2000;23: 109–112.
- Perkiomaki JS, Koistinen MJ, Yli Mayry S, Huikury HV. Dispersion of QT interval in patients with and without susceptibility to ventricular tachyarrhythmias after previous myocardial infarction. *J Am Coll Cardiol.* 1995;26:174–179.
- Unsal C, Kaplan OK, Soner SMZ, Uzunlar B, Uyarel H, Caliskan MP. Wave and QT dispersion in patients with generalized anxiety disorder. *Kosuyolu Kalp Derg.* 2013;16(3):214–219.
- Nahshoni E, Gur S, Marom S, Levin JB, Weizman A, Hermesh H. QT dispersion in patients with social phobia. J Affect Disord. 2004; 78(1):21–26.

- Atmaca M, Yavuzkir M, Izci F, Gurok MG, Adiyaman S. QT wave dispersion in patients with panic disorder. *Neurosci Bull*. 2012;28(3):247–252.
- Stampfer HG. The relationship between psychiatric illness and the circadian pattern of heart rate. *Aust N Z J Psychiatry*. 1998;32(2):187–198.
- Barsky AJ, Cleary PD, Coeytaux RR, Ruskin JN. Psychiatric disorders in medical outpatients complaining of palpitations. *J Gen Intern Med*. 1994;9(6):306–313.
- Ozturk O. Ruh sağlığı ve bozuklukları [Mental health and disorders]. 8th ed. Ankara: Hekimler Yayin Birligi Press; 2001:389–425.
- 9. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. Fourth ed. Washington DC, USA: American Psychiatric Association; 2000. [Text Revision].
- 10. Salgırtay A. Ankara üniversitesi psikiyatri kliniğine bir yıl içinde başvuran acil hastalar üzerinde bir araştırma. Yayınlanmamış Uzmanlık Tezi [A research on emergency patients admitted within one year of the university psychiatric clinic. Unpublished Master Thesis], Ankara Üniversitesi Tıp Fakültesi [Ankara University Medical School], Ankara, 1979.
- Bediz U, Aydemir C, Başterzi AD, Kısa C, Cebeci S, Göka E. Hekimlerin Konversiyon Bozukluğuna Yaklaşımını Etkileyen Faktörler [Affecting factors in approach to conversion disorder of physicians]. *Klinik Psikiyatri*. 2004;7:73–79.
- Alkım T, Tunca Z. In: Köroğlu E, editor. Konversiyon bozukluğu. Psikiyatri Temel Kitabi [Conversion disorder. The Basic Book of Psychiatry]. 1st ed. Ankara: Hekimler Yayın Birliği; 1997:537–544.
- Guggenheim FG. In: Sadock BJ, editor. Somatoform Disorders. Kaplan and Sadock's Synopsis of Psychiatry. 7. ed, (2). Philadelphia: Lippincott Williams and Wilkins; 2000:1504–1514.
- First MB, Spitzer RL, Gibbon M, et al. *Structured Clinical Interview* for DSM-IV Clinical Version (SCID-I/CV). Washington D.C: American Psychiatric Press; 1997.
- Çorapçıoğlu A, Aydemir Ö, Yildiz M. DSM-IV Eksen I Bozukluklari İçin Yapilandirilmiş. Klinik Görüşme Kullanim Kilavuzu (SCID-I) Ankara, Turkey. [Structured Clinical Interview for DSM-IV, Clinical Version] Turkey: Hekimler Yayın Birliği; 1999.
- Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol*. 1988; 56:893–897.
- Ulusoy M, Şahin N, Erkmen H. Turkish version of Beck anxiety inventory: psychometric properties. *J Cogni Psychother*. 1998;12(2): 163–172.
- Beck AT. An inventory for measuring depression. Arch Gen Psychiatry. 1961;7:151–169.
- Hisli N. Beck Depresyon Envanterinin üniversite öğrencileri için geçerliği, güvenirliği [Validity reliability of the Beck Depression Inventory for college students]. *Psikoloji Derg.* 1989;7:3–13.
- Kaplan HI, Sadock BJ, Grebb JA. Somatoform Disorders Synopsis of Psychiatry. 7th ed. Baltimore: Williams and Wilkins; 1994:617–632.
- Saxe GN, Chinman G, Berkowitz R, et al. Somatization in patients with dissociative disorders. *Am J Psychiatry*. 1994;151:1329–1334.
- 22. Özen S, Özbulut Ö, Altındağ Z, Arıcıoğulları Z. Acil serviste konversiyon bozukluğu tanısı konan hastaların sosyodemografik özellikleri, stres faktörleri, I. ve II. Eksen eş tanıları [In the emergency department, diagnosis of conversion disorder socio-demographic characteristics of the patients, stress factors, I. and II. Axis comorbidity]. *Türkiye'de Psikiyatri*. 2000;2:87–96.

Therapeutics and Clinical Risk Management

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peerreviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS,

Submit your manuscript here: http://www.dovepress.com/therapeutics-and-clinical-risk-management-journal and the second sec

- Chandrasekaran R, Goswami U, Sivakumar V, Chitralekha J. Hysterical neurosis – a follow-up study. *Acta Psychiatr Scand*. 1994;89:78–80.
- Bediz AU, Aydemir Ç, Başterzi AD, et al. Attitudes of physician towards patients with conversion disorder. In: *37th National Congress* of Psychiatry Congress Handbook, Istanbul; 2001:78–79.
- Canpolat BI, Göka E, Örsel S, et al. Sociocultural aspects of emergency psychiatry in today's Turkey. In: 5th World Congress International Association For Emergency Psychiatry Congress Handbook; 1998.
- Connor KM, Davidson JR. Generalized anxiety disorder: neurobiological and pharmacotherapeutic perspectives. *Biol Psychiatry*. 1998;44:1286–1294.
- Stanley MA, Beck JG. Anxiety disorders. *Clin Psychol Rev.* 2000; 20(14):731–754.
- Uzbay İT. Anksiyetenin nörobiyolojisi [Neurobiology of anxiety]. Klinik Psikiyatri Dergisi. 2002;5(suppl1):5–13.
- Roelofs K, Naring GW, Moene FC, Hoogduin CA. The question of symptom lateralization in conversion disorder. *J Psychosom Res.* 2000; 49:21–25.
- Yavuzkir M, Atmaca M, Dagli N, et al. P wave dispersion in panic disorder. *Psychosom Med.* 2007;69:344–347.
- Wagner GS. In: Dursun AN, editor. *Marriott'un Pratik Elektrokardiyografisi* [Marriott's Pratic Electrocardiography]. 10th ed. 2001:2–64.
- Castellanos A, Kessler MK, Myerburg RJ. The resting electrocardiogram. In: Sclant CR, Alexander RW, editors. *Hurst's The Heart*. New York: McGraw-Hill; 1994:321.
- Dilaveris PE, Gialafos EJ. Clinical and electrocardiographic predictors of recurrent atrial fibrillation. *Pacing Clin Electrophysiol.* 2000; 23(3):352–358.
- Dilaveris PE, Gialafos EJ. P wave dispersion: a novel predictor of paroxysmal AF. Ann Noninvasive Electrocardiol. 2001;6:159–165.
- 35. Uyarel H, Kasikcioglu H, Dayi SU, et al. Anxiety and P wave dispersion in a healthy young population. *Cardiology*. 2005;104:162–168.
- Atmaca M, Korkmaz H, Korkmaz S. P wave dispersion in patients with hypochondriasis. *Neurosci Lett.* 2010;485(3):148–150.
- Piccirillo G, Viola E, Bucca C, et al. QT interval dispersion and autonomic modulation in subjects with anxiety. *J Lab Clin Med.* 1999; 133(5):461–468.
- Piccirillo G, Elvira S, Viola E, et al. Autonomic modulation of heart rate and blood pressure in hypertensive subjects with symptoms of anxiety. *Clin Sci.* 1998;95:43–52.
- Uyarel H, Okmen E, Cobanoğlu N, Karabulut A, Cam N. Effects of anxiety on QT dispersion in healthy young men. *Acta Cardiol*. 2006;61(1): 83–87.
- Takimoto Y, Yoshiuchi K, Akabayashi A. Effect of mood states on QT interval and QT dispersion in eating disorder patients. *Psychiatry Clin Neurosci.* 2008;62(2):185–189.
- Kelmanson IA. High anxiety in clinically healthy patients and increased QT dispersion: a meta-analysis. *Eur J Prev Cardiol.* 2014;21(12): 1568–1574.
- Nahshoni E, Aizenberg D, Strasberg B, et al. QT dispersion in the surface electrocardiogram in elderly patients with major depression. *J Affect Disorder*. 2000;60:197–200.

Dovepress

EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.