Internet-based adherence interventions for treatment of chronic disorders in adolescents

Alexandria M Bass¹
Michael E Farhangian¹
Steven R Feldman¹⁻³

¹Center for Dermatology Research, Department of Dermatology, Wake Forest School of Medicine, Winston-Salem, NC, USA; ²Department of Pathology, Wake Forest School of Medicine, Winston-Salem, NC, USA; ³Department of Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, NC, USA

Background: Treatment adherence is a ubiquitous challenge in medicine, particularly in the adolescent population with chronic disorders. Web-based adherence interventions may be particularly useful in adolescents, due to their familiarity with and frequent use of the Internet.

Objective: To review web-based interventions used to improve adherence to medication in adolescent patients with chronic disorders.

Methods: A PubMed search was performed for full-text, English, clinical trials in adolescents using keywords “adherence” or “compliance”, “Internet” or “web”, and “treatment” from inception until November 2014. Articles were selected if they involved using the Internet to provide support to adolescents to help improve their adherence to treatment, excluding those focused on solely providing medical services through the Internet and articles focusing on preventative care, rather than treatment of an illness.

Results: Fourteen studies were found concentrating on chronic adolescent disorders. Interventions included online surveys, physician chat lines, monitoring programs, and interactive programs. All interventions experienced either greater improvement in adherence or another disease control measure or no statistically significant difference compared with the control group (in-clinic visits).

Limitations: Few clinical trials studying web-based interventions to improve adherence in adolescents were found. Due to not having one standard outcome measured in all of the studies, it was also difficult comparing the effectiveness of the interventions.

Conclusion: Web-based interventions play a role in improving adherence in adolescents with chronic disorders by being more time saving and cost effective compared with in-clinic visits. Further research focusing on adherence interventions using the Internet is necessary to determine the ideal ways of utilizing the Internet to improve adherence in adolescents with chronic disorders.

Keywords: web, compliance, teenagers, acne, diabetes, pediatrics

Introduction

Over the last few decades in medicine, advances in the research of chronic disorders have changed how physicians manage once fatal conditions in patients. This has led to millions of adolescents living with chronic disorders such as type 1 diabetes, asthma, and HIV due to improved screening, management, and treatment regimens available.¹ However, treatment for these disorders often requires long-term cumbersome regimens consisting of multiple different medications taken multiple times a day, creating a hurdle to good adherence and high treatment failure rates.

Adherence is a challenge faced in many realms of medicine with an adherence rate of only 50%–60% across all patients.² Poor adherence not only leads to poorer

Correspondence: Michael E Farhangian
Center for Dermatology Research,
Department of Dermatology,
Wake Forest School of Medicine,
Medical Center Boulevard, Winston-Salem,
NC 27157-1071, USA
Tel +1 336 716 7740
Fax +1 336 716 7732
Email mfarhang@wakehealth.edu

treatment outcomes, but also leads to increases in health care spending due to emergency department visits, hospitalizations, and office visits. As of 2010, 83% of our health care resources were consumed by people with chronic disorders and non-adherence attributed to 33%–69% of all hospital admissions.³ Children and adolescents also account for a huge part of this problem with 50%–88% of those with chronic conditions being non-adherent to their medication.³

Today's adolescents have bonded with the Internet, with 27% of adolescents using the Internet for over 12 hours some days.⁴ Web-based applications could play a crucial role in supporting adolescent patients to improve their adherence to medication. Our objective was to review web-based interventions used to improve adherence to medication in adolescent patients with chronic disorders.

Methods

A PubMed search was performed using keywords “adherence” or “compliance” and “web” or “Internet” and “treatment” up until November 2014. Studies were filtered to include only full-text clinical trials performed with human subjects

between the ages of 13–18 years old, in the English language. Articles were selected if they involved using the Internet or a web-based program to provide support to adolescents to help improve their adherence to treatment. Studies excluded were those that focused solely on providing medical services through the Internet and articles focusing on preventative care, rather than treatment of an illness. Twelve studies were found utilizing the database. In an effort to find more clinical trials, references of the included studies were screened and two additional articles were included (Figure 1). Only seven studies were found directly measuring adherence, so other disease outcomes, such as severity, quality of life, proper usage of treatments, knowledge about their disease, problem-solving capabilities, and limitations in daily activities were included. We classified the studies in this review based on the type of web-based intervention used to help improve adherence to treatment.

Results

Different web-based approaches such as emails containing surveys, chat lines, and online interactive programs

Figure 1 Preferred reporting items for systematic reviews and meta-analysis (PRISMA) 2009 flow chart.

were evaluated to increase education and adherence of adolescents to their treatment (Table 1). The 14 studies focused on a variety of chronic adolescent disorders, such as acne,⁵ type 1 diabetes,^{6–11} asthma,^{12–16} juvenile idiopathic arthritis,¹⁷ and HIV.¹⁸ The outcomes measured among the studies also varied with only a few studies directly measuring adherence.^{5,9,12,13,15,17} Other studies measured severity,^{6,7,14} quality of life,^{8,11} knowledge and self-efficacy,¹⁸ and/or problem-solving management¹⁰ instead of directly measuring adherence. It was assumed that improvements seen in these measures were due to increased adherence to treatment. The results are summarized according to type of intervention (Table 1).

Web-based online survey

Only one study was found examining the impact of the Internet in improving adherence in patients with acne.⁵ Adolescents, all using a benzoyl peroxide cream, received a weekly email containing a link to an online survey assessing their severity and treatment. At the end of the study, the intervention group had a greater overall adherence rate and a median adherence rate than the control group. The intervention group also experienced a greater improvement in the severity of their acne than the control group; however, these results were not statistically significant. Small sample size could have attributed to insignificant results in changes in severity.⁵

Web-based monitoring programs

Four studies examined monitoring system programs in adolescents with either type 1 diabetes or asthma.^{6,12,13,16} One study examined a website titled “Medtronic Carelink” for adolescents with type 1 diabetes.⁶ This website was an Internet-based blood glucose monitoring system that instructed participants in the intervention group to upload their glucose levels weekly over a 6-month period to a diabetes care team, who could make changes in the patient’s regimen according to their data. After 6 months, hemoglobin A_{1c} (HbA_{1c}) in the intervention group declined and increased in the control group; however, these results were not statistically significant ($P=0.54$). Out of 36 participants in the intervention group, 12 did not submit their blood glucose levels as instructed, leaving few compliant patients to retrieve results from during the study.⁶

Two similar studies by Chan et al in 2003¹² and 2007¹³ also examined an Internet-based monitoring system to improve adherence to treatment, but in asthma patients. The differences were that in the 2003 study, the participants were

required to record and upload their data (such as usage of their steroid inhaler, peak flow meter readings, and asthma symptom diary) to a website to be reviewed by a case manager. In the 2007 study, similar data were studied but instead of uploading it to a website, communication was via virtual visits for a year with the exception of three in-person visits. The control group for both studies received education and monitoring via traditional in-clinic scheduled appointments. After the completion of both studies, no significant differences were seen in adherence, quality of life, or severity. In the 2007 study, the intervention group experienced greater symptom-free days than the control group.^{12,13}

Another study using online monitoring for adolescents with asthma was examined in 2012.¹⁶ Using a website, participants were encouraged to upload weekly parameters of asthma control, as well as have access to two online educational sessions and a personalized online asthma action plan. In conjunction with the website, patients continued to visit with their physician every 3–6 months as needed and for any exacerbations. The control group only had in-clinic visits every 3–6 months with their physician without access to the online monitoring program. After 1 year, no statistically significant differences were seen between the two groups in adherence, severity, quality of life, and knowledge. After 3 months, greater improvement was seen in severity and quality of life in the intervention group; however, these differences were not statistically significant by the completion of the study.¹⁶

Physician chat line

One study examined the efficacy of a weekly 1.5-hour physician mediated chat line for patients with type 1 diabetes.⁷ The topic of each chat, voted on by participants, focused on quality of life and metabolic control such as diabetes management and anxiety about the future. After 2 years, greater improvements in severity were appreciated in the intervention group; however, these results were not statistically significant ($P=0.056$). Overall, all quality of life scores were higher in the intervention group with the greatest improvement found in the “Worries about Diabetes” scores.⁷

Online interactive programs

Eight studies were found evaluating various web-based management programs intended for adolescents with chronic diseases such as Juvenile Idiopathic Arthritis (JIA),¹⁷ type 1 diabetes,^{8–11} HIV,¹⁸ and asthma.^{14,15} Most programs focused on education, management, and problem-solving strategies to improve adherence to treatment. Overall, the interactive

Table 1 Summary of data from 14 studies on web-based interventions to improve adherence in adolescents

Study; year	Internet intervention	Disease studied	Age range and sample size	Was adherence to treatment directly measured? If so, what was the result?	Outcome measurements (other than adherence)	Outcome results (excluding adherence)
Web-based online survey						
Yentzer et al; ⁵ 2011	Internet-based online survey: weekly email containing a URL link to an online survey for 12 weeks	Acne	Age: 13–18 years Size: 15	Yes: MEMS caps; Intervention group: 58%–132% adherence rate; median adherence rate of 74%; Control group: 4%–80% adherence rate, median adherence rate of 32%	Severity: Acne Global Assessment and inflammatory and non-inflammatory lesion counts at baseline, week 6, and week 12	Severity: greater improvement in the intervention group was not statistically significant ($P=0.56$ total lesion counts) ($P=0.49$ non-inflammatory lesions)
Web-based monitoring programs						
Landau et al; ⁶ 2012	Medtronic Carelink: glucose meter data uploaded weekly to this website over a 6-month period	Type 1 diabetes	Age: 11–20 years Size: 70	Not directly measured	Severity: HbA _{1c} level at baseline and at 6 months	Severity: greater improvement in the intervention group was not statistically significant ($P=0.54$)
Chan et al; ¹² 2003	Internet-based video monitoring system: education and monitoring of symptoms, medication usage, and inhaler technique online vs traditional scheduled office visits (control group) over a 6-month period	Asthma	Age: 6–17 years Size: 10	Yes: controller medication usage and inhaler technique; no statistically significant difference between the two groups	Severity: symptom diaries, LFTs, rescue inhaler usage, and utilization of services; Quality of life: PAQLQ	Severity and quality of life: No statistically significant differences
Chan et al; ¹³ 2007	Asthma in-home monitoring: three in-person visits over 1 year with the rest as virtual visits (discussing symptoms, proper inhaler technique, and education) vs six traditional in-clinic visits (control group)	Asthma	Age: 6–17 years Size: 120	Yes: controller medication usage; no statistically significant difference between the two groups	Severity: asthma symptom diaries, LFTs, utilization of services, and rescue therapy usage; Quality of life: PAQLQ	Severity: intervention group experienced greater symptom-free days than the control group (61.1 days vs 51.7 days). No significant differences in other measures of severity or in quality of life
Rikkers-Mutsaerts et al; ¹⁶ 2012	Self-monitoring program: website with two education sessions, weekly monitoring of asthma control, and a personalized electronic action plan in conjunction with regular in-clinic appointments every 3–6 months over a 12-month period	Asthma	Age: 12–18 years Size: 90	Yes: self-reported medication adherence; no significant difference between the two groups ($P=0.12$)	Severity: ACQ and FEV ₁ Quality of Life: PAQLQ; Education: asthma knowledge (via the Consumer Asthma Knowledge Questionnaire) and inhaler technique (via the standardized checklist of the Dutch Asthma Foundation)	Severity: greater improvement in the intervention group at 3 months for asthma control and FEV ₁ but no significant differences between the groups at 12 months ($P=0.75$ and $P=0.35$, respectively) Quality of life: greater improvement in the intervention group at 3 months by 0.40 but no significant differences between the groups at 12 months ($P=0.85$) Education: greater improvement in the intervention group was not statistically significant ($P=0.39$)

Physician chat line						
lafusco et al; ⁷ 2011	Chat line: weekly physician mediated online chat line over a 2-year period	Type I diabetes	Age: 10–18 years Size: 193	Not directly measured	Severity: HbA _{1c} measurements Quality of life: DQOLY	Severity: greater improvement in the intervention group was not statistically significant ($P=0.056$) Quality of life: improvement in the intervention group especially in the “Worries about diabetes” scores compared with a significant impairment found in the control group. “Satisfaction with life” and “Impact of diabetes on life” scores also improved gradually in the intervention group where no change was seen in the control group
Online interactive programs						
Stinson et al; ¹⁷ 2010	“Teens Taking Charge: Managing Arthritis Online”: 12 online modules over 12 weeks consisting of self-management strategies, disease-specific information, and social support	JIA	Age: 12–18 years Size: 46	Yes: CARQ and PARQ scores; no significant differences between the two groups ($P>0.05$)	Quality of life: JAQQ; Pain: RPI; Stress: PSQ; JIA knowledge: MEPS; Self-efficacy: CASE	Pain: intervention group experienced a decrease in average pain intensity compared with an increase seen in the control group; Medical knowledge: increase in intervention group (effect size of 1.32); Quality of life, stress levels, and self-efficacy: No statistically significant differences Intervention group demonstrated greater improvement in quality of life ($P=0.20$), perceived stress ($P=0.20$), coping skills ($P=0.07$), self-efficacy ($P=0.20$), and reported it more enjoyable ($P=0.12$) than the control group
Whittemore et al; ¹¹ 2010	TEENCOPE: interactive website with five weekly sessions focused on improving coping skills vs the control group (Managing Diabetes: online educational program with four weekly sessions) for 6 months	Type I diabetes	Age: 13–16 years Size: 12	Not directly measured	Quality of life: PedsQL; Stress: Perceived Stress Scale; Coping: Issues in Coping with IDDM-Child scale; Self-efficacy: Self-efficacy for Diabetes scale; Psychosocial adjustment: Children's Depression Inventory; Acceptability: 5-item satisfaction survey Severity: HbA _{1c} at baseline and 11 weeks; Problem solving: Diabetes Problem Solving Behaviors scale	
Mulvaney et al; ⁹ 2010	YourWay: web-based support program using videos, expert advice, a personalized webpage, and problem-solving cycles to improve problem solving, self-management, and glycemic control over 11 weeks	Type I diabetes	Age: 13–17 years Size: 72	Yes: Behavior Rating Scale; improved in the intervention group by 0.64 SDs compared with the control group	Severity: HbA _{1c} remained constant in the intervention group compared with an increase in the control group (-0.01% vs $+0.33\%$, respectively); Problem solving: Intervention group improved by 0.3 SDs compared with the control group	
Mulvaney et al; ¹⁰ 2011	YourWay: web-based problem-solving program using videos, problem-solving cycles, social networking, and weekly prompts to improve self-management over 10 weeks	Type I diabetes	Age: 13–17 years Size: 41	Not directly measured	Problem-solving improvement: participant's self-reported progress using the website and a telephone interview at week 10	77% of participants reported the website provided adequate support to solve their self-management and was the fourth best resource (29%) for solving their diabetes problems; 96% reported that reading comments posted by other users was helpful; 45% (mostly) and 45% (totally) were successful in resolving their identified self-management problem

(Continued)

Table 1 (Continued)

Study; year	Internet intervention	Disease studied	Age range and sample size	Was adherence to treatment directly measured? If so, what was the result?	Outcome measurements (other than adherence)	Outcome results (excluding adherence)
Newton and Ashley; ⁸ 2013	Diabetes Teen Talk: website with blogs, discussion forums, and a chat room used to facilitate discussions about a weekly topic over a course of 7 weeks. Participants were encouraged to log in at least three times per week	Type 1 diabetes	Age: 13–18 years Size: 59	Not directly measured	Quality of life: DQOLY, Self-efficacy: Self-Efficacy of Diabetes Self-Management questionnaire; Outcome Expectations: Diabetes Self-Management questionnaire	Outcome expectations: higher positive outcome expectations in the control group than in the intervention group; however, this was seen in both pre- and post-treatment control groups; Quality of life, self-efficacy, and negative outcomes: no statistically significant differences ($P=0.63$, $P=0.53$, $P=0.31$, respectively) Knowledge: improvement (post-intervention mean at 13.1 compared with 11.4 pre-intervention); Self-efficacy: improvement in taking medications in all situations (at the right time every day [9.1 vs 7.8], when away from home [8.5 vs 7.2], when busy at work/school/parties [8.6 vs 6.9], and getting family and friends to help with remembering [8.25 vs 5.6]); Perceived importance of adherence: taking meds at the right time every day (9.3 vs 8.4); Intentions to adhere to treatment: No significant differences
Shegog et al; ¹⁸ 2012	+CLICK: tailored web-based training program providing interactive activities, including animation and peer/expert videos to be accessed during their routine medical visit	HIV	Age: 14–22 years Size: 10	Not directly measured	Knowledge of HIV and ART, self-efficacy, perceived importance of adherence to treatment; Intentions to adhere to treatment for the next 3 months: 5-point rating scale	Knowledge: improvement (post-intervention mean at 13.1 compared with 11.4 pre-intervention); Self-efficacy: improvement in taking medications in all situations (at the right time every day [9.1 vs 7.8], when away from home [8.5 vs 7.2], when busy at work/school/parties [8.6 vs 6.9], and getting family and friends to help with remembering [8.25 vs 5.6]); Perceived importance of adherence: taking meds at the right time every day (9.3 vs 8.4); Intentions to adhere to treatment: No significant differences
Guendelman et al; ¹⁵ 2002	Health Buddy: 12-week self-management and education program that (via a device connected to the home telephone) asked daily questions about severity and asthma trivia. (Control group used a hard-copy asthma diary to monitor their symptoms)	Asthma	Age: Inner city children 8–16 years Size: 134	Yes: questionnaires assessing adherence to medication and use of their asthma symptom diary Intervention group: more likely to take medications without reminders compared with the control group (25 vs 16 participants, respectively)	Limitations in activity and Severity (perceived symptoms, missed school days due to asthma, peak flow readings in the yellow or red zone, use of health care services): from answers to Health Buddy questions	Limitations in activity: fewer in the intervention group (42 participants in the intervention group vs 32 in the control group); Severity: less odds of having peak flow readings in the yellow or red zone in the intervention group than in the control group (38 vs 27 participants, respectively). No significant differences in usage of health services
Joseph et al; ¹⁴ 2013	Puff City: 180-day tailored web-based program focusing on adherence and smoking cessation delivered by a radio D] over four sessions. (Control group received four generic 30 minutes asthma education sessions)	Asthma	Age: 13–18 years (98% African American) Size: 422	Not directly measured	Severity (symptom days, days missed from school due to asthma, utilization of services): survey at baseline, 6 months, and 12 months	Severity: fewer symptom days in the intervention group than in the control group (3.9 vs 5.2 days). No significant differences in emergency department visits, hospitalizations, and days missed from school

Abbreviations: ACQ, Asthma Control Questionnaire; ART, anti-retroviral treatment; CARQ, Child Adherence Report Questionnaire; CASE, Children's Arthritis Self-Efficacy scale; DQOLY, Diabetes Quality of Life for Youth Inventory; IDDM, insulin dependent diabetes mellitus; MEMS caps, Medication Event Monitoring System caps to record the date and time the medication was opened; MEPS, Medical Issues, Exercise, Pain, and Social Support Questionnaire; JAQQ, Juvenile Arthritis Quality of Life Questionnaire; JJA, Juvenile Idiopathic Arthritis; LFTs, Lung Function Tests; including spirometry and peak flow measurements; FEV₁, forced expiratory volume in 1 second; PARQ, Parent Adherence Report Questionnaire; PAQLQ, Pediatric Asthma Quality of Life questionnaire; PedsQL, Pediatric Quality of Life Inventory; PSQ, Perceived Severity of Stress Questionnaire; SDs, standard deviations; HbA_{1c}, hemoglobin A_{1c}; RPI, recalled pain inventory;

programs experienced the most improvements in measured outcomes compared with the control group. All eight studies had greater improvement in at least one measured outcome in the intervention group such as adherence,^{9,15} severity,^{9,14,15} quality of life,¹¹ problem-solving management,^{9,10} coping,¹¹ knowledge,^{17,18} pain,¹⁷ self-efficacy,^{11,18} and outcome expectations⁸ compared with the control group. The other outcomes measured in these studies had no statistically significant difference between the intervention group and the control group.

One program called “Teens Taking Charge: Managing Arthritis Online” was used for adolescents with JIA.¹⁷ This web-based self-management program focused on diagnosis, treatment, and management and was evaluated in 2010 to see whether it improved adherence to treatment. After 12 weeks, there was no significant difference between the two groups in adherence, quality of life, stress levels, or self-efficacy. The intervention group had a decrease in average pain intensity scores compared with the increase seen in the control group, in addition to an increase in medical knowledge.¹⁷ Limitations to this study were having a small sample size, which may have led to insignificant results.

Four interactive programs were used to improve adherence in adolescents with type 1 diabetes.^{8–11} One study evaluated an online interactive website focused on coping skills, called “TEENCOPE”, vs the control group, an online educational website called “Managing Diabetes” over a 6-month period.¹¹ At the completion of the study, the intervention group demonstrated greater improvement in quality of life, perceived stress, coping skills, self-efficacy, and reported it to be more enjoyable than the control group.¹¹

Two studies, performed by Mulvaney et al in 2010⁹ and 2011,¹⁰ focused on a self-management online program titled “YourWay”, indicated for adolescents with type 1 diabetes. The program consisted of multimedia stories, problem-solving cycles, problem-solving expert advice, social networking, and weekly emails to encourage participation. The differences in the two studies were the outcomes measured. After 11 weeks of the 2010 study, adherence and problem solving were higher in the intervention group compared with the control group. Also, severity was maintained in the intervention group compared with worsening severity in the control group.⁹ The 2011 study examined the participants’ opinions of the quality of self-management support, rather than adherence or severity. After 10 weeks, 77% of all of the participants felt that the website provided adequate support to solve their self-management problems. In addition, 45% (mostly) and 45% (totally) were successful in resolving their identified self-management problem.¹⁰ Even though the 2011

study does not focus on improvements in quality of life, severity, or adherence, it provides common adolescent barriers to adherence and an additional way that these problems can be overcome.

“Diabetes Teen Talk” was the third interactive program found for adolescents with type 1 diabetes.⁸ This website consisted of three main areas: blogs, discussion forums, and a chat room used to facilitate discussion about a weekly topic over a course of 7 weeks. No significant difference was seen in quality of life between the two groups, self-efficacy, or negative outcome expectations. A significant difference was seen with the control group reporting a higher positive outcome expectation than individuals in the intervention group; however, this was also seen in both pre- and post-treatment control groups. However, the sample size may have been too small, limiting the study and leading to insignificant results.⁸

“+CLICK” was an online program intended to enhance adherence to treatment among HIV-positive youth based on motivational interviewing principles.¹⁸ The program was customized based on the patient’s reported missed doses, blood counts, self-efficacy, psychosocial issues of perceived importance, and their goals regarding adherence. It provides interactive activities to be accessed during their routine office visit. At the end of the study, improvement was seen in knowledge about HIV and anti-retroviral therapy adherence at post-intervention compared with pre-intervention, in perceived importance of taking medications at the correct time every day, and in self-efficacy about taking their medications. No significant differences were appreciated in psychosocial variables and behavioral intentions to adhere to medications. Although these results do not provide direct information regarding improved adherence, the program may encourage patients to become more involved with their health and in facilitating discussion with health care providers.¹⁸

“Health Buddy”, a program designed to improve asthma self-management in inner city children, was evaluated in 2002.¹⁵ The interactive 12-week program was a device, connected to a home telephone, which was accessed daily. It was programmed to present ten questions pertaining to symptoms, peak flow readings, use of services, activity limitations, and asthma facts and trivia and then record the participant’s answers. The intervention group reported greater improvements in most measures of severity (such as fewer limitations in activity and less odds of peak flow meter readings in the yellow or red zone) and adherence compared with the control group. No significant difference was appreciated between the two groups in usage of health services.¹⁵

In 2013, another Internet-based asthma prevention program was evaluated titled “Puff City”.¹⁴ It was intended for 9th–12th grade students with asthma and focused on three behaviors: controller medication adherence, keeping an inhaler nearby, and smoking reduction/cessation. The program was divided into four sessions (15–30 minutes) over a 180-day period delivered by a radio DJ with tailored information presented to each teenager. No significant differences were appreciated between the two groups except for fewer symptom days were reported in the intervention group compared with the control group.¹⁴

Discussion

Several types of web-based adherence interventions for many chronic disorders were assessed in this review. Due to the variation in outcomes measured among the studies, the comparison between the interventions was rather difficult. However, common trends of effectiveness were seen with adherence and severity measures.

Out of the seven studies that directly measured adherence^{5,9,12,13,15–17} three of them had improvements in adherence in the intervention group^{5,9,15} with the other four having no significant differences between the intervention and control groups.^{12,13,16,17} The three with improvements were interventions using an online survey⁵ and two of the online interactive programs.^{9,15}

Another outcome measured frequently was severity, being measured in all the studies except for five of the interactive program intervention studies.^{8,10,11,17,18} Greater improvements in severity in the intervention group compared with the control group were seen in four of the nine studies evaluating one of the online monitoring programs¹³ and three of the interactive programs.^{9,14,15} No statistically significant differences were seen in the intervention using online surveys,⁵ three of the online monitoring programs,^{6,12,16} and the physician chat line.⁷ All of the online interactive programs that measured severity as an outcome saw significant improvement in some aspect of severity in the intervention group.

Overall, the web-based interventions reviewed were beneficial to patients. In studies where adherence or disease outcomes were measured, there were clear (albeit sometimes not statistically significant) improvements compared with traditional in-clinic appointments used in the control groups. The intervention groups experienced improvements in adherence and/or other aspects of their disease such as severity, quality of life, problem solving, coping, education, and pain levels. Further clinical trials performed with web-based

interventions on improving adherence could provide more answers on their true effectiveness.

Intervention groups differed from the control groups in that they were participating in some form of additional communication, education, and/or monitoring in-between their in-clinic visits. These methods kept patients as active participants in managing their disease, which likely was a large contributor to why the patients in the intervention group experienced greater improvement. Maintaining accountability with patients using web-based interventions may be explained by a similar concept termed “white coat compliance”. This phenomenon is when patients tend to increase their adherence to treatment around the time of follow-up appointments, possibly explaining why decreasing the time between follow-up appointments is an effective way to increase adherence to treatment.¹⁹ Allowing patients to log in online, complete a learning module, or upload data to a website instead of returning for weekly in-clinic follow-up appointments could provide insight to a more convenient method of maintaining adherence in patients.

In addition, even though the interventions reviewed were created to supplement traditional in-clinic visits, utilizing these interventions in lieu of office visits could provide a more time saving and cost-effective way of keeping up with patients to maintain adherence.²⁰ Patients could schedule follow-up appointments farther apart and could avoid missing school/activities, the hassle of driving to the physician’s office, waiting at the office, and paying for the office visit.

There were a couple of limitations to this review article. Most importantly, we found few clinical trials studying web-based interventions to improve adherence in adolescents. In addition, many of the studies did not directly measure adherence, so the assumption was made that improvements in other outcomes measured were due to improved adherence to treatment. Due to not having one standard outcome measured in all of the studies, it was also difficult to compare the effectiveness of the interventions.

Conclusion

Web-based interventions projected on improving adherence in adolescents may play a huge future role in the future of the treatment of chronic disorders. These interventions provide patients with a sense of accountability for managing their disorders in a more accessible way than in the past. Utilizing a source that so many adolescents already use on a daily basis could be the key to success in monitoring and treating patients effectively. Further research focusing on adherence

interventions using the Internet is necessary to determine the ideal ways of implementing the Internet to improve adherence in adolescents with chronic disorders.

Disclosure

The Center for Dermatology Research is supported by an unrestricted educational grant from Galderma Laboratories, LP. Dr Steven R Feldman is a speaker for Janssen and Taro. He is a consultant and speaker for Galderma, Stiefel/GlaxoSmithKline, Abbott Labs and Leo Pharma Inc. He has received grants from Galderma, Janssen, Abbott Labs, Amgen, Stiefel/GlaxoSmithKline, Celgene, and Anacor. He is a consultant for Amgen, Baxter, Caremark, Gerson Lehman Group, Guidepoint Global, Hanall Pharmaceutical Co, Ltd, Kikaku, Lilly, Merck and Co, Inc., Merz Pharmaceuticals, Mylan, Novartis Pharmaceuticals, Pfizer Inc., Qurient, Suncare Research and Xenoport. He is on an advisory board for Pfizer Inc. He is the founder and holds stock in Causa Research and holds stock and is majority owner in Medical Quality Enhancement Corporation. He receives Royalties from UpToDate and Xlibris. Alexandria M Bass and Michael Farhangian report no conflicts of interest in this work.

References

- Compas BE, Jaser SS, Dunn MJ, Rodriguez EM. Coping with chronic illness in childhood and adolescence. *Annu Rev Clin Psychol*. 2012;8:455–480.
- Modi AC, Ingerski LM, Rausch JR, Glauser TA, Drotar D. White coat adherence over the first year of therapy in pediatric epilepsy. *J Pediatr*. 2012;161(4):695–699. e691.
- McGrady ME, Hommel KA. Medication adherence and health care utilization in pediatric chronic illness: a systematic review. *Pediatrics*. 2013;132(4):730–740.
- Kiatrungrit K, Hongsanguansri S. Cross-sectional study of use of electronic media by secondary school students in Bangkok, Thailand. *Shanghai Arch Psychiatry*. 2014;26(4):216–226.
- Yentzer BA, Wood AA, Segransky MJ, et al. An Internet-based survey and improvement of acne treatment outcomes. *Arch Dermatol*. 2011;147(10):1223–1224.
- Landau Z, Mazor-Aronovitch K, Boaz M, et al. The effectiveness of Internet-based blood glucose monitoring system on improving diabetes control in adolescents with type 1 diabetes. *Pediatr Diabetes*. 2012;13(2):203–207.
- Iafusco D, Galderisi A, Nocerino I, et al. Chat line for adolescents with type 1 diabetes: a useful tool to improve coping with diabetes: a 2-year follow-up study. *Diabetes Technol Ther*. 2011;13(5):551–555.
- Newton KT, Ashley A. Pilot study of a web-based intervention for adolescents with type 1 diabetes. *J Telemed Telecare*. 2013;19(8):443–449.
- Mulvaney SA, Rothman RL, Wallston KA, Lybarger C, Dietrich MS. An internet-based program to improve self-management in adolescents with type 1 diabetes. *Diabetes Care*. 2010;33(3):602–604.
- Mulvaney SA, Rothman RL, Osborn CY, Lybarger C, Dietrich MS, Wallston KA. Self-management problem solving for adolescents with type 1 diabetes: intervention processes associated with an Internet program. *Patient Educ Couns*. 2011;85(2):140–142.
- Whittemore R, Grey M, Lindemann E, Ambrosino J, Jaser S. Development of an Internet coping skills training program for teenagers with type 1 diabetes. *Comput Inform Nurs*. 2010;28(2):103–111.
- Chan DS, Callahan CW, Sheets SJ, Moreno CN, Malone FJ. An Internet-based store-and-forward video home telehealth system for improving asthma outcomes in children. *Am J Health Syst Pharm*. 2003;60(19):1976–1981.
- Chan DS, Callahan CW, Hatch-Pigott VB, et al. Internet-based home monitoring and education of children with asthma is comparable to ideal office-based care: results of a 1-year asthma in-home monitoring trial. *Pediatrics*. 2007;119(3):569–578.
- Joseph CL, Ownby DR, Havstad SL, et al. Evaluation of a web-based asthma management intervention program for urban teenagers: reaching the hard to reach. *J Adolesc Health*. 2013;52(4):419–426.
- Guendelman S, Meade K, Benson M, Chen YQ, Samuels S. Improving asthma outcomes and self-management behaviors of inner-city children: a randomized trial of the Health Buddy interactive device and an asthma diary. *Arch Pediatr Adolesc Med*. 2002;156(2):114–120.
- Rikkers-Mutsaerts ER, Winters AE, Bakker MJ, et al. Internet-based self-management compared with usual care in adolescents with asthma: a randomized controlled trial. *Pediatr Pulmonol*. 2012;47(12):1170–1179.
- Stinson JN, McGrath PJ, Hodnett ED, et al. An internet-based self-management program with telephone support for adolescents with arthritis: a pilot randomized controlled trial. *J Rheumatol*. 2010;37(9):1944–1952.
- Shegog R, Markham CM, Leonard AD, Bui TC, Paul ME. “+CLICK”: pilot of a web-based training program to enhance ART adherence among HIV-positive youth. *AIDS Care*. 2012;24(3):310–318.
- Feinstein AR. On white-coat effects and the electronic monitoring of compliance. *Arch Intern Med*. 1990;150(7):1377–1378.
- van der Meer V, van den Hout WB, Bakker MJ, et al. Cost-effectiveness of Internet-based self-management compared with usual care in asthma. *PLoS One*. 2011;6(11):e27108.

Adolescent Health, Medicine and Therapeutics

Publish your work in this journal

Adolescent Health, Medicine and Therapeutics is an international, peer-reviewed, open access journal focusing on health, pathology, and treatment issues specific to the adolescent age group. All aspects of health maintenance, preventative measures and disease treatment interventions are addressed within the journal and practitioners from

Submit your manuscript here: <http://www.dovepress.com/adolescent-health-medicine-and-therapeutics-journal>

Dovepress

all disciplines are invited to submit their work as well as healthcare researchers and patient support groups.. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.