

Umbilical cord blood transplantation supplemented with the infusion of mesenchymal stem cell for an adolescent patient with severe aplastic anemia: a case report and review of literature

Chengxin Luan
Runzhe Chen
Baohan Chen
Jiahua Ding
Ming Ni

Department of Hematology and Oncology (Key Department of Jiangsu Medicine), Zhongda Hospital, Medical School, Southeast University, Nanjing, Jiangsu Province, People's Republic of China

Abstract: Delayed hematopoietic recovery and increased rate of engraftment failure limit the use of umbilical cord blood transplantation (UCBT). We describe a case of severe aplastic anemia treated by UCBT combined with mesenchymal stem cells. Our case reveals that infusing mesenchymal stem cells early (about 40 days) after UCBT may promote hematopoietic recovery. This experience will guide clinical scientists, especially hematologists, to deal with similar situations and encourage them to widen this strategy.

Keywords: cord blood transplantation, mesenchymal stem cell, aplastic anemia

Introduction

The first successful case of umbilical cord blood transplantation (UCBT) was reported in the 1980s.¹ Nowadays, UCBT is recognized as one of the most significant branches of the hematopoietic stem cell transplantation (HSCT) field. However, the delay of hematopoietic reconstruction and increasing probability of engraftment failure resulting from relatively low counts of hematopoietic stem cells contained in a single cord blood (CB) unit prevent the UCBT from widespread use.²⁻⁴ Many strategies such as double-unit unrelated mismatched UCBT,⁴⁻⁸ direct bone marrow infusion,⁹⁻¹¹ and ex vivo expansion¹²⁻¹⁴ have been studied to overcome the above-mentioned limitations.¹⁵ The first description of mesenchymal stem cells (MSCs) by Friedenstein in 1966^{16,17} drew the attention of the biologic field, and the properties of supporting hematopoiesis of MSCs were also shown by Friedenstein in 1974.^{18,19} MSCs are multipotent stromal cells that can differentiate into a variety of cell types, including osteoblasts, chondrocytes, adipocytes, and so on.²⁰⁻²² Due to their multilineage differentiation capacity, hematopoiesis-supporting nature, immunomodulation, and secretion of proregenerative factors,^{23,24} MSCs have been in the focus of intense research for decades.^{21,25} Here we report a case of severe aplastic anemia (SAA) successfully treated by UCBT combined with MSCs.

Case presentation

A 13-year-old Chinese girl who presented with repeated petechia and ecchymoses for 3 days accompanied by nasal bleeding once was admitted to a local hospital. Routine examination of blood revealed that the white blood cell (WBC) count was $2.64 \times 10^9/L$, hemoglobin (HGB) was 66.4 g/L, platelet (PLT) count was $9 \times 10^9/L$, neutrophil count

Correspondence: Baohan Chen
Department of Hematology and Oncology, Zhongda Hospital, Medical School, Southeast University, Dingjiaqiao 87, Gulou District, Nanjing 210009, Jiangsu Province, People's Republic of China
Tel +86 25 8327 2006
Fax +86 25 8327 2011
Email cba8888@hotmail.com

was $0.18 \times 10^9/L$, and reticulocyte was 0.3%. Bone marrow examination showed that the bone marrow was extremely hypoplastic, megakaryocytes were absent, non-hematopoietic cells including plasmocyte, fibrocyte, and lymphocyte were increased, fat cells were increased significantly, hematopoietic cells were rare, and the chromosome karyotype was unremarkable; myelodysplastic syndrome (MDS) was not considered. There was no typical hemoglobinuria, Ham test was negative, and paroxysmal nocturnal hemoglobinuria (PNH) was not considered. She was diagnosed as suffering from SAA. After receiving antithymocyte globulin (ATG), cyclosporin, androgen, and component blood transfusion for 8 months, her condition did not improve.

She was then transferred to our hospital. Bone marrow examination showed that the bone marrow was hypoplastic, pancytopenia was noted, megakaryocytes were absent, and the chromosome karyotype was unremarkable. She matched with her sister's umbilical CB (HLA-A, HLA-B, HLA-DR). She was preconditioned with FC regimen (fludarabine 34 mg/m^2 , day 1 to day 6; cyclophosphamide 53 mg/kg , day 5 to day 6, with day 1 defined as the first day before UCBT day, and so on). She received 33 mL umbilical CB from her sibling sister with mononuclear cell $9.8 \times 10^6/\text{kg}$ and WBC count $14.1 \times 10^9/L$ on October 15, 2013. After transplantation, cyclosporin and mycophenolate mofetil were administered to prevent graft-versus-host disease (GVHD), alprostadil to prevent hepatic vein occlusion disease, and acyclovir to prevent cytomegalovirus (CMV) infection. Fluid infusion, alkalization of urine, and mesna were administered to prevent hemorrhagic cystitis. After transplantation, GVHD manifestation such as diarrhea, skin rash, and liver damage did not occur. Intermittent composition blood transfusion was needed. Routine blood examination on day 1 (day 1 is defined as the first day after UCBT day, and so on) showed WBC $0.15 \times 10^9/L$, HGB 57 g/L, and

PLT $34 \times 10^9/L$; other blood analysis results from day 2 to day 41 are shown in Figure 1. Routine blood examination on November 25, 2013 showed WBC $0.62 \times 10^9/L$, HGB 62 g/L, and PLT $6 \times 10^9/L$. She was still not grafted, and her condition did not improve. She received MSCs (total number 2×10^7 , $4.35 \times 10^5/\text{kg}$, from umbilical cord; Alliancells Bioscience Co., Ltd., People's Republic of China) on November 27, 2013. She had no obvious discomfort during the process or adverse reaction after the process. Routine blood examination before she was discharged showed WBC $3\text{--}5 \times 10^9/L$, HGB 80–90 g/L, and PLT $100\text{--}124 \times 10^9/L$. The shimeric state showed complete donor phenotype. The changing trends of her WBC counts before and after MSC infusion are shown in Figure 1, the HGB counts in Figure 2, and PLT counts in Figure 3. The figures show that after MSC infusion, counts of WBC, HGB, and PLT increased and fluctuated around their normal levels; she was at complete remission and followed up.

Discussion

Compared to bone marrow transplantation (BMT), UCBT provides many advantages. The acquisition of CB is easy, and the collection is harmless both to the mother and the newborn infant. Also, HSCs from the CB graft can be cryopreserved and transplanted to the host after thawing without losing its reproducing ability. Broxmeyer's study found no significant differences of nucleated cells, granulocyte-macrophage (CFU-GM), erythroid (BFU-E), and multipotential (CFU-GEMM) progenitors after cryopreservation of CB for 10 years.²⁶ Yamamoto et al reported that the recovery rate of total nucleated cell (TNC) count, CD34⁺ cell count, and CFU-GM number was not significantly different between the study group (18 CB units, collected between April 1998 and September 1998) and control group (18 CB units, collected between May 2008




Figure 1 Counts of WBC from day 1 to day 131.

Note: After MSCs infusion, counts of WBC increased and fluctuated around normal level.

Abbreviations: WBC, white blood cell; MSC, mesenchymal stem cell.




Figure 2 Counts of HGB from day 1 to day 131.

Note: After MSCs infusion, counts of HGB increased and fluctuated near normal level.

Abbreviations: HGB, hemoglobin; MSC, mesenchymal stem cell.

and June 2008).²⁷ The latest study by Mitchell et al showed that UCB units after cryopreservation for at least 10 years had no impact on the clinical outcome.²⁸ The incidence of GVHD related to UCBT is generally lower than in BMT. According to Hwang et al GVHD incidence in unrelated UCBT is not higher than unrelated-donor BMT in pediatric patients: chronic GVHD (cGVHD) is lower in UCBT, and acute GVHD (aGVHD) (III–IV) is no different.²⁹ Zhang et al reported that the incidence rates of both aGVHD and cGVHD were significantly lower in the UBMT group compared with the BMT group in acute leukemia patients.³⁰ But Chen et al reported that the incidence of Grade 2–4 and 3–4 aGVHD was increased in the UBMT group compared with the BMT group and that the incidence of extensive cGVHD was significantly lower in UCBT.³¹ Another obvious advantage of UCBT compared with BMT is the relatively low human leukocyte antigen (HLA)-matching requirements. A high degree of HLA matching is generally required in BMT, namely 6 of 6 (HLA-A, HLA-B, and HLA-DR loci),

and only 30% of patients needing allogeneic HSCT can have access to a matched sibling donor.² For UCBT, a minimum 4 of 6 HLA allele matching is required, but 5 of 6 is ideal.³²

While hematopoiesis recovery is slow,^{2–4} and infectious complication incidence is high in CB recipients.^{33,34} Delayed hematopoietic recovery and increased rate of engraftment failure limit the use of UCBT. Delayed hematopoietic recovery could result in infection complication and increased transplantation-related mortality (TRM). UCBT outcomes have been proven to be closely associated with CD34⁺ dose or TNC dose.³⁵ Wagner et al³⁶ reported that the dose of CD34⁺ cells was a significant factor related to the rate of engraftment, TRM, and survival in Cox regression analyses. Eapen et al³⁷ found that with the TNC >3.0×10⁷/kg, the overall outcome significantly improved in children with acute leukemia. The CD34⁺ dose in a CB graft is on average 1–2 log smaller than in an unrelated BM or peripheral blood stem cell graft.³⁸ Many strategies mentioned in the Introduction are utilized to overcome the low dose limitation in CB.




Figure 3 Counts of PLT from day 1 to day 131.

Note: After MSCs infusion, counts of PLT increased and fluctuated around normal level.

Abbreviations: PLT, platelet; MSC, mesenchymal stem cell.




Figure 4 Treatment schema of SAA.

Abbreviations: HLA, human leukocyte antigen; SAA, severe aplastic anemia; IST, immunosuppressive therapy; HCT, hematopoietic cell transplantation.

MSCs can be found in various tissues, such as peripheral blood, umbilical cord and placenta, bone marrow. Because of their properties such as multilineage differentiation capacity, hematopoiesis-supporting nature, immunomodulation, and secretion of proregenerative factors, MSCs have been regarded as potential therapeutic agents in many clinical diseases, in particular for the treatment of hematologic and immunological disorders. MSCs are used to treat autoimmune diseases as well. Constantin et al³⁹ reported that MSCs derived from adipose administered intravenously before onset of the disease significantly mitigated the severity of autoimmune encephalomyelitis (EAE) due to their immunomodulation properties. As regard to hematological disorders, MSCs are reported to be studied and used in various disorders, maybe due to their properties of hematopoiesis support and immunomodulation. Robinson et al⁴⁰ showed that a 10–20-fold increase in TNCs, a 7–18-fold increase in progenitor cells, and a 16–37-fold increase in CD34⁺ cells could be achieved by coculturing CB cells with MSCs. Nauta et al⁴¹ reported that significant enhancement of long-term engraftment with better tolerance to host and donor antigens was achieved by the addition of host MSCs. The underlying

mechanism of the immunomodulation of MSCs is not very clear, but growing evidence reveals that MSCs exhibit immunosuppressive activity on T cells,^{42–46} B cells,^{47,48} and natural killer (NK) cells.^{49–51} The underlying mechanism of hematopoiesis support of MSCs is supposed to be the effect of supporting the reconstruction of hematopoietic microenvironment in addition to direct hematopoiesis.^{52–54}

Aplastic anemia (AA) is a disease in which the bone marrow and the blood stem cells are damaged, which causes pancytopenia, including anemia, leukopenia, and thrombocytopenia. Its causes are various, and it is regarded as an autoimmune disorder⁵⁵ in which immune cells (mainly T cells)⁵⁶ attack the bone marrow and cause pancytopenia. Its first-line treatment consists of hematopoietic cell transplantation and administration of immunosuppressive drugs. SAA is life-threatening and needs immediate treatment. Adolescents and young adults (age <30 years) with SAA who have an HLA-matched sibling donor should proceed directly to hematopoietic cell transplantation;^{57–59} the treatment scheme⁵⁷ is shown in Figure 4. But this treatment scheme does not consider the situation in which the patients do not respond to hematopoietic cell transplantation.

Table 1 Basic clinical parameters of the other four UCBT patients

	Patient 1	Patient 2	Patient 3	Patient 4
Age (years)	12	25	41	41
Sex	Female	Male	Male	Female
Diagnosis	M5	CML	M4	CML
Diagnosis time	9/2003	11/2005	2/2012	9/2002
Main therapeutic schedule	MA and DA	Hydroxyurea	DA	Hydroxyurea
UCBT time, volume, and dose	3/3/2004, 30 mL, missing	11/7/2008, 42 mL, 1.2×10 ⁷ /kg	4/26/2012, 5 mL, 2.69×10 ⁷ /kg	3/8/2005, N/A, N/A
Interval between UCBT and death	20 days	45 days	85 days	40 days
Whether grafted	Not grafted	Grafted with acute GVHD	Grafted with acute GVHD	Not grafted
Main complication	Central nervous system infection, intracranial hemorrhage	Septic shock, DIC	Respiratory failure, severe pneumonia, acute renal failure	Interstitial pneumonia and acute left heart failure

Abbreviations: M5, acute monocytic leukemia; CML, chronic myelocytic leukemia; M4, acute myelomonocytic leukemia; MA, mitoxantrone + Ara-c; DA, daunorubicin + Ara-c; UCBT, umbilical cord blood transplantation; DIC, disseminated intravascular coagulation; GVHD, graft-versus-host disease; N/A, not applicable.

There have been reports in the literature on the use of MSCs during allogeneic transplantation to enhance engraftment.^{60–62} Our patient was an SAA case. She matched with her sister's umbilical CB and was without transplantation-associated contraindications, but she did not respond to immunosuppressive therapy (IST) or UCBT. After failure of the initial UCBT treatment, considering AA is of pancytopenia and an immunity-related disease and the limitation of delayed hematopoietic recovery and increased rate of engraftment failure of UCBT discussed above, we decided to apply MSCs to this patient for its hematopoiesis-supporting properties and immunomodulation. After MSC infusion, the WBC count increased and fluctuated around the normal level; she was on complete remission and followed up.

Before this case, our department had handled four cases of UCBT and they all failed. Their basic clinical parameters are listed in Table 1. The first patient died of central nervous system infection and intracranial hemorrhage. The second one died of septic shock and disseminated intravascular coagulation. The third died of respiratory failure, severe pneumonia, and acute renal failure. The fourth died of interstitial pneumonia and acute left heart failure. Infection was a common cause of death of the four patients after receiving UCBT, and the mean interval between UCBT and death was 47.5 days.

It is still challenging to deal with delayed hematopoietic recovery and to increase the rate of engraftment after UCBT. As discussed above, delayed hematopoietic recovery easily results in infection, and it is hard to control infection if hematopoietic recovery is delayed.

Our case reveals that infusing MSCs early (about 40 days) after UCBT may promote hematopoietic recovery. However, as this is a single case study without a controlled trial, we

cannot be sure whether the MSC infusion had any effect in promoting hematopoietic recovery, or whether the initial CB engraftment was just delayed or infusion of MSC with engraftment was just coincidental. Further studies such as controlled trials and more cases are needed to prove the effect of MSC infusion to promote hematopoietic recovery in UCBT. But surely, the hematopoietic-supporting and immunomodulation properties of MSC are increasingly attracting the attention of scientists and clinicians, and our experience will encourage clinical scientists, especially hematologists, to elaborate this strategy while dealing with a similar situation or another.

Acknowledgments

This work was supported by the National Natural Science Foundation of People's Republic of China (Grant no 81170492, 81370673), the National High Technology Research and Development Program 863 of People's Republic of China (Grant no 2012AA022703), the National Key Basic Research Program 973 of People's Republic of China (Grant no 2010CB732404), the Key Medical Projects of Jiangsu Province (Grant no BL2014078), and the Key Discipline of Jiangsu Province (2011–2015).

Disclosure

The authors declare no conflicts of interest in this work.

References

1. Gluckman E, Broxmeyer HA, Auerbach AD, et al. Hematopoietic reconstitution in a patient with Fanconi's anemia by means of umbilical-cord blood from an HLA-identical sibling. *N Engl J Med*. 1989;321(17):1174–1178.
2. Oran B, Shpall E. Umbilical cord blood transplantation: a maturing technology. *Hematology Am Soc Hematol Educ Program*. 2012;2012:215–222.

3. Rocha V, Gluckman E. Improving outcomes of cord blood transplantation: HLA matching, cell dose and other graft- and transplantation-related factors. *Br J Haematol*. 2009;147(2):262–274.
4. De Lima M, St John LS, Wiedner ED, et al. Double-chimaerism after transplantation of two human leucocyte antigen mismatched, unrelated cord blood units. *Br J Haematol*. 2002;119(3):773–776.
5. Sideri A, Neokleous N, Brunet De La Grange P, et al. An overview of the progress on double umbilical cord blood transplantation. *Haematologica*. 2011;96(8):1213–1220.
6. Yin Y, Ren HY, Cen XA, et al. Long-term outcomes in adults with leukemia treated with transplantation of two unrelated umbilical cord blood units. *Chin Med J (Engl)*. 2011;124(16):2411–2416.
7. Huang SL, Zhou DH. 非血缘异基因脐血造血干细胞移植现状: 问题与对策 [Unrelated allogeneic umbilical cord blood transplantation: present status, problems and countermeasures]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi*. 2009;17(1):1–7. Chinese.
8. Hashem H, Lazarus HM. Double umbilical cord blood transplantation: relevance of persistent mixed-unit chimerism. *Biol Blood Marrow Transplant*. 2015;21(4):612–619.
9. Rocha V, Labopin M, Ruggeri A, et al. Unrelated cord blood transplantation: outcomes after single-unit intrabone injection compared with double-unit intravenous injection in patients with hematological malignancies. *Transplantation*. 2013;95(10):1284–1291.
10. Frasson F, Gualandi F, Podesta M, et al. Direct intrabone transplant of unrelated cord-blood cells in acute leukaemia: a phase I/II study. *Lancet Oncol*. 2008;9(9):831–839.
11. Brunstein CG, Barker JN, Weisdorf DJ, et al. Intra-BM injection to enhance engraftment after myeloablative umbilical cord blood transplantation with two partially HLA-matched units. *Bone Marrow Transplant*. 2009;43(12):935–940.
12. Flores-Guzman P, Fernandez-Sanchez V, Mayani H. Concise review: ex vivo expansion of cord blood-derived hematopoietic stem and progenitor cells: basic principles, experimental approaches, and impact in regenerative medicine. *Stem Cells Transl Med*. 2013;2(11):830–838.
13. Ge J, Cai H, Li Q, Du Z, Tan WS. Effects of telomerase activity and apoptosis on ex vivo expansion of cord blood CD34(+) cells. *Cell Prolif*. 2013;46(1):38–44.
14. Horwitz ME, Frasson F. Improving the outcome of umbilical cord blood transplantation through ex vivo expansion or graft manipulation. *Cytotherapy*. 2015;17(6):730–738.
15. Yu X, Gu Z, Wang Y, Wang H. New strategies in cord blood cells transplantation. *Cell Biol Int*. 2013;37(9):865–874.
16. Friedenstein AJ, Piatetzky S 2nd, Petrakova KV. Osteogenesis in transplants of bone marrow cells. *J Embryol Exp Morphol*. 1966;16(3):381–390.
17. Chao YH, Wu HP, Chan CK, Tsai C, Peng CT, Wu KH. Umbilical cord-derived mesenchymal stem cells for hematopoietic stem cell transplantation. *J Biomed Biotechnol*. 2012;2012:759503.
18. Friedenstein AJ, Chailakyan RK, Latsinik NV, Panasyuk AF, Keiliss-Borok IV. Stromal cells responsible for transferring the microenvironment of the hemopoietic tissues. Cloning in vitro and retransplantation in vivo. *Transplantation*. 1974;17(4):331–340.
19. Resnick I, Stepensky P, Elkin G, et al. MSC for the improvement of hematopoietic engraftment. *Bone Marrow Transplant*. 2010;45(3):605–606.
20. Harris DT. Umbilical cord tissue mesenchymal stem cells: characterization and clinical applications. *Curr Stem Cell Res Ther*. 2013;8(5):394–399.
21. Sharma RR, Pollock K, Hubel A, McKenna D. Mesenchymal stem or stromal cells: a review of clinical applications and manufacturing practices. *Transfusion*. 2014;54(5):1418–1437.
22. Soleymaninejadian E, Pramanik K, Samadian E. Immunomodulatory properties of mesenchymal stem cells: cytokines and factors. *Am J Reprod Immunol*. 2012;67(1):1–8.
23. Serakinci N, Fahrioglu U, Christensen R. Mesenchymal stem cells, cancer challenges and new directions. *Eur J Cancer*. 2014;50(8):1522–1530.
24. Vanikar AV, Trivedi HL, Kumar A, Gopal SC, Kute VB. Mesenchymal stem cells and transplant tolerance. *Nephrology (Carlton)*. 2014;19(7):369–374.
25. Bieback K, Wuchter P, Besser D, et al. Mesenchymal stromal cells (MSCs): science and friction. *J Mol Med (Berl)*. 2012;90(7):773–782.
26. Broxmeyer HE, Cooper S. High-efficiency recovery of immature haematopoietic progenitor cells with extensive proliferative capacity from human cord blood cryopreserved for 10 years. *Clin Exp Immunol*. 1997;107(suppl 1):45–53.
27. Yamamoto S, Ikeda H, Toyama D, et al. Quality of long-term cryopreserved umbilical cord blood units for hematopoietic cell transplantation. *Int J Hematol*. 2011;93(1):99–105.
28. Mitchell R, Wagner JE, Brunstein CG, et al. Impact of long-term cryopreservation on single umbilical cord blood transplantation outcomes. *Biol Blood Marrow Transplant*. 2015;21(1):50–54.
29. Hwang WY, Samuel M, Tan D, Koh LP, Lim W, Linn YC. A meta-analysis of unrelated donor umbilical cord blood transplantation versus unrelated donor bone marrow transplantation in adult and pediatric patients. *Biol Blood Marrow Transplant*. 2007;13(4):444–453.
30. Zhang H, Chen J, Que W. A meta-analysis of unrelated donor umbilical cord blood transplantation versus unrelated donor bone marrow transplantation in acute leukemia patients. *Biol Blood Marrow Transplant*. 2012;18(8):1164–1173.
31. Chen YH, Xu LP, Liu DH, et al. Comparative outcomes between cord blood transplantation and bone marrow or peripheral blood stem cell transplantation from unrelated donors in patients with hematologic malignancies: a single-institute analysis. *Chin Med J (Engl)*. 2013;126(13):2499–2503.
32. Wang F, He J, Chen S, et al. HLA-A, HLA-B, HLA-DRB1 allele and haplotype frequencies in 6,384 umbilical cord blood units and transplantation matching and engraftment statistics in the Zhejiang cord blood bank of China. *Int J Immunogenet*. 2014;41(1):13–19.
33. Parody R, Martino R, Rovira M, et al. Severe infections after unrelated donor allogeneic hematopoietic stem cell transplantation in adults: comparison of cord blood transplantation with peripheral blood and bone marrow transplantation. *Biol Blood Marrow Transplant*. 2006;12(7):734–748.
34. Danby R, Rocha V. Improving engraftment and immune reconstitution in umbilical cord blood transplantation. *Front Immunol*. 2014;5:68.
35. Gluckman E, Rocha V, Arcese W, et al. Factors associated with outcomes of unrelated cord blood transplant: guidelines for donor choice. *Exp Hematol*. 2004;32(4):397–407.
36. Wagner JE, Barker JN, DeFor TE, et al. Transplantation of unrelated donor umbilical cord blood in 102 patients with malignant and non-malignant diseases: influence of CD34 cell dose and HLA disparity on treatment-related mortality and survival. *Blood*. 2002;100(5):1611–1618.
37. Eapen M, Rubinstein P, Zhang MJ, et al. Outcomes of transplantation of unrelated donor umbilical cord blood and bone marrow in children with acute leukaemia: a comparison study. *Lancet*. 2007;369(9577):1947–1954.
38. Lindemans CA, Van Besien K. Topping it up: methods to improve cord blood transplantation outcomes by increasing the number of CD34+ cells. *Cytotherapy*. 2015;17(6):723–729.
39. Constantin G, Marconi S, Rossi B, et al. Adipose-derived mesenchymal stem cells ameliorate chronic experimental autoimmune encephalomyelitis. *Stem Cells*. 2009;27(10):2624–2635.
40. Robinson SN, Ng J, Niu T, et al. Superior ex vivo cord blood expansion following co-culture with bone marrow-derived mesenchymal stem cells. *Bone Marrow Transplant*. 2006;37(4):359–366.
41. Nauta AJ, Westershuis G, Kruisselbrink AB, Lurvink EG, Willemze R, Fibbe WE. Donor-derived mesenchymal stem cells are immunogenic in an allogeneic host and stimulate donor graft rejection in a nonmyeloablative setting. *Blood*. 2006;108(6):2114–2120.
42. Di Nicola M, Carlo-Stella C, Magni M, et al. Human bone marrow stromal cells suppress T-lymphocyte proliferation induced by cellular or nonspecific mitogenic stimuli. *Blood*. 2002;99(10):3838–3843.

43. Zhao K, Lou R, Huang F, et al. Immunomodulation effects of mesenchymal stromal cells on acute graft-versus-host disease after hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2015;21(1):97–104.
44. Yan Z, Zhuansun Y, Chen R, Li J, Ran P. Immunomodulation of mesenchymal stromal cells on regulatory T cells and its possible mechanism. *Exp Cell Res*. 2014;324(1):65–74.
45. Bartholomew A, Sturgeon C, Siatskas M, et al. Mesenchymal stem cells suppress lymphocyte proliferation in vitro and prolong skin graft survival in vivo. *Exp Hematol*. 2002;30(1):42–48.
46. Abumaree M, Al Jumah M, Pace RA, Kalionis B. Immunosuppressive properties of mesenchymal stem cells. *Stem Cell Rev*. 2012;8(2):375–392.
47. Corcione A, Benvenuto F, Ferretti E, et al. Human mesenchymal stem cells modulate B-cell functions. *Blood*. 2006;107(1):367–372.
48. Deng W, Han Q, Liao L, You S, Deng H, Zhao RC. Effects of allogeneic bone marrow-derived mesenchymal stem cells on T and B lymphocytes from BXS mice. *DNA Cell Biol*. 2005;24(7):458–463.
49. Wang HF, Shi YJ, Ren HY. 骨髓间充质干细胞对异体自然杀伤细胞调节作用的研究 [Bone marrow-derived mesenchymal stem cells regulate the proliferation and activity of natural killer cells]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi*. 2012;20(2):438–442. Chinese.
50. Li Y, Qu YH, Wu YF, et al. Bone marrow mesenchymal stem cells suppressing activation of allogeneic cytokine-induced killer/natural killer cells either by direct or indirect interaction. *Cell Biol Int*. 2015;39(4):435–445.
51. Spaggiari GM, Capobianco A, Becchetti S, Mingari MC, Moretta L. Mesenchymal stem cell-natural killer cell interactions: evidence that activated NK cells are capable of killing MSCs, whereas MSCs can inhibit IL-2-induced NK-cell proliferation. *Blood*. 2006;107(4):1484–1490.
52. Pontikoglou C, Deschaseaux F, Sensebe L, Papadaki HA. Bone marrow mesenchymal stem cells: biological properties and their role in hematopoiesis and hematopoietic stem cell transplantation. *Stem Cell Rev*. 2011;7(3):569–589.
53. Huo SW, Zhang Y. 间充质干细胞在造血调控中的作用 [Mesenchymal stem cells in hematopoietic regulation – review]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi*. 2006;14(1):187–190. Chinese.
54. Hao L, Sun H, Wang J, Wang T, Wang M, Zou Z. Mesenchymal stromal cells for cell therapy: besides supporting hematopoiesis. *Int J Hematol*. 2012;95(1):34–46.
55. Young NS. Current concepts in the pathophysiology and treatment of aplastic anemia. *Hematology Am Soc Hematol Educ Program*. 2013;2013:76–81.
56. Weston W, Gupta V, Adkins R, Jurecic R. New therapeutic approaches for protecting hematopoietic stem cells in aplastic anemia. *Immunol Res*. 2013;57(1–3):34–43.
57. DeZern AE, Guinan EC. Aplastic anemia in adolescents and young adults. *Acta Haematol*. 2014;132(3–4):331–339.
58. Hartung HD, Olson TS, Bessler M. Acquired aplastic anemia in children. *Pediatr Clin North Am*. 2013;60(6):1311–1336.
59. Kurre P, Johnson FL, Deeg HJ. Diagnosis and treatment of children with aplastic anemia. *Pediatr Blood Cancer*. 2005;45(6):770–780.
60. Ball LM, Bernardo ME, Roelofs H, et al. Cotransplantation of ex vivo expanded mesenchymal stem cells accelerates lymphocyte recovery and may reduce the risk of graft failure in haploidentical hematopoietic stem-cell transplantation. *Blood*. 2007;110(7):2764–2767.
61. Le Blanc K, Samuelsson H, Gustafsson B, et al. Transplantation of mesenchymal stem cells to enhance engraftment of hematopoietic stem cells. *Leukemia*. 2007;21(8):1733–1738.
62. Macmillan ML, Blazar BR, DeFor TE, Wagner JE. Transplantation of ex-vivo culture-expanded parental haploidentical mesenchymal stem cells to promote engraftment in pediatric recipients of unrelated donor umbilical cord blood: results of a phase I-II clinical trial. *Bone Marrow Transplant*. 2009;43(6):447–454.

Patient Preference and Adherence

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focuses on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize

Submit your manuscript here: <http://www.dovepress.com/patient-preference-and-adherence-journal>

clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress