

# Daily rhythm variations of the clock gene *PER1* and cancer-related genes during various stages of carcinogenesis in a golden hamster model of buccal mucosa carcinoma

Hua Ye  
Kai Yang  
Xue-Mei Tan  
Xiao-Juan Fu  
Han-Xue Li

Department of Oral and Maxillofacial Surgery, The First Affiliated Hospital of Chongqing Medical University, Chongqing, People's Republic of China

**Background:** Recent studies have demonstrated that the clock gene *PER1* regulates various tumor-related genes. Abnormal expressions and circadian rhythm alterations of *PER1* are closely related to carcinogenesis. However, the dynamic circadian variations of *PER1* and tumor-related genes at different stages of carcinogenesis remain unknown. This study was conducted to investigate the daily rhythm variation of *PER1* and expression of tumor-related genes *VEGF*, *KI67*, *C-MYC*, and *P53* in different stages of carcinogenesis.

**Materials and methods:** Dimethylbenzanthracene was used to establish a golden hamster model of buccal mucosa carcinogenesis. Hamsters with normal buccal mucosa, precancerous lesion, and cancerous lesion were sacrificed at six different time points during a 24-hour period of a day. Pathological examination was conducted using routine hematoxylin and eosin staining. *PER1*, *VEGF*, *KI67*, *C-MYC*, and *P53* mRNAs were detected by real-time reverse transcriptase polymerase chain reaction, and a cosinor analysis was applied to analyze the daily rhythm.

**Results:** *PER1*, *VEGF*, *C-MYC*, and *P53* mRNA exhibited daily rhythmic expression in three carcinogenesis stages, and *KI67* mRNA exhibited daily rhythmic expression in the normal and precancerous stages. The daily rhythmic expression of *KI67* was not observed in cancerous stages. The mesor and amplitude of *PER1* and *P53* mRNA expression decreased upon the development of cancer ( $P < 0.05$ ), whereas the mesor and amplitude of *VEGF*, *KI67*, and *C-MYC* mRNA increased upon the development of cancer ( $P < 0.05$ ). Compared with the normal tissues, the acrophases of *PER1*, *VEGF*, and *C-MYC* mRNA occurred earlier, whereas the acrophases of *P53* and *KI67* mRNA lagged remarkably in the precancerous lesions. In the cancer stage, the acrophases of *VEGF* and *C-MYC* mRNA occurred earlier and later, respectively, compared with the normal stage.

**Conclusion:** Variations in the daily rhythm characteristics of the clock gene *PER1* and tumor-related genes *VEGF*, *KI67*, *C-MYC*, and *P53* correlate with the development of cancer. Additional studies might provide new insights and methods to explore carcinogenic mechanisms and cancer treatment.

**Keywords:** circadian rhythm, clock gene, *PER1*, tumor, carcinogenesis

Correspondence: Kai Yang  
Department of Oral and Maxillofacial Surgery, The First Affiliated Hospital of Chongqing Medical University, No 1, Youyi Road, Yuzhong District, Chongqing 400016, People's Republic of China  
Tel +86 23 8901 2569  
Fax +86 23 8901 2569  
Email cqfyyk@hotmail.com

## Introduction

"Circadian" is a Latin word that means "about a day".<sup>1</sup> Living organisms on earth have evolved with an endogenous timing system, specifically the circadian clock.<sup>1-3</sup> Current research has demonstrated that numerous vital physiological processes that occur in mammals in vivo, such as sleeping and waking, hormone secretion, and immune activity, exhibit a circadian fluctuation of approximately 24 hours.<sup>1,4-7</sup>

The generation of a circadian rhythm in vivo depends on circadian variations in clock genes expression. At least 14 known core clock genes have been reported, including *PER1*, *PER2*, *PER3*, *CRY1*, *CRY2*, *TIM*, *CK1 $\epsilon$* , *CLOCK*, *BMAL1*, *RORs*, *REV-ERBs*, *NPAS2*, *DEC1*, and *DEC2*.<sup>8–11</sup> These clock genes are located in the suprachiasmatic nuclei of the anterior hypothalamus and probably in all peripheral tissue cells.<sup>12–14</sup> The former is called the central circadian clock, and the latter is the peripheral circadian clock.<sup>15,16</sup> Circadian rhythms in the central circadian and peripheral circadian clocks are generated by the interaction of positive and negative transcription–translation feedback loops at the molecular level. The transcriptional regulators *CLOCK* and *BMAL1* form *CLOCK/BMAL1* heterodimers, and the heterodimers subsequently bind to the E-box of the *PER* (*PER1*, *PER2*, and *PER3*) and *CRY* (*CRY1* and *CRY2*) genes to activate the expression of these genes in the nuclei. These processes are the main components of the positive feedback circadian rhythm loop.<sup>17,18</sup> *PER* and *CRY* proteins form heterodimers that translocate from the cytoplasm to the nucleus, where they interfere with *CLOCK/BMAL1* heterodimer formation and inhibit the expressions of *PER* and *CRY*, thereby forming a negative feedback loop.<sup>11,19,20</sup> The clock genes generate and sustain rhythmic expressions in a 24-hour period via these feedback loops.<sup>2,12</sup> The peripheral circadian clock can be regulated and influenced by the central circadian clock; meanwhile, the peripheral circadian clock can also generate and sustain a 24-hour circadian rhythm independently.<sup>2,16</sup> Approximately 5%–10% of the genes of the mammal genome are regulated by clock genes,<sup>3,21</sup> which exhibit 24-hour periodic expression, and these genes are known as clock-controlled genes (CCGs).<sup>12,22</sup> Importantly, the circadian expression of clock genes and CCGs maintains a high level of synchronization and orderliness for normal physiological activities.<sup>23,24</sup>

Current reports have demonstrated that the *PER1* gene is an important clock gene that exhibits circadian rhythms in normal cells.<sup>25,26</sup> *PER1* is a cancer suppressor gene that is downregulated in various solid tumor cells, including prostatic cancer, melanoma, and oral squamous cell carcinoma.<sup>8,27,28</sup> Many important cancer-related genes are CCGs; *PER1* is an important clock gene that regulates many cancer-related genes. Gery et al reported that *PER1* overexpression in human colon cancer cells (HCT116) increased *P53* and *C-MYC* expression.<sup>29</sup> Soták et al demonstrated that reduced *PER1* mRNA expression in mice colorectal tumors promoted the upregulation of *C-MYC* and downregulation of *P21*.<sup>15</sup>

Current studies have demonstrated that the clock gene *PER1* regulates numerous downstream cancer-related genes to influence cell proliferation and apoptosis as well as tumor angiogenesis; thus, *PER1* is closely related to the occurrence and development of carcinoma.<sup>15,29,30</sup> Carcinogenesis is a process of pathological change involving multiple genes and steps. The dynamic circadian rhythm variations (including the expressions of mesor, amplitude, and acrophase) of the clock gene *PER1* and cancer-related CCGs involved in the stages of carcinogenesis and the development of carcinoma are unclear. The daily profiles of the expression of clock gene *PER1* and cancer-related genes *VEGF*, *KI67*, *C-MYC*, and *P53* during three stages of carcinogenesis in a golden hamster model of buccal mucosa carcinoma were detected in this study; this research provides new concepts for further research on the mechanism of carcinogenesis and the development of cancer as well as the development of cancer therapies based on the recovery of circadian rhythms.

## Materials and methods

### Experimental animals

Ninety male-specific pathogen-free Syrian golden hamsters (6–7 weeks, 90–120 g) were purchased from Vital River Company (Beijing, People's Republic of China). All experimental procedures were approved by the Experimental Animal Use and Management Committee in Experimental Animal Research Institute, Chongqing Medical University.

### Establishment of a Syrian golden hamster model of buccal mucosa carcinoma by dimethylbenzanthracene

Ninety golden hamsters were randomly housed in separate cages (five hamsters per cage). The golden hamsters were synchronized for 3 weeks under 12-hour light/12-hour dark cycles (at a temperature of 24°C $\pm$ 1°C and humidity of 60% $\pm$ 10%) before the study. The hamsters' bedding, food, and water were sterilized. In the 12-hour light/12-hour dark cycles, the time was described as hours after light onset (HALO). For example, "0 HALO" was denoted as the time to turn on the light, whereas "12 HALO" was denoted as the time to turn off the light. On the last day of the third week, five hamsters were sacrificed by cervical dislocation at each time point, including 4 HALO, 8 HALO, 12 HALO, 16 HALO, 20 HALO, and 24 HALO during a 24-hour period, and normal left buccal mucosa was simultaneously obtained from the normal group. The remaining 60 hamsters were housed under a 12-hour light/12-hour dark cycle. The hamsters were maintained in a self-made box, and their

mouths were opened and painted with a 0.5% dimethylbenzanthracene acetone solution (Sigma, USA) with a No 5 flat brush on the left buccal mucosa every Monday, Wednesday, and Friday. On the last day of the sixth and 14th week after dimethylbenzanthracene painting, five hamsters were sacrificed by cervical dislocation at each time point, including 4 HALO, 8 HALO, 12 HALO, 16 HALO, 20 HALO, and 24 HALO during a 24-hour period; the left buccal diseased tissue was obtained simultaneously. The tissues obtained in the sixth week were included in the precancerous lesions group, whereas tissues obtained in the 14th week were included in the cancer group. Then, each tissue sample was divided into two parts. One portion was fixed in 4% paraformaldehyde, dehydrated, and embedded in paraffin blocks, whereas the other portion was snap-frozen in liquid nitrogen.

## Pathological examination using HE staining

We obtained 4  $\mu\text{m}$  thick sections from each paraffin block, and the hematoxylin and eosin (HE)-stained sections were prepared for observation under an optical microscope (Olympus BX51, Japan). Classification of histological pathology was accomplished according to the classification standard of WHO Collaborating Centre for Oral Precancerous Lesions.<sup>31</sup>

## Real-time RT-PCR

Real-time reverse transcriptase polymerase chain reaction (RT-PCR) was performed according to the manufacturer's protocol. First, the total RNA was isolated using RNAiso Plus (Takara, Japan). Second, the cDNA was synthesized from 10  $\mu\text{L}$  of total RNA using a Prime Script RT reagent kit (Takara). Finally, RT-PCR was performed on a real-time PCR machine (CFX96TM Real-Time PCR Detection System; BioRad, USA) in a 25  $\mu\text{L}$  final volume containing 12.5  $\mu\text{L}$  of 2 $\times$  SYBR Premix Ex TaqTM II (Takara), 8.5  $\mu\text{L}$  of RNase-free water, 1  $\mu\text{L}$  of 0.4  $\mu\text{mol/L}$  forward primer and reverse primer, and 2  $\mu\text{L}$  of 50 ng/ $\mu\text{L}$  cDNA. Specific primers for  $\beta$ -actin (a housekeeping gene that served as an internal control), *PER1*, *VEGF*, *KI67*, *C-MYC*, and *P53* were designed using Oligo 17.0 software. The primer sequences are listed in Table 1. The following RT-PCR cycling parameters were used: 1.5 minutes at 95°C followed by 40 rounds of 10 seconds at 95°C and 30 seconds at 60°C. The data were acquired as the threshold cycle (Ct) value. The relative mRNA expression values of *PER1*, *VEGF*, *KI67*, *C-MYC*, and *P53* were calculated using the  $2^{-\Delta\Delta C_t}$  method. Each sample was performed in triplicate to ensure the accuracy of the data.

**Table 1** Primers used for real-time PCR amplification of gene expression

Gene	Primer	Primer sequence (5' to 3')
PER1	Forward	AGCAACAGCCACGGTTCTC
	Reverse	CAGTCCACACAAGCCATCAC
VEGF	Forward	GACTTTGTTTGTCCAAGATCC
	Reverse	TCACATCTGCAAGTACGTTTCG
KI67	Forward	TGTATCCTTTGGTGGTCGTCTA
	Reverse	GCTGGAGTGTGAGTGGTGAG
C-MYC	Forward	GCCTACATCCTGTCCATCCA
	Reverse	TCACGCACCAGAGTTTCG
P53	Forward	ATGCCGAATACCTGGATGAC
	Reverse	GCGTGATGATGGTGAGGATA
$\beta$ -actin	Forward	GGCAGGCAAAGGTTACTCTG
	Reverse	TGGTGACAGGTGACAAGAT

**Abbreviation:** PCR, polymerase chain reaction.


## Statistical analysis

One-way analysis of variance and Student–Newman–Keuls test were used to analyze the differences in the mRNA expression of each gene among six time points and the differences in the mesor and amplitude of each gene in three groups with SPSS 17.0 statistical software (IBM Corporation, Armonk, NY, USA). Mean and standard deviation were used to describe the results, and a *P*-value of  $<0.05$  was considered statistically significant. The daily rhythm was assessed by a single cosine test using Time Series Analysis Cosinor 6.3 software (Expert Soft Technologie Laboratory of Applied Statistics and BioMedical Computing, Richelieu, France). A value of  $P<0.05$  indicated that the expression of the target gene showed a daily rhythm. A cosine-fitted curve was generated with SigmaPlot 10.0. The daily rhythm was characterized by the mesor, amplitude, and acrophase. The mesor is the mean value of all the statistics in a circadian fluctuation within a 24-hour period, the amplitude indicates the maximum degree above or below the mesor of the circadian fluctuation, and the acrophase represents the time point when the circadian fluctuation achieved the peak value.

## Results

### Pathological examination using HE staining

The pathological examination conducted using HE staining indicates that the 30 cases in the normal group exhibited normal buccal mucosa tissue. Among the 30 cases in the precancerous lesions group, 25 displayed moderate atypical hyperplasia, three exhibited mild atypical hyperplasia, and two exhibited severe atypical hyperplasia. The 30 cases in the cancer group exhibited squamous carcinoma tissue (Figure 1).



**Figure 1** The golden hamster buccal mucosa carcinogenesis model.

**Notes:** (A) Images of the golden hamster buccal mucosa from various carcinogenesis stages. (B) Pathological slice photos (HE staining,  $\times 400$ ) from various carcinogenesis stages.

**Abbreviation:** HE, hematoxylin and eosin.

## Daily rhythm variations of the clock gene PER1 mRNA in various stages of buccal mucosa carcinogenesis

Significant differences in PER1 mRNA expression were noted among the six different time points in the normal, precancerous lesions, and cancer groups (Table 2). The cosine analysis

revealed that PER1 exhibited daily rhythmic expression in the normal, precancerous lesions, and cancer groups (Table 2). The cosine-fitted curves are presented in Figure 2.

The daily rhythms demonstrated the following characteristics (Table 3). 1) Mesor and amplitude: The mesor and amplitude of PER1 mRNA expression in the precancerous lesions group


**Table 2** The mRNA expression of PER1, VEGF, KI67, C-MYC, and P53 in normal buccal mucosa, precancerous lesion, and cancer at six different time points in 24 hours (mean  $\pm$  SD,  $n=5$ )

Gene	Group	Time point						$P_1$ -value	$P_2$ -value	$P_3$ -value
		4 HALO	8 HALO	12 HALO	16 HALO	20 HALO	24 HALO			
PER1	N	0.82 $\pm$ 0.07	0.91 $\pm$ 0.08	1.84 $\pm$ 0.12	2.58 $\pm$ 0.08	1.83 $\pm$ 0.13	1.60 $\pm$ 0.05	0.000	0.000	0.000 <sup>a</sup>
	P	1.93 $\pm$ 0.17	1.80 $\pm$ 0.03	1.28 $\pm$ 0.21	0.98 $\pm$ 0.61	0.41 $\pm$ 0.07	0.81 $\pm$ 0.11	0.000	0.000	( $F=73.324$ )
	C	0.32 $\pm$ 0.11	0.91 $\pm$ 0.22	0.66 $\pm$ 0.22	1.39 $\pm$ 0.19	1.63 $\pm$ 0.17	0.38 $\pm$ 0.22	0.000	0.003	
VEGF	N	2.12 $\pm$ 0.14	2.53 $\pm$ 0.18	3.91 $\pm$ 0.16	1.39 $\pm$ 0.15	1.81 $\pm$ 0.14	1.79 $\pm$ 0.12	0.000	0.0062	0.000 <sup>a</sup>
	P	5.81 $\pm$ 0.36	4.87 $\pm$ 0.22	3.99 $\pm$ 0.12	1.42 $\pm$ 0.11	2.13 $\pm$ 0.21	2.84 $\pm$ 0.36	0.000	0.000	( $F=469.509$ )
	C	7.43 $\pm$ 0.89	6.34 $\pm$ 0.97	5.89 $\pm$ 0.54	4.67 $\pm$ 0.51	7.33 $\pm$ 0.63	9.51 $\pm$ 0.84	0.000	0.0002	
KI67	N	1.37 $\pm$ 0.37	0.54 $\pm$ 0.18	0.77 $\pm$ 0.19	0.37 $\pm$ 0.10	0.97 $\pm$ 0.27	1.10 $\pm$ 0.44	0.010	0.0115	0.000 <sup>a</sup>
	P	0.64 $\pm$ 0.25	1.36 $\pm$ 0.27	0.78 $\pm$ 0.43	2.03 $\pm$ 0.93	2.25 $\pm$ 0.56	0.99 $\pm$ 0.36	0.011	0.0246	( $F=31.144$ )
	C	2.62 $\pm$ 0.53	1.77 $\pm$ 0.15	1.71 $\pm$ 0.28	1.62 $\pm$ 0.31	2.59 $\pm$ 0.39	1.03 $\pm$ 0.25	0.001	0.9183	
C-MYC	N	2.37 $\pm$ 0.43	2.38 $\pm$ 0.38	2.14 $\pm$ 0.20	1.28 $\pm$ 0.45	0.96 $\pm$ 0.36	1.13 $\pm$ 0.31	0.001	0.000	0.000 <sup>b</sup>
	P	5.88 $\pm$ 0.65	4.30 $\pm$ 1.59	1.72 $\pm$ 0.86	2.33 $\pm$ 0.58	2.69 $\pm$ 0.68	3.45 $\pm$ 1.79	0.008	0.0026	( $F=19.542$ )
	C	5.41 $\pm$ 2.29	3.04 $\pm$ 0.30	2.66 $\pm$ 0.24	2.85 $\pm$ 0.29	5.70 $\pm$ 3.59	5.67 $\pm$ 0.56	0.052	0.0128	
P53	N	1.09 $\pm$ 0.26	1.86 $\pm$ 0.27	2.64 $\pm$ 0.22	3.82 $\pm$ 0.62	4.44 $\pm$ 1.40	6.20 $\pm$ 1.59	0.000	0.0032	0.000 <sup>a</sup>
	P	2.64 $\pm$ 0.63	2.06 $\pm$ 0.33	1.23 $\pm$ 0.28	2.15 $\pm$ 0.20	2.63 $\pm$ 0.62	3.28 $\pm$ 0.87	0.011	0.007	( $F=48.358$ )
	C	1.37 $\pm$ 0.21	0.84 $\pm$ 0.17	0.71 $\pm$ 0.10	1.63 $\pm$ 0.17	1.65 $\pm$ 0.42	1.40 $\pm$ 0.34	0.003	0.0021	

**Notes:**  $P<0.05$  represents that there is a significant difference.  $P_1$  represents the results of the differences in the expression of each gene at six time points in each group analyzed by one-way ANOVA.  $P_2$  represents the results of the daily rhythm of expressions of each gene in each group analyzed by single cosine test.  $P_3$  represents the results of the average differences in the expressions of each gene in three groups analyzed by one-way ANOVA. Superscripts "a" and "b" represent the results of the intergroup differences analyzed by Student–Newman–Keuls test after the one-way ANOVA, as follows: <sup>a</sup>There are mutual significant differences in three groups. <sup>b</sup>There are significant differences between P or C group and N group, but there is no significant difference between P and C groups.

**Abbreviations:** SD, standard deviation; n, numbers; HALO, hours after light onset; N, normal buccal mucosa; P, precancerous lesion; C, cancer; ANOVA, analysis of variance.





**Figure 2** Cosine-fitted curves of *PER1*, *VEGF*, *KI67*, *C-MYC*, and *P53* expression in normal buccal mucosa, precancerous lesions, and cancer tissues.

**Note:** The solid line in the figure represents the cosine-fitted curve, whereas the dotted line connects the mean values of the mRNA expression of each gene at six time points during a 24-hour period.

**Abbreviation:** HALO, hours after light onset.

and cancer group were significantly reduced compared with the normal group ( $P < 0.05$ ). The mesor and amplitude of *PER1* mRNA in the cancer group were remarkably reduced compared with the precancerous lesions group ( $P < 0.05$ ). 2) Acrophase: The acrophases expressed by *PER1* mRNA during various stages of carcinogenesis are presented in Table 3. In the cancer group, the acrophase of *PER1* mRNA was approximately identical to that observed in the normal group. In the precancerous

lesions group, the acrophase of *PER1* mRNA expression was 9.66 hours earlier than that in the normal group.

### Daily rhythm variations of mRNA of cancer-related genes during different stages of buccal mucosa carcinogenesis

Significant differences in *VEGF*, *KI67*, and *P53* mRNA were noted among the six daily time points in the normal,

**Table 3** The daily rhythm characteristics of PER1, VEGF, KI67, C-MYC, and P53 mRNA in normal buccal mucosa, precancerous lesion, and cancer

Gene	Cosinor parameters	Tissue			P-value
		N	P	C	
PER1	Mesor	1.60±0.11	1.20±0.06	0.88±0.04	0.000 <sup>a</sup> (F=48.137)
	Amplitude	0.81±0.01	0.71±0.01	0.54±0.04	0.000 <sup>a</sup> (F=92.659)
	Acrophase (HALO)	16.93±0.04	7.27±0.06	16.73±0.18	0.000 <sup>c</sup> (F=7,294.436)
VEGF	Mesor	2.26±0.21	3.51±0.33	6.86±0.59	0.000 <sup>a</sup> (F=71.351)
	Amplitude	0.82±0.01	2.06±0.23	1.91±0.06	0.000 <sup>b</sup> (F=71.938)
	Acrophase (HALO)	9.93±0.91	6.21±0.24	1.04±0.95	0.000 <sup>a</sup> (F=100.324)
KI67	Mesor	0.85±0.42	1.34±0.76	–	0.000 <sup>a</sup> (F=12.387)
	Amplitude	0.39±0.07	0.66±0.03	–	0.013 <sup>a</sup> (F=17.970)
	Acrophase (HALO)	1.65±0.71	19.93±0.74	–	0.003 <sup>a</sup> (F=42.501)
C-MYC	Mesor	1.71±0.10	3.39±0.03	4.22±0.11	0.000 <sup>a</sup> (F=543.026)
	Amplitude	0.83±0.06	1.74±0.01	1.87±0.04	0.000 <sup>b</sup> (F=574.424)
	Acrophase (HALO)	7.78±0.57	3.92±0.55	23.93±0.06	0.000 <sup>a</sup> (F=1,607.264)
P53	Mesor	3.34±0.19	2.33±0.26	1.27±0.21	0.000 <sup>a</sup> (F=82.040)
	Amplitude	1.92±0.13	0.86±0.08	0.45±0.07	0.000 <sup>a</sup> (F=109.770)
	Acrophase (HALO)	20.47±0.81	23.87±0.89	21.06±0.73	0.005 <sup>c</sup> (F=14.995)

**Notes:** The daily rhythm is described as mesor, amplitude, and acrophase.  $P < 0.05$  represents that there is a significant difference. P-value represents the results of differences in each cosinor parameter of each gene in three groups analyzed by one-way ANOVA. Superscripts "a", "b", and "c" represent the results of the intergroup differences analyzed by Student–Newman–Keuls test after the one-way ANOVA, as follows: <sup>a</sup>There are mutual significant differences in three groups. <sup>b</sup>There are significant differences between P or C group and N group, but there is no significant difference between P and C groups. <sup>c</sup>There are significant differences between P group and C or N group, but there is no significant difference between C and N groups.

**Abbreviations:** N, normal buccal mucosa; P, precancerous lesion; C, cancer; HALO, hours after light onset; ANOVA, analysis of variance.

precancerous lesions, and cancer groups, respectively ( $P < 0.05$ ). C-MYC mRNA exhibited significant differences at the six daily time points in the normal and precancerous lesions groups ( $P < 0.05$ ); however, no differences were noted in the cancer group ( $P > 0.05$ ), as shown in Table 2. The cosine analysis indicated that VEGF, P53, and C-MYC mRNA, respectively, exhibited daily rhythmic expression in the normal, precancerous lesions, and cancer groups ( $P < 0.05$ ). KI67 mRNA exhibited daily rhythmic expression in the normal and precancerous lesions groups ( $P < 0.05$ ), whereas this expression pattern was not observed in the cancer group ( $P > 0.05$ ) (Table 2 and Figure 2).

The daily rhythms exhibited the following characteristics (Table 3). 1) Mesor: The mesor of VEGF and C-MYC mRNA expression in the precancerous lesions and cancer groups, respectively, was significantly increased compared with the normal group ( $P < 0.05$ ). In the cancer group, the mesor of VEGF and C-MYC mRNA was remarkably increased compared with that in the precancerous lesions group ( $P < 0.05$ ). The mesor of KI67 mRNA in the precancerous lesions group was remarkably increased compared with the normal group ( $P < 0.05$ ). The mesor of P53 mRNA in the precancerous lesions group and cancer group significantly decreased compared with the normal group ( $P < 0.05$ ). In the cancer group, the mesor was significantly reduced compared with the precancerous lesions group ( $P < 0.05$ ).

2) Amplitude: The amplitudes of VEGF and C-MYC mRNA in the precancerous lesions and cancer groups, respectively, were significantly increased compared with the normal group ( $P < 0.05$ ). No remarkable difference was noted between the precancerous lesions and cancer groups ( $P > 0.05$ ). The amplitude of KI67 mRNA in the precancerous lesions group was remarkably increased compared with the normal group ( $P < 0.05$ ). The amplitude of P53 mRNA in the precancerous lesions and cancer groups was significantly reduced compared with the normal group ( $P < 0.05$ ), and the amplitude in the cancer group was remarkably decreased compared with the precancerous lesions group ( $P < 0.05$ ). 3) Acrophase: The acrophases of VEGF, KI67, C-MYC, and P53 mRNA in various carcinogenesis periods are presented in Table 3. The acrophases of VEGF mRNA in the precancerous lesions and cancer groups were 3.72 hours and 8.89 hours earlier, respectively, compared with the normal group. The acrophase of KI67 mRNA in the precancerous lesions group was delayed 18.28 hours compared with the normal group. The acrophases of P53 mRNA were similar in the normal and cancer groups. In the precancerous lesions group, the acrophase was 3.4 hours later than that observed in the normal group. The acrophases of C-MYC mRNA in the precancerous lesions and cancer groups were 3.86 hours earlier and 16.15 hours later, respectively, compared with the normal group.

## Discussion

Current studies have indicated that the clock gene *PER1* is a cancer suppressor gene, and its circadian variations exhibit a close relationship with the occurrence and development of cancers.<sup>8,15,27–29,32</sup> Bjarnason et al reported that *PER1* expression in human buccal mucosa cells exhibits a circadian rhythm.<sup>25</sup> Our previous research demonstrated decreased and circadian rhythmic *PER1* expression in oral squamous cell carcinomas.<sup>28,33</sup> This study reports that the expression of *PER1* in three stages of oral squamous cell carcinoma exhibited daily rhythms. The mesor and amplitude of *PER1* mRNA expression decreased remarkably as carcinogenesis progressed. These results demonstrate that the role of *PER1* in cancer inhibition was gradually reduced, probably resulting in malignant transformation. In terms of the acrophase, the timing of *PER1* mRNA expression peak values was similar in the normal buccal mucosa and carcinoma stages, and the patterns of *PER1* mRNA daily rhythmic expression were similar. In the precancerous lesions stage, the acrophase of *PER1* mRNA was 9.66 hours earlier than that observed in the normal buccal mucosa. The expression pattern of the daily rhythms of *PER1* mRNA in the normal buccal mucosa was in contrast to that observed in precancerous lesions. The mechanism of these variations requires further study.

Carcinogenesis is dependent on tumor angiogenesis, and *VEGF* plays a central role in tumor-induced angiogenesis.<sup>34,35</sup> Koyanagi et al reported that *VEGF* mRNA expression in transplanted sarcoma cells of mice exhibits circadian rhythms. In hepatic cells and plasma of mice with tumors, *VEGF* protein exhibits circadian rhythmic expression.<sup>36</sup> Our research found that *VEGF* mRNA expression exhibits daily rhythms in the three carcinogenesis stages in oral squamous cell carcinoma. The mesor and amplitude of *VEGF* mRNA expression were significantly elevated upon the development of cancer, thus promoting tumor angiogenesis and carcinogenesis. The acrophase of *VEGF* mRNA constantly progressed with the occurrence and development of cancer. The mechanism of such variations in carcinogenesis requires further exploration and serves as an important reference value for the administration of cancer chemotherapies using *VEGF* as a biomarker.

*P53* is a cancer suppressor gene, and increased *P53* expression inhibits tumor growth and induces cell apoptosis.<sup>37,38</sup> *KI67* and *C-MYC* are cell proliferation genes that promote cell proliferation via increases in their expression.<sup>37,39</sup> Our research demonstrated that *P53* and *C-MYC* mRNA exhibit daily rhythmic expression in three stages of buccal mucosa carcinogenesis. However, *KI67* mRNA exhibited daily

rhythmic expression in the normal and precancerous lesions groups, whereas *KI67* mRNA expression was dysregulated in the cancer group. During the occurrence and development of cancer, the mesor and amplitude of *P53* mRNA expression typically decreased gradually, indicating a weakened effect on cancer inhibition and apoptotic induction. *KI67* and *C-MYC* mRNA expression increased gradually, resulting in enhanced reinforcement of cell proliferation and leading to an imbalance between cell proliferation and apoptosis. Regarding the acrophase, *P53* mRNA expression was approximately identical in the normal and cancer groups; *P53* mRNA expression in the precancerous lesions group lagged by 3.4 hours compared with the normal group. In contrast, the acrophase of *KI67* mRNA in the precancerous lesions group was 18.28 hours later than the normal group. *C-MYC* mRNA expression was 3.86 hours earlier and 16.15 hours later in the precancerous lesions and cancer groups, respectively. The uncoordinated changes in the acrophases of the apoptosis gene *P53* and cell proliferation genes *KI67* and *C-MYC* probably further aggravate the imbalance between cell proliferation and apoptosis, thus promoting carcinogenesis.

## Conclusion

The specific mechanisms by which the clock gene *PER1* regulates and controls cancer-related genes, such as *VEGF*, *KI67*, *C-MYC*, and *P53*, and the characteristic changes in the circadian rhythms of these genes remain unclear; however, this study was the first to demonstrate the daily rhythmic expression patterns of the clock gene *PER1* and cancer-related genes, such as *VEGF*, *KI67*, *C-MYC*, and *P53*, in different stages of carcinogenesis. These results indicate that the mesors, amplitudes, and acrophases of the clock gene *PER1* and cancer-related genes *VEGF*, *KI67*, *C-MYC*, and *P53* undergo substantial changes upon the development of cancer. This study provides novel ideas and methods for the exploration of carcinogenic mechanisms and cancer treatments from the perspective of the relationship between circadian variations in gene expression and carcinogenesis.

## Acknowledgments

We thank Miss Ling Li for her help in preparing the manuscript and Mr Baisong Li for assisting in the statistical analysis of the data.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Zieker D, Jenne I, Koenigsrainer I, et al. Circadian expression of clock- and tumor suppressor genes in human oral mucosa. *Cell Physiol Biochem*. 2010;26(2):155–166.
- Borgs L, Beukelaers P, Vandenbosch R, Belachew S, Nguyen L, Malgrange B. Cell “circadian” cycle: new role for mammalian core clock genes. *Cell Cycle*. 2009;8(6):832–837.
- Bozek K, Relógio A, Kielbasa SM, et al. Regulation of clock-controlled genes in mammals. *PLoS One*. 2009;4(3):e4882.
- Wulff K, Porcheret K, Cussans E, Foster RG. Sleep and circadian rhythm disturbances: multiple genes and multiple phenotypes. *Curr Opin Genet Dev*. 2009;19(3):237–246.
- Oster H, Damerow S, Hut RA, Eichele G. Transcriptional profiling in the adrenal gland reveals circadian regulation of hormone biosynthesis genes and nucleosome assembly genes. *J Biol Rhythms*. 2006;21(5):350–361.
- Keller M, Mazuch J, Abraham U, et al. A circadian clock in macrophages controls inflammatory immune responses. *Proc Natl Acad Sci U S A*. 2009;106(50):21407–21412.
- Huang XL, Fu CJ, Bu RF. Role of circadian clocks in the development and therapeutics of cancer. *J Int Med Res*. 2011;39(6):2061–2066.
- Cao Q, Gery S, Dashti A, et al. A role for the clock gene *per1* in prostate cancer. *Cancer Res*. 2009;69(19):7619–7625.
- Bjarnason GA, Jordan RC, Sothorn RB. Circadian variation in the expression of cell-cycle proteins in human oral epithelium. *Am J Pathol*. 1999;154(2):613–622.
- Rana S, Mahmood S. Circadian rhythm and its role in malignancy. *J Circadian Rhythms*. 2010;8:3.
- Savvidis C, Koutsilieris M. Circadian rhythm disruption in cancer biology. *Mol Med*. 2012;18:1249–1260.
- Schibler U, Sassone-Corsi P. A web of circadian pacemakers. *Cell*. 2002;111(7):919–922.
- Fu L, Pelicano H, Liu J, Huang P, Lee C. The circadian gene *Period2* an important role in tumor suppression and DNA damage response in vivo. *Cell*. 2002;111:41–50.
- Schibler U, Ripperger J, Brown SA. Peripheral circadian oscillators in mammals: time and food. *J Biol Rhythms*. 2003;18(3):250–260.
- Soták M, Polidarová L, Ergang P, Sumová A, Pácha J. An association between clock genes and clock-controlled cell cycle genes in murine colorectal tumors. *Int J Cancer*. 2013;132(5):1032–1041.
- Yoo SH, Yamazaki S, Lowrey PL, et al. *PERIOD2*: luciferase real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci U S A*. 2004;101(15):5339–5346.
- Shearman LP, Sriram S, Weaver DR, et al. Interacting molecular loops in the mammalian circadian clock. *Science*. 2000;288(5468):1013–1019.
- Gekakis N, Staknis D, Nguyen HB, et al. Role of the *CLOCK* protein in the mammalian circadian mechanism. *Science*. 1988;280(5369):1564–1569.
- Kume K, Zylka MJ, Sriram S, et al. *mCRY1* and *mCRY2* are essential components of the negative limb of the circadian clock feedback loop. *Cell*. 1999;98(2):193–205.
- Sato TK, Yamada RG, Ukai H, et al. Feedback repression is required for mammalian circadian clock function. *Nat Genet*. 2006;38(3):312–319.
- Storch KF, Lipan O, Leykin I, et al. Extensive and divergent circadian gene expression in liver and heart. *Nature*. 2002;417(6884):78–83.
- Meyer V, Lerchl A. Evidence for species-specific clock gene expression patterns in hamster peripheral tissues. *Gene*. 2014;548(1):101–111.
- Rana S, Munawar M, Shahid A, et al. Deregulated expression of circadian clock and clock-controlled cell cycle genes in chronic lymphocytic leukemia. *Mol Biol Rep*. 2014;41(1):95–103.
- Fu L, Kettner NM. The circadian clock in cancer development and therapy. *Prog Mol Biol Transl Sci*. 2013;119:221–282.
- Bjarnason GA, Jordan RC, Wood PA, et al. Circadian expression of clock genes in human oral mucosa and skin: association with specific cell-cycle phases. *Am J Pathol*. 2001;158(5):1793–1801.
- Zheng L, Seon YJ, McHugh J, Papagerakis S, Papagerakis P. Clock genes show circadian rhythms in salivary glands. *J Dent Res*. 2012;91(8):783–788.
- Lengyel Z, Lovig C, Kommedal S, et al. Altered expression patterns of clock gene mRNAs and clock proteins in human skin tumors. *Tumor Biol*. 2013;34(2):811–819.
- Zhao N, Yang K, Yang G, et al. Aberrant expression of clock gene *period1* and its correlations with the growth, proliferation and metastasis of buccal squamous cell carcinoma. *PLoS One*. 2013;8(2):e55894.
- Gery S, Komatsu N, Baldjian L, Yu A, Koo D, Koeffler HP. The circadian gene *per1* plays an important role in cell growth and DNA damage control in human cancer cells. *Mol Cell*. 2006;22(3):375–382.
- Pluquet O, Dejeans N, Chevet E. Watching the clock: endoplasmic reticulum-mediated control of circadian rhythms in cancer. *Ann Med*. 2014;46(4):233–243.
- WHO Collaborating Centre for Oral Precancerous Lesions. Definition of leukoplakia and related lesions: an aid to studies on oral precancer. *Oral Surg Oral Med Oral Pathol*. 1978;46:518–539.
- Lee S, Donehower LA, Herron AJ, Moore DD, Fu L. Disrupting circadian homeostasis of sympathetic signaling promotes tumor development in mice. *PLoS One*. 2010;5(6):e10995.
- Chen R, Yang K, Zhao NB, et al. Abnormal expression of *PER1* circadian-clock gene in oral squamous cell carcinoma. *Onco Targets Ther*. 2012;5:403–407.
- Wang CY, Wen MS, Wang HW, et al. Increased vascular senescence and impaired endothelial progenitor cell function mediated by mutation of circadian gene *Per2*. *Circulation*. 2008;118(21):2166–2173.
- Wood PA, Du-Quito J, You S, Hrushesky WJ. Circadian clock coordinates cancer cell cycle progression, thymidylate synthase, and 5-fluorouracil therapeutic index. *Mol Cancer Ther*. 2006;5(8):2023–2033.
- Koyanagi S, Kuramoto Y, Nakagawa H, et al. A molecular mechanism regulating circadian expression of vascular endothelial growth factor in tumor cells. *Cancer Res*. 2003;63(21):7277–7283.
- Granda TG, Liu XH, Smaalend R, et al. Circadian regulation of cell cycle and apoptosis proteins in mouse bone marrow and tumor. *FASEB J*. 2005;19(2):304–306.
- Rengarajan T, Nandakumar N, Rajendran P, Haribabu L, Nishigaki I, Balasubramanian MP. D-pinitol promotes apoptosis in MCF-7 cells via induction of p53 and Bax and inhibition of Bcl-2 and NF-κB. *Asian Pac J Cancer Prev*. 2014;15(4):1757–1762.
- Scholzen T, Gerdes J. The Ki-67 protein: from the known and the unknown. *J Cell Physiol*. 2000;182(3):311–322.

### OncoTargets and Therapy

### Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.