Prevalence of presbyopia and spectacle correction coverage in a rural population of North West Nigeria

Murtala Muhammad Umar¹
Nasiru Muhammad²
Mahmoud B Alhassan³

¹Zamfara State Eye Care Programme, King Fahad IBN Abdul-Azeez Women and Children Hospital, Zamfara State, Nigeria; ²Ophthalmology Unit, Surgery Department, College of Health Sciences Usmanu Danfodiyo University, Sokoto, Nigeria;

³Department of Vitreoretina, National Eye Centre, Kaduna, Nigeria

Purpose: To determine the prevalence of presbyopia, and near vision spectacle coverage in a rural population of Northwestern Nigeria.

Study design: Cross sectional prevalence study.

Subjects and methods: Six hundred and fifty people of at least 40 years of age, in 13 clusters (50 per cluster) were examined using a multi-stage random sampling with probability proportional to size. The survey was conducted from April 7 to 28, 2012 at Bungudu Local Government Area of Zamfara State, Nigeria. Presbyopia was defined as the inability to read N8 at 40 cm. Presbyopic Spectacle Correction Coverage (PSCC) was calculated, and information on barriers to using near vision spectacles identified.

Results: The crude prevalence of presbyopia was 30.4%, 95% CI: (26.8%–34.1%). The prevalence was significantly higher in females ($P=0.0005$) and individuals with at least secondary education ($P=0.022$). The age specific prevalence of presbyopia was three times (63.5%) more among those aged 70 years and above, as compared to those within 40–49 years age group (19.3%). The met need was 0.2%, the unmet need 30.2%, and a PSCC of 0.7%. The major barriers reported as reasons for not obtaining near vision spectacles were unawareness and lack of felt need.

Conclusion: The prevalence of presbyopia in Bungudu is relatively low compared to other reports with major risk factors being increasing age, female sex and attainment of higher education. The presbyopic spectacles correction coverage is very low with high unmet need thus there is a need to create awareness, and provide affordable and accessible optical services in the affected population.

Keywords: presbyopia, spectacle coverage, Nigeria

Introduction

Presbyopia is an inability to focus at near distance as a result of loss of accommodation with age.¹

With age, presbyopia eventually affects everyone but is generally measured and diagnosed only when an individual becomes symptomatic and presents to an eye care provider with need for near-vision correction. In clinical practice, presbyopia is diagnosed by measurements of accommodative amplitude, near subjective refraction, and/or patient's reported symptoms.²

While new treatments are being developed for presbyopia, spectacles represent an effective, economic option for low- and middle-income countries.³ Use of contact and accommodative intraocular lenses have the advantage of making presbyopes spectacle independent. Lenticular and corneal surgical manipulations have been reported to alleviate presbyopia; these include multifocal and monovision corneal refractive surgical

Correspondence: Murtala Muhammad Umar
Zamfara State Eye Care Programme,
King Fahad IBN Abdul-Azeez Women
and Children Hospital, P.M.B 1129,
Gusau, Zamfara State, Nigeria
Email murtalau@gmail.com

procedures, or near-vision conductive keratoplasty, or a pinhole (PH) corneal inlay.⁴ These devices and surgical procedures are however not accessible to our study population.

It is now increasingly recognized that presbyopia is an aspect of refractive error that needs to be addressed. Good vision is important, even among populations who use it for tasks other than reading and writing.³

Holden et al⁵ estimated in 2005 that 1.04 billion people worldwide were presbyopic with an expected increase to 1.4 billion and 1.8 billion by the years 2020 and 2050, respectively. They deduced the prevalence in North America, Europe, and Japan to be 83% among individuals 45 years and above. Population-based studies in East Africa reported prevalence of 61.7%–85.4%,^{6,7} with low spectacle coverage in individuals aged 40 years and older.

Few population-based studies^{8–10} carried out in Nigeria reported presbyopia prevalence of 33%–53.5%; majority of reported studies being institution-based.^{11–14}

Zamfara state located in Northwestern Nigeria has had a VISION 2020: the Right to Sight-compliant eye care program since 2009 that provides eye care services including optical services. Although implemented through the primary health care structure, optical services are largely restricted to major towns where skilled workers are available to provide the service. The purpose of this study was to provide data on presbyopia and spectacle correction coverage and barriers that could serve as baseline data for stakeholders of the Zamfara state eye care program and nearby states. The data could be used for advocacy, planning, and resource allocation to achieve a more efficient and effective optical service delivery.

Subjects and methods

Study design

This was a population-based survey of people 40 years and above conducted from April 7 to 28, 2012.

Ethical considerations

Human Research and Ethics Committee of the National Eye Centre in Nigeria granted ethical approval. Bungudu Local Government Area (LGA) and Zamfara State Ministry of Health also gave approval for conducting the study.

Sample size determination

A minimum sample size of 646 people was calculated using the formula:

$$n = \frac{Z^2 pq}{d^2} \quad (1)$$

where n = required sample size, Z = standard normal deviation, p = expected prevalence, $q=(1-p)$, d = degree of accuracy and multiplied by the design effect.

$$(Z=1.96 \text{ (95\%)}, p=55\%^8, d=0.05 \text{ (5\%)}, \text{ design effect}=1.7) \quad (2)$$

Study population

Bungudu LGA is one of the 14 LGAs in Zamfara State with 2,356 km² land mass. The local government headquarters is about 20 km from Gusau, the state capital. It is located at 12°16'00"N 6°33'24"E. It has a population of 257,917 with 20% aged 40 years and above who are mainly agrarian Hausas, Fulanis, and a few other minority ethnic groups of Nigeria. The LGA has eleven administrative wards with one general hospital and 57 primary health centers. There is no eye clinic in any of the health facilities in the LGA. The nearest eye care facility is located 20 km away in the state capital. In 2011, the State Eye Care Programme trained ten community health extension workers as integrated eye care workers who are spread across the health facilities.

Inclusion criteria

1) Individuals 40 years and above. 2) Resident who has spent at least 6 months in the community continuously.

Exclusion criteria

1) Person(s) whose presenting distance visual acuity (VA) is less than 6/60 of Snellen chart and did not improve with PH. 2) Individuals with mental or other incapacitating illnesses whose vision cannot be tested.

Sampling technique

Using a cluster size of 50 people, 13 (650/50) clusters were selected using a multi-stage random sampling with probability proportional to size. In each cluster, sample selection was by "spin-the-bottle method" at the center of the cluster that was followed by random walk to select households for inclusion into the study. All eligible people in a household were included in the survey until the required sample size in a cluster was obtained.

Survey team and training

The survey team comprised of an ophthalmologist (principal investigator [PI]), an ophthalmic nurse (ON) and a research assistant (interviewer). A pilot survey was conducted in a non-selected community in the LGA. The VA measurement achieved an inter-observer agreement of more than 90%

between the ON and the PI. The purpose of the interview was masked to the research assistant (interviewer) to reduce bias and an agreement of more than 90% was obtained with the PI on how to administer the questionnaire. Areas of difficulty were re-discussed and further practiced to minimize errors in data collection. The International Center for Eye Health of the London School of Hygiene and Tropical Medicine validated the survey questionnaire as used in an earlier study in Nigeria.⁹

Data collection procedures

The trained interviewer obtained enumeration and demographic information comprising of age, sex, occupation, and educational level after obtaining consent. Historical events were used to estimate age where necessary.

The ON assessed the distance VA of subjects using the Snellen tumbling E chart at six meters in ambient outdoor illumination under shade. PH VA was done on all subjects who had VA less than 6/18 in either eye. Refractive error was defined as improvement of vision to 6/18 or better with PH in either eye.

The ophthalmologist then carried out objective and subjective distance refraction for subjects presenting with VA less than 6/18 who had an improvement of at least one Snellen acuity line with a PH in either eye.

Near vision was tested at 40 cm (wearing distance correction where necessary) using logMAR (logarithm of the minimum angle of resolution) near E chart under ambient indoor illumination. Correct identification of three out of five characters constituted success in reading a line. The end point of near vision testing was N8 optotypes. Presbyopia (functional) was defined in a subject who could not read the N8 optotype at 40 cm with the distance correction in place, if required. Under-corrected presbyopia is said to be present in a subject presenting with near vision spectacles but who fails to read N8.

Spectacles for near correction were provided at no cost to the presbyopes (uncorrected or under-corrected), subject to availability of the required power and willingness of participants to accept. This was done after completion of data collection in an entire cluster without prior knowledge of the participants to avoid bias. Subjects whose near spectacle prescription was not available or had other ocular pathologies were referred to the eye clinic for appropriate management.

Participants with presbyopia who do not present with near vision spectacle correction were asked to state reason(s) for not having a presbyopic spectacle correction.

Data management

Collected data were entered into a software program designed in SPSS for windows version 16.0 (SPSS Inc., Chicago, IL, USA) and analyzed with support from a statistician. The prevalence, confidence intervals and *P*-values (significant at the $P<0.05$ level), percentages of met need, unmet need, and presbyopia spectacles correction coverage (PSCC) were calculated. Analysis of variance (ANOVA) was used to determine the association between age, sex, and literacy with near visual impairment. “Met need” was calculated as the number of presbyopic subjects who are able to read N8 with presenting near vision spectacles out of the total people examined. “Unmet need” was the number of presbyopic people without near vision spectacles or unable to read N8 with spectacles (where available) out of the total people examined. PSCC was calculated using the formula:

$$PSCC = \frac{\text{Met need}}{\text{Met need} + \text{Unmet need}} \times 100. \quad (3)$$

Results

General characteristics

Of the 650 subjects enumerated, 635 were examined constituting a response rate of 97.7%. The mean age of participants was 53.59 years (range 40–97 years). As shown in Table 1, there was no significant age differences among people examined and those not examined (refused and absent), ($P=0.448$). The baseline characteristics of subjects examined are depicted in Table 2.

The number of males and females and their proportions within the age groups was significantly different ($P=0.000$). Majority (42.4%) of the sample was 40–49 years of age. Majority of the participants had Qur’anic education and only 4.1% attained tertiary education (Table 3).

Out of all the females examined only 3.2% (8/249) had formal education. Farming was the major occupation among the males (48.5%) while the majority of the females were housewives.

Prevalence of presbyopia

The crude prevalence of presbyopia was 30.4%, (95% confidence interval: 26.8%–34.1%). The mean dioptric add required to read N8 increased significantly with age ($P=0.000$).

The prevalence of presbyopia was two times higher in females ($P=0.000$). The prevalence of presbyopia is higher among hyperopes (100%) than myopes (64%) and emmetropes (39%) (Figure 1). There is a weak association of presbyopia

Table 1 Baseline characteristics of sample population with age distribution of presbyopia

Characteristics	Examined		Not examined n (%)	Total n (%)
	n (%)	n (%)		
Age group (years)	Presbyopic	Non-presbyopic		
40–49	52 (19.3)	217 (80.7)	8 (1.2)	277 (42.7)
50–59	50 (30.1)	116 (69.9)	5 (0.8)	171 (26.3)
60–69	51 (37.2)	86 (62.8)	2 (0.3)	139 (21.4)
≥70	40 (63.5)	23 (36.5)	0 (0.0)	63 (9.7)
Total	193 (30.4)	442 (69.6)		650 (100)
Sex				
Male	143 (37)	243 (63)	2 (0.3)	388 (59.7)
Female	121 (49)	128 (51)	13 (2.0)	262 (40.3)
Total				650 (100)

with educational status as people with at least secondary education were more likely to be presbyopic ($P=0.022$).

PSCC, met, and unmet needs

Only six people with presbyopia presented with near vision correction spectacles, and five could not read N8 at 40 cm with their current spectacles. The met need was 0.2% ($1/635 \times 100$), unmet need of 30.2% ($192/635 \times 100$) and a PSCC of 0.7% (met need/met need + unmet need). There was no statistically significant difference among participants in terms of their age group, sex, literacy level, and occupation in use of near vision spectacle correction. The near vision spectacles were

obtained by three participants from the roadside spectacle vendors while the rest were from an eye clinic.

Barriers to use of near vision spectacles

The major barriers to obtaining near vision spectacles were “not aware of problems”, “no felt need”, “not a priority”, and “lack of money” (Table 4).

Women significantly reported lack of awareness as a barrier to obtaining near vision spectacles compared with men ($P=0.002$). Other demographic characteristics such as age group, literacy level, and occupation of the participants did not show any statistically significant difference

Table 2 Baseline characteristics of study subjects

Characteristics	Male N=386 (%)	Female N=249 (%)	Total N=635 (%)	P-value
Age group (years)				
40–49	148 (23.3)	121 (19.1)	269 (42.4)	0.011*
50–59	104 (26.9)	62 (9.8)	166 (26.2)	0.567
60–69	86 (22.3)	51 (8.0)	137 (21.5)	0.591
≥70	48 (7.5)	15 (2.4)	63 (9.9)	0.008*
Total			635 (100)	
Literacy level				
None	74 (11.6)	80 (12.6)	154 (24.2)	0.000*
Qur'anic	244 (38.4)	161 (25.4)	405 (63.8)	0.711
Primary	30 (4.7)	5 (0.8)	35 (5.5)	0.002*
Secondary	12 (1.9)	3 (0.5)	15 (2.4)	0.123
Tertiary	26 (4.1)	0 (0.0)	26 (4.1)	0.000*
Total			635 (100)	
Occupation				
Unemployed	1 (0.2)	1 (0.2)	2 (0.4)	1.000
House wife	0 (0.0)	222 (34.9)	222 (34.9)	
Farming	304 (47.9)	4 (0.6)	308 (48.5)	0.000*
Other manual work	15 (2.3)	1 (0.2)	16 (2.5)	0.013*
Skilled (self-employed)	11 (1.7)	0 (0.0)	11 (1.7)	0.0045*
Civil servant/public servant	46 (7.2)	0 (0.0)	46 (7.2)	0.000*
Retired civil servant/public servant	4 (0.6)	0 (0.0)	4 (0.6)	0.004
Others	5 (0.9)	21 (3.3)	26 (4.2)	0.000*
Total			635 (100)	

Note: *Statistically significant.

Table 3 Distribution of presbyopia by literacy level

Literacy level	Presbyopes N (%)	Non-presbyopes N (%)
None	51 (33.1)	103 (66.9)
Qura'nic	130 (32.1)	275 (67.9)
Primary	4 (11.4)	31 (88.6)
Secondary	5 (33.3)	10 (66.7)
Tertiary	3 (11.5)	23 (88.5)
Total	193 (30.4)	442 (69.6)

in the barriers reported. Of the six participants who have had near vision spectacles, four bought it for less than N2000.00 (US\$12) while two bought theirs for between N2,000.00 (US\$12) and N3,999.00 (US\$24). All of them said they could pay a maximum of the same amount for replacement.

Discussion

This study had a good response rate that was achieved through adequate community mobilization. The results of this study could be a reflection of presbyopia in the district and possibly nearby population. The results obtained in this study will help in the ongoing implementation of Zamfara State Eye Care Programme which hitherto had no baseline data on presbyopia in the state.

The mean age of participants in this study is similar to a study conducted in Abuja, Nigeria.⁹ Out of the 15 people who were absent during examination or refused, 13 (86%) were females. The refusal rate was higher in women because their male spouses did not give consent for examination. The literacy level of participants was low which is at variance with the Abuja and Southeastern Nigeria studies.^{8,9} Only 12% of the people examined had at least secondary education which was far lower than the average of 41.4%¹⁶ for the Zamfara state central senatorial district to which LGA study belongs.

The prevalence of 30.4% obtained in this study translates to a magnitude of between 13,824 and 17,590 people 40 years and above with presbyopia in Bungudu LGA. This is similar to that reported from studies in Southeast Nigeria (33%)⁸ and Timor-Leste (32.3%).¹⁵ The Timor-Leste study was similar to ours as they were both conducted in people 40 years and above unlike the Nigerian study that included younger people. Studies in Nigeria⁸ and other African countries^{6,7} reported prevalence higher than 50%. The variation may be due to significantly lower number of female participants who are known to have higher prevalence of presbyopia.^{6,7,9} A male female ratio of 1:1.7 was obtained in this study suggesting the need to give priority to females in eye care

Figure 1 The distribution of presbyopia based on refractive status.

Table 4 Barriers to use of near vision spectacles

Barriers	Yes N (%)	No N (%)	Total N (%)
Not aware of problems	148 (76.7)	45 (23.3)	193 (100)
No felt need	135 (69.9)	58 (30.1)	193 (100)
Not a priority	11 (5.7)	182 (94.3)	193 (100)
Lack of money	7 (3.6)	186 (96.4)	193 (100)
Do not know where to go	2 (1.0)	191 (99.0)	193 (100)
Services are too far away	1 (0.5)	192 (99.5)	193 (100)

services aimed at improving near vision. The low literacy level might have contributed to low prevalence, as higher prevalence has been associated with high literacy level.^{7,9} Majority of the people who attained secondary education in this study are presbyopic which may be related with higher visual demand for near tasks such as reading.

The mean dioptric power required to read N8 at 40 cm increases with age among participants in this study, which corroborates results from several other studies.^{6,7,15} This is due to decline in amplitude of accommodation with increasing age.^{1,17} All participants with hypermetropia were presbyopic compared to those with myopia and emmetropia. Hypermetropia is known to be associated with early onset presbyopia. However, residual hypermetropia not fully corrected may induce pseudo-presbyopia.

PSCC, met, and unmet needs

The PSCC, met and unmet needs obtained in this study are very low compared to results from similar studies in Nigeria⁹ and other developing countries.^{6,15} The Nigerian⁹ study reported a PSCC, met need and unmet need of 21%, 11.2%, and 42.1%, respectively. The low literacy level in this study population may have contributed to the low PSCC of 0.7%. A global survey showed that educated individuals are more likely to have eye examinations in their lifetime.¹⁸ An Indian study reported that the odds of seeking ophthalmic treatment were significantly higher among literate individuals.¹⁹ Only one (16.7%) out of six participants who presented with near vision spectacles read N8 at 40 cm, thus giving 83% under-corrected presbyopia. The absence of an eye clinic in the LGA and unfelt need for near vision spectacles might have accounted for this. Sex, literacy level, and occupation did not influence ownership of near vision spectacles in our study. There have been reports of high PSCC among males¹⁵ and people with higher education.⁸ Majority of the participants in this study read the Holy Qur'an, usually written in large Arabic script. This may delay the need for reading spectacles. Health education training and integration of nurses and community health

extension workers as refractionists, primary eye care workers, and redeploying them to rural areas will increase access to optical services thereby raising the PSCC. Incorporation of refraction services into the cataract outreach program may also enhance presbyopic spectacle correction uptake especially among women and the elderly who have high prevalence of presbyopia.

The major reasons given by participants for not owning near vision spectacles were “lack of awareness” and, “unfelt need”, lack of priority which compares with findings in studies from resource poor nations.^{7,8,20} This similarity may be due to common prevailing socio-economic conditions in the developing countries. None of the six participants with presbyopia who presented with spectacles interviewed in this study was willing to pay more than N4,000.00 (US\$24) for near vision spectacles. Provision of affordable and accessible optical services may enhance spectacle utilization in Bungudu LGA.

Of the six people who use spectacles in this study, three purchased it from roadside vendors. Ayanniyi et al²¹ reported that 16 (7%) out of 214 spectacle wearers interviewed in six health centers in Ilorin and Ekiti states obtained them from roadside vendors. This attitude should be discouraged through health education, because incorrectly prescribed glasses may act as a barrier to continuous use of spectacles if discomfort and poor vision persist. There is the need for provision of an affordable and accessible optical service to optimize spectacle correction uptake and improve quality of life. However, the fact that lack of felt need was the commonest barrier reported in this study means unless the people are health educated on the benefit of using near spectacles, utilization of the latter may remain low.

The onset of presbyopia influences near work habits of individuals and as such no uniform method can accurately detect it among different people. In this study the inability to read N8 at 40 cm with logMAR near chart may have under-diagnosed participants with presbyopia whose habitual near work distance is less than 40 cm. There is therefore a need for the stakeholders to develop short- and long-term plans to provide optical services to the population. The short-term strategy should create demand using radio and television jingles and also utilize available ONs in the state capital as refractionists diagnosing and dispensing spectacles for simple refractive errors including presbyopia using outreach service delivery strategy; and a long-term plan should be the establishment of an eye clinic in the local government area providing regular comprehensive eye care services including optical services.

Conclusion

The PSCC is very low in the study district and impliedly in Zamfara state, Nigeria with high unmet need.

Acknowledgment

We thank the management of National Eye Centre, Kaduna, Nigeria for partial funding, members of the survey team for the sacrifices during the conduct of the survey, and Prof CO Bekibele for his immense contribution and support during the conduct of this work that was submitted to the National Postgraduate Medical College of Nigeria as dissertation for the award of the Fellowship of the College.

Disclosure

The authors have no conflicts of interest to disclose.

References

1. Glasser A and Kaufman PL. Accommodation and presbyopia. In: Kaufman PL and Alm A, editors. *Adler's physiology of the Eye, clinical application*. 10th ed. Philadelphia: Mosby; 2003:197–233.
2. Hickenbotham A, Roorda A, Steinmaus C, Glasser A. Meta-Analysis of Sex Differences in Presbyopia. *Invest Ophthalmol Vis Sci*. 2012;53(6):3215–3220.
3. Patel I, West SK. Presbyopia: prevalence, impact and interventions. *Community Eye Health*. 2007;20(63):40–41.
4. Baikoff G. Surgical treatment of presbyopia: scleral, corneal, and lenticular. *Curr Opin Ophthalmol*. 2004;15(4):365–369.
5. Holden BA, Fricke TR, Ho SM, et al. Global Vision Impairment due to Uncorrected Presbyopia. *Arch Ophthalmol*. 2008;126(12):1731–1739.
6. Sherwin JC, Keffe JE, Kuper H, Islam FM, Muller A, Mathenge W. Functional presbyopia in a rural Kenyan population: the unmet presbyopic need. *Clin Experiment Ophthalmol*. 2008;36(3):245–251.
7. Burke AG, Patel I, Beatriz M, et al. Population-based study of presbyopia in rural Tanzania. *Ophthalmology*. 2006;113(5):723–727.
8. Nwosu SN. Ocular problems of young adults in rural Nigeria. *Int Ophthalmol*. 1998;22(5):259–263.
9. Chiroma MR. *Prevalence of presbyopia and the impact of uncorrected presbyopia on the quality of life in rural Gwagwalada, Abuja, Nigeria* [dissertation]. International Center for Eye Health (ICEH): London School of Hygiene and Tropical Medicine (LSHTM); 2008.
10. Kio FE, Ostia-Emina M. Pattern of refractive anomalies in Warri metropolis, Delta State of Nigeria. *Journal of the Nigerian Optometric Association*. 2003;10:30–38.
11. Ashaye AO, Asuzu MC. Ocular findings seen among the staff of an institution in Lagos Nigeria. *West Afr J Med*. 2005;24(2):96–99.
12. Bagaiya T, Pam V. Refractive error in Kaduna, Nigeria. *The Nigerian Journal of Surgical Research*. 2003;5(3–4):106–109.
13. Amadi AN, Nwankwo BO, Ibe AI, et al. Common ocular problems in Aba metropolis of Abia State, Eastern Nigeria. *Pakistan Journal of Social Sciences*. 2009;6(1):32–33.
14. Ekwonchi EE, Ezegwui I, Umeh R. Pattern of Eye disorders among inmates of a Nigerian Prison. *Orient Journal of Medicine*. 2009;2:(1–4). Available from: www.ajol.info/journal/home.
15. Ramke J, Toit RD, Palagyi A, Brian G, Naduvilath T. Correction of refractive error and presbyopia in Timor-Leste. *Br J Ophthalmol*. 2007;91(7):860–866.
16. National Bureau of Statistics Nigeria [homepage on the Internet]. Report of the National Literacy Survey 2010. Available from: nigerianstat.gov.ng/pages/download/4. Accessed March 10, 2013.
17. Elkington AR, Frank HJ, Greaney MJ, editors. Presbyopia. In: Elkington AR, Frank HJ, Greaney MJ, editors. *Clinical Optics*. 3rd ed. London: Wiley-Blackwell; 1999:141–151.
18. Vela C, Samson E, Zunzunegui MV, Haddad S, Aubin MJ, Freema EE. Eye care utilization by older adults in low, middle, and high income countries. *BMC Ophthalmol*. 2012;12:5.
19. Kovai V, Krishnaiah S, Shamanna BR, Thomas R, Rao GN. Barriers to accessing eye care services among visually impaired populations in rural Andhra Pradesh, South India. *Indian J Ophthalmol*. 2007;55(5):365–371.
20. Laviers H. The prevalence of presbyopia and feasibility of community distribution of near spectacles in adults in Zanzibar, East Africa. *J Community Eye Health*. 2007;20(64):73.
21. Ayanniyi AA, Fadamiro CO, Adeyemi, Folorunso FN, Uzukwu SC. Common refractive errors among the Ekitis of southwestern. *Journal of Medicine and Medical Sciences*. 2010;1(9):401–416.

Clinical Ophthalmology

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on

Submit your manuscript here: <http://www.dovepress.com/clinical-ophthalmology-journal>

Dovepress

PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.