Combined 2-deoxy glucose and metformin improves therapeutic efficacy of sodium-iodide symporter-mediated targeted radioiodine therapy in breast cancer cells

Sushmita Chatterjee
Nirmal Thaker
Abhijit De

Molecular Functional Imaging
Laboratory, Advanced Centre for
Treatment, Research and Education
in Cancer, Tata Memorial Centre,
Kharghar, Navi Mumbai, India

Abstract: Radiosensitization using either metformin or 2-deoxy-D-glucose (2-DG) in various cancer cells has been reported. The present study reveals novel information on combining these drugs to enhance radiosensitization effect in breast cancer (BC) cells. Responses to low-dose Cobalt60 radiation, as well as a newly emerged radioiodine therapy target for BC, that is, sodium-iodide symporter (NIS or SLC5A5) protein, are tested. As therapeutic potential of NIS in BC is often limited due to low uptake and fast efflux rate of iodine, the scope of these two radiosensitizers to further improve NIS-mediated ^{131}I therapeutic efficacy is explored. Two BC cell lines, MCF-7, and MDA MB231 are tested to optimize minimal drug doses required for radiosensitization. A combination of 2 mM metformin and 20 mM 2-DG with 2 grey (Gy) Cobalt60 radiation shows significant radiosensitization effect ($P=0.0002$). In cells treated with the combination therapy, increased $\gamma\text{H2A.X}$ foci formation was noted. Further, MCF-7 BC cells overexpressing NIS (MCF-7 NIS) was established, and using the optimized drug concentrations, significant radiosensitization ($P=0.0019$) by $50\ \mu\text{Ci }^{131}\text{I}$ usage was found to be the case as well. Apoptosis data corroborates with the result of clonogenic assay showing significant increase in apoptotic population upon dual drug-mediated radiosensitization. In case of metformin treatment, lowered adenosine triphosphate (ATP) content of the cell has been observed. The encouraging radiosensitization effect observed using combined 2-DG and metformin may aid in reducing Cobalt60 radiation exposure or for targeted radioiodine therapy in BC cells with NIS expression. This study indicates high potential of this drug combination in sensitizing BC cells for NIS-mediated-targeted radioiodine therapy, which otherwise may have lacked efficacy.

Keywords: sodium-iodide symporter, 2-DG, ^{131}I , radiosensitizers

Introduction

Breast cancer (BC) is the most common form of cancer in female population all over the world. Major treatment strategy for BC includes surgery, chemotherapy, hormonal therapy, and radiation therapy. Radiation therapy usually extends patients' survival; however, it has several side effects. Thus, reducing the dose of radiation to effectively kill the cancer cells with the aid of radiosensitizers has long been of great research interest. In addition, efficacious targeted radiotherapy option for various subtypes of BC is yet to develop. One such emerging strategy under active research investigation is the endogenous aberrant expression of sodium-iodide symporter (NIS or SLC5A5) protein, which can serve the purpose of the targeted radioiodine therapy. Human NIS is an integral membrane protein that is involved in active uptake of iodide ions inside the

Correspondence: Abhijit De
KS325, Molecular Functional Imaging
Laboratory, Advanced Centre for
Treatment, Research and Education in
Cancer, Tata Memorial Centre, Sector
22, Kharghar, Navi Mumbai 410210, India
Tel +91 22 2740 5038
Fax +91 22 2740 5085
Email ade@actrec.gov.in

thyroid cells, which has a crucial function in delivering iodine content required for the synthesis of thyroid hormones T3 and T4. NIS overexpression also serves the basis for the targeted radioiodine therapy of patients with thyroid cancer since past 60 years.^{1,2} Study of NIS in BC was pioneered by Tazebay et al, showing NIS-positive expression in 80% patients with BC, whereas normal breast tissue samples examined from the patients who underwent reductive mamoplasty showed negative expression.³ Following this study, many reports across the world confirmed high NIS expression in BC by various molecular analysis methods.⁴⁻⁹ In our own study, NIS expression was observed in 70% BC cases, where its expression was predominantly intracellular except in a few cases where NIS staining at the cell surface was also observed.⁹

Translational-medicine-based studies have also been carried out for testing the efficacy of NIS-based therapy and diagnosis of malignant breast tissue. Moon et al reported significant $^{99m}\text{TcO}_4^-$ uptake in four out of 25 patients with BC by scintigraphic scanning method.¹⁰ In a very important study, Wapnir et al detected radioiodide uptake in metastatic BC by scintigraphic analysis using radioiodides (^{123}I and ^{131}I) or $^{99m}\text{TcO}_4^-$.⁷ In this study, iodine uptake was noted in 25% of NIS-positive metastatic breast tissues (two out of eight). In addition, this study also showed significant suppression of thyroid radioiodine uptake by administration of agents like thyroxine and methimazole in trial patients. In another study, NIS expression was observed in 15 out of 23 patients with triple-negative BC, and membrane expression was observed in four patients with BC.⁴ Further, radioiodide uptake was also demonstrated in tumor tissue by ^{123}I scintigraphy in a patient with large primary BC; however, it was reported that had the isotope been ^{131}I , the cumulative concentration of iodine in the tumor may not deliver sufficient therapeutic benefits.⁴

Overall, these reports together suggest that NIS expression is high in BC as compared with normal breast tissue, but its expression is strong in a very small subset (<25%) of positive cases. This discrepancy widens the scope of multiple studies that have focused on modulating the expression level of NIS with agents such as retinoic acids and histone deacetylases before ^{131}I therapy.¹¹⁻¹³ Another strategy that is still unexplored is to enhance the ^{131}I treatment efficacy itself by using radiosensitizer drugs. Use of radiosensitizers may enhance cytotoxicity of ^{131}I therapy in case of a very low iodine uptake when NIS expression is low. At the same time, with the use of radiosensitizers, a lower concentration of ^{131}I can be used when NIS expression is high, which may benefit by minimizing the side effects of radiation in other NIS-expressing tissues, where entry of radioiodine cannot

be avoided. Further, without the presence of a defined iodide organification mechanism in BC cells, radiosensitizers may enhance radio-ablative effect even during the short presence of ^{131}I inside these cells. Since radiation therapy is known to show cytotoxicity via free radical-mediated oxidative damage, it has been proposed that combining radiation therapy with agents that cause oxidative stress-induced injury may sensitize cells to the cytotoxic effect of radiation.^{14,15} Thus, 2-deoxy-D-glucose (2-DG) shows good radiosensitization effect by disrupting thiol metabolism and increasing oxidative stress.¹⁶⁻¹⁹ Another drug used in this study is metformin (a bi-guanide drug and activator of AMP-activated protein kinase [AMPK]), which has been reported as a radiosensitizer for BC cells and showed eradication of radioresistant cancer stem cells by activating AMPK and suppressing mTOR.²⁰

To date, this is the first study to show the combination of 2-DG and metformin as radiosensitizers for the enhancement of Cobalt60 or ^{131}I radio-ablative effect on the cancer cells. We focus on combining the two radiosensitizer drugs against the two different radiation exposure methods and measuring the benefit using BC cells. We show that a combination of metformin and 2-DG along with Cobalt60 radiation or NIS-mediated ^{131}I radiation shows significantly enhanced cytotoxicity. To the best of our knowledge, this is also the first report where radiosensitization effect alone has been shown to enhance NIS-mediated ^{131}I therapy efficacy in BC cell model.

Materials and methods

Materials

MCF-7 and MDA MB231 BC cell lines were from ATCC (Manassas, VA, USA). RPMI1640 (Cat # 31800-022) and L15 (41300021) media, fetal bovine serum (10082-147), Trypsin-ethylenediaminetetraacetic acid (EDTA) (25300-062), lipofectamine 2000 (11668-027), and Hank's Balanced Salt Solution (HBSS, 24020-117) were obtained from Thermo Fisher Scientific (Waltham, MA, USA). pttB-CAG plasmid vector was obtained from Addgene (Cambridge, MA, USA). 2-DG (D8375), metformin (PHR 1084), propidium iodide (PI, P4170), 4',6-diamidino-2-phenylindole (DAPI, D8417), and MTT (M5655) were obtained from Sigma-Aldrich Co (St Louis, MO, USA). Monoclonal human NIS antibody (ab17795) and annexin V FITC (fluorescein isothiocyanate) apoptosis detection kit (ab14085) were obtained from Abcam (Cambridge, UK). H2A.X monoclonal antibody (MA5-15130) and anti-mouse (35512) and anti-rabbit Dylite633 (35562) antibody were obtained from Thermo Fisher Scientific. Enliten ATP assay kit (FF2000) was from Promega

Corporation (Fitchburg, WI, USA). The authors advise no ethical approval was required for this study, as they did not use any human tissue samples, and the complete study was performed using well established breast cancer cell lines.

MTT assay

A total of 5,000 cells/well were seeded in 96-well plate. Cells were allowed to adhere overnight before they were treated with mentioned drug doses of metformin or 2-DG. After 48 hours of treatment, cell viability was measured by MTT assay, and data are represented as percentage cell viability as compared with the untreated cells.

Preparation of NIS overexpression plasmid

NIS gene was inserted immediately downstream of chicken β -actin promoter in pcDNA3.1+ attB-CAG-FTT (Fluc2-tdtomato-thymidine kinase) vector system. Briefly, NIS gene was amplified by polymerase chain reaction using primers containing Nhe-I and Not-I restriction sites. The pcDNA3.1+ attB vector was digested with Nhe-I and Not-I, which released the FTT reporter from vector backbone. Further, NIS gene and attB vector backbone were ligated. Positive colonies were screened for NIS gene insertion by enzyme digestions and sequencing.

Preparation of NIS expressing BC cell line

MCF-7 BC cells were co-transfected by lipofectamine 2000 with pcDNA 3.1+ attB-CAG-NIS vector and integrase vector. We used integrase-based transfection method, where the plasmid vector contained bacterial attachment site (attB) upstream to chicken β -actin promoter.²¹ As the plasmid contained neomycin gene, positive clones were selected using 500 μ g G418. The isolated cell colonies were expanded and checked for NIS expression by immunofluorescence assay and functionally validated for iodine uptake. Cells obtained showed different levels of NIS expression and thus different levels of iodine uptake. The single cell colony (MCF-7 NIS) showing high expression of NIS protein, as well as good iodine uptake, was used for radiosensitization experiments.

Iodine uptake assay

A total of 50,000 MCF-7 NIS cells were seeded in 24-well plate in duplicates and incubated with 2 μ Ci 125 I/mL for 1 hour in uptake buffer (10 mM, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid [HEPES] in HBSS). At the end of 1 hour, medium was removed and cells were washed

once with cold uptake buffer, trypsinized, and collected. The iodine uptake in the collected cells is measured by beta counter. For showing NIS-specific iodine uptake, the sodium transporter was blocked by addition of 30 μ M KClO₄ along with 125 I.

Iodine efflux assay

A total of 50,000 cells were seeded in 24-well plate in duplicates and incubated with 125 I for 1 hour. At the end of 1 hour, 125 I-containing medium was removed and replaced with plain medium without 125 I. Cells were collected at the mentioned time points and remaining 125 I activity was measured by beta counter.

Clonogenic assay

In this assay, 2×10^6 cells were seeded in 25 mm flask and incubated with 50 μ Ci 131 I/mL in HBSS for 5–6 hours. At the end of the time point, 131 I-containing medium was removed, and cells were washed with cold HBSS. Cells were trypsinized, and 1,000 cells were seeded in six-well plates in triplicates. Medium was changed at regular intervals, and after 14 days, cells were fixed with 10% formaldehyde and stained with 4% crystal violet in methanol. Stained colonies were counted under stereomicroscope, and survival fraction was calculated according to the protocol mentioned in a previous publication.²² All the drug treatments were given 12 hours before the 131 I treatment/radiation therapy and were replaced by plain medium after 12 hours of 131 I treatment, and assay was performed after 24 hours of treatment.

Immunofluorescence for NIS and γ H2A.X foci assay

Cells were seeded on cover slip in 35 mm plate, and after drug/radiation/ 131 I treatment, they were washed with 1 $\times$ phosphate buffered saline (PBS) for three times and fixed with 4% para formaldehyde in PBS at 37°C. An additional permeabilization step with 0.2% TritonX 100 in 4% para formaldehyde was carried out for foci staining. Non-specific sites were blocked using 2% bovine serum albumin in PBS. Cells were then incubated with γ H2A.X or NIS antibody in 1:400/1:50 dilutions, respectively, for overnight at 4°C in a humid chamber. After incubation with primary antibody, cells were washed with 1 $\times$ PBS and incubated with either anti-rabbit Dylight633 or anti-mouse Dylight633 secondary antibody, respectively (1:200) for 1 hour at room temperature in dark humid chamber. Cells were washed with 1 $\times$ PBS, and nucleus was stained with DAPI. Cells were mounted

with Vecta Shield, and images were captured under confocal microscope (LSM510, Zeiss). Foci assay was performed at the immediate end of radiation/ ^{131}I treatment. For quantification of the foci, intensity of staining in red channel was measured by ImageJ software. For each treatment, 50 cells were analyzed, and average fluorescence intensity was plotted.

Cell cycle analysis and apoptosis assay

At the mentioned time points, 1×10^6 cells were trypsinized, collected, and washed with $1 \times \text{PBS}$. For cell cycle analysis, cells were fixed with 70% ethanol and incubated overnight at -20°C . Next day, cells were pelleted and washed with $1 \times \text{PBS}$. Washed cells were treated with $1 \mu\text{g/mL}$ RNase A for 10 minutes at 37°C . Then, $300 \mu\text{L}$ of $50 \mu\text{g/mL}$ propidium iodide was added in each tube, incubated for 30 minutes at 37°C , and acquired in FACSCaliburTM.

For apoptotic cell count, annexin V apoptosis detection kit was used, and the protocol mentioned in the kit was followed.

ATP measurement

Total cellular ATP was determined using manufactured recommended protocol of luciferase-based ATP detection

kit. Total ATP was measured after 24 hours of radiation or ^{131}I therapy.

Statistics

Data are presented as mean $\pm$ standard error. Student's *t*-test was used to judge the significance of difference. All tests were two-sided, and $P < 0.05$ was considered as significant. Graph pad Prizm software was used for all the statistical analysis.

Results

A combination of 2-DG and metformin enhances tumor cell ablation upon Cobalt60 radiation treatment

To determine the cytotoxic effect of 2-DG and metformin, BC cell lines were first exposed to various concentrations of these two drugs ranging between 0.1 mM and 100 mM, and viable cell count was measured by MTT assay (Figure S1). Further, based on the MTT data, 10–50 mM dose of 2-DG and 2–10 mM dose of metformin were combined with 1 or 2 grey (Gy) Cobalt60 radiation to determine the minimal optimum drug dose required for

Figure 1 Effect of 2-DG, metformin, and their combination with Cobalt60 radiation on MCF-7 cells.

Notes: (A and B) MCF-7 cells were treated with different dose of 2-DG or metformin with or without radiation exposure for 24 hours, and clonogenic assay was performed. (C) Combination of 20 mM 2-DG and 2 mM metformin with 2 Gy radiation dose showing further decrease in survival fraction. $*P < 0.05$; $**P < 0.01$. (D) Immunofluorescence analysis of $\gamma\text{H2A.X}$ foci showing increased expression in combination therapy as compared with control. Scale bars represent $10 \mu\text{m}$. (E) Metformin, but not 2-DG, treatment shows significant decrease in ATP content of the cells. Error bars indicate SEM. $**P < 0.01$.

Abbreviations: 2-DG, 2-deoxy-D-glucose; ATP, adenosine triphosphate; Gy, grey; Met F, metformin; SEM, standard error of mean.

maximum radiosensitization, and cell survival analysis was performed by clonogenic assay in MCF-7 cells. We observed that a combination of 20 mM 2-DG with 1 Gy reduced twofold cell survival as compared with 1 Gy radiation alone, while its combination with 2 Gy radiation showed 1.6-fold decrease in cell survival than radiation alone (Figure 1A). Further increment in 2-DG dose did not show any significant enhancement in radiosensitization effect. Similarly, treatment of cells with metformin showed dose-dependent radiosensitization effect (Figure 1B). Combination of 2 mM metformin with 1 Gy radiation dose showed 1.5-fold decrease in cell survival as compared with 1 Gy radiation alone, while the combination of 2 mM metformin with 2 Gy radiation resulted in 1.9-fold decrease in cell survival as compared with 2 Gy radiation alone.

Upon deciding the effective minimal drug doses, we also combined both the drugs and tested in combination with 2 Gy Cobalt60 radiation. Cell survival showed significant drop in cells treated with dual drug radiosensitization to that of cells treated with radiation alone ($P=0.0094$, CI =95%) (Figure 1C). In addition to MCF-7 cells, we also evaluated the effect of radiosensitizers on Cobalt60 radiotherapy in MDA MB231 cells (Figure S2). In MDA MB231 cells, 4 mM metformin and 5 mM 2-DG dose showed radiosensitization with 1 Gy, as well as 2 Gy,

radiation dose. Further, combination of both the drugs with 2 Gy radiation showed significant decrease in cell survival as compared with 2 Gy radiation alone ($P=0.0020$, CI =95%). Further, γ H2A.X foci assay was performed to ensure increased cell death is associated to increased DNA double-strand break (Figure 1D), and quantification of the immunofluorescence images indicated increase in foci intensity in cells treated with drug for 12 hours followed by the measured radiation exposure as compared with cells treated with similar radiation dose alone (Figure S3). In case of untreated cells or cells treated with drugs alone, foci formation is recorded as negligible. Since metformin and 2-DG therapy are known to reduce ATP content of the cell, we also analyzed ATP content with or without drug treatment and observed a significant drop in cells treated with metformin ($P=0.0058$, CI =95%), whereas this drop was not significant when cells were treated with 2-DG alone (Figure 1E). Further decrease in ATP content was noted when both the drugs were combined.

Cell cycle analysis was also performed at different time points after the combination therapy, which revealed S-phase arrest after 12 hours of treatment in combination as well as in case of only radiation therapy (Figure 2). It was also observed that after 24 hours and 48 hours of combination therapy, cells remained arrested in S phase, whereas in case of only radiation therapy, this arrest was relieved at these

Figure 2 Cell cycle status at different time points of treatment of MCF-7 cell.

Notes: Data showing elongation of S phase in combination treatment at 24 and 48 hours. However, when treated with radiation alone, S phase elongation is relieved after 12 hours.

Abbreviations: 2-DG, 2-deoxy-D-glucose; Gy, grey; Met F, metformin.

Figure 3 Characterization of MCF-7 cells stably expressing NIS.

Notes: (A) Diagrammatic representation of AttB-CAG-NIS DNA vector used for establishing NIS expressing cell lines. (B) Western blot analysis data showing major band of NIS at 97 kDa for membrane NIS expressing clone indicating dominance of completely glycosylated form of NIS. However, in the clone where NIS is localized in cytoplasm, the band intensity of glycosylated form of NIS (97 kDa band) is less. (C) NIS expression in MCF-7 control cells showing faint staining, and strong cytoplasmic or membrane expression in NIS overexpressing cells. Scale bar represents 20 μ m. (D) Chart showing significantly high ¹²⁵I uptake in MCF-7 NIS cells than in cytoplasmic NIS expressing or parental cells. Blocking by 30 μ M KClO₄ shows complete inhibition of iodine uptake. ** $P < 0.01$. (E) Efflux assay showing maximum efflux of iodine within 15 minutes. Error bars indicate SEM.

Abbreviations: AttB, attachment site of bacteria; CAG, synthetic promoter made by fusion of various elements; Cyto, cytoplasmic; DAPI, 4',6-diamidino-2-phenylindole; NIS, sodium-iodide symporter; SEM, standard error of mean.

time points, indicating prolonged cytotoxic effect of combination therapy.

Established BC cell line overexpressing cell membrane NIS protein demonstrates good iodine influx for short duration

For effective iodide accumulation inside cells, the presence of NIS transporter protein on cell membrane is crucial. Keeping this fact in mind, we engineered a mammalian expression vector for expressing NIS (Figure 3A) and isolated MCF-7 BC clonal cell variants with abundant NIS protein localized either on the cell membrane or in the cytoplasm. NIS transcript and protein expression were confirmed by real-time PCR (data not shown), western blot analysis (Figure 3B), and immunofluorescence (Figure 3C). As shown in Figure 3D, when NIS protein was expressed on the membrane, significantly high iodine accumulation took place as compared

with the cells where NIS was localized predominantly in the cytoplasm ($P = 0.0028$, CI = 95%) or the baseline MCF-7. Treatment of NIS-positive cells with KClO₄, a competitive iodide inhibitor, showed complete inhibition of iodine uptake in cells, indicating NIS-specific iodine uptake. Further, to estimate the duration of iodine retention in membrane expressing clones (MCF-7 NIS), efflux assay was performed, and the results showed that 90% iodine was released within 15 minutes of incubation (Figure 3E).

We have also evaluated the therapeutic efficacy of ¹³¹I treatment on MCF-7 NIS membrane expressing cell line by measuring the survival fraction by clonogenic assay. As detailed in the "Materials and methods" section, a brief exposure of these cells to 50 μ Ci ¹³¹I was found to reduce cell survival significantly ($P < 0.0001$, CI = 95%) than the untreated cells (Figure 4A). Radiation injury causing cellular apoptosis was assessed by annexin V staining

Figure 4 Cytotoxic effect of radioiodine therapy in MCF-7 cells stably expressing NIS.

Notes: (A) Clonogenic assay showing significant decrease in colony formation post ^{131}I treatment. Error bars indicate SEM. $^{**}P < 0.01$. (B) Apoptosis assay showing significant loss in live cell population. (C) Immunofluorescence staining for $\gamma\text{H2A.X}$ foci showing increased number of foci formation in ^{131}I treated cells than in untreated controls.

Abbreviations: NIS, sodium-iodide symporter; SEM, standard error of mean.

protocol, which showed 46.5% live cells in case of ^{131}I treatment as compared with 86% in control (Figure 4B). To confirm that such brief and low-dose exposure of MCF-7 NIS cell still cause DNA double-strand break, $\gamma\text{H2A.X}$ foci formation assay was performed at different time intervals after ^{131}I treatment. Formation of a significantly

large number of foci in treated cells was noted as compared with the untreated cells at 12 hours post-treatment time point (Figure 4C). However, $\gamma\text{H2A.X}$ time kinetics did show that at 48 hours, the number and size of foci significantly diminished (data not shown). Thus, together our results confirm that although major amount of iodine

Figure 5 Effect of 2-DG, metformin, and their combination with ^{131}I treatment on different prospects of MCF-7 NIS cells.

Notes: (A) MCF-7 NIS cell population were treated with 20 mM 2-DG and 4 mM metformin with or without ^{131}I (50 μCi) for 24 hours, and clonogenic assay was performed. Graph showing significant decrease in survival fraction in case of combination therapy as compared with control or only ^{131}I treated cells. $^{**}P < 0.01$; $^{***}P < 0.001$. (B) Immunofluorescence analysis of $\gamma\text{H2A.X}$ foci showing increased expression in combination therapy than various treatment controls as marked. Scale bars represent 10 μm . (C) Apoptotic assay showing significant decrease in live cell population in combination therapy than other treatment controls. (D) Graph showing radiosensitization effect of the combination therapy on Cobalt60 or ^{131}I . Radiosensitization effect by these drugs is more pronounced on radioiodine therapy than Cobalt60 radiation exposure. $^{*}P < 0.05$.

Abbreviations: 2-DG, 2-deoxy-D-glucose; Met F, metformin; NIS, sodium-iodide symporter; ns, not significant.

was effluxed out by these cells within a short duration, still sufficient radio-ablative effects on MCF-7 NIS cell were evident.

Treatment efficacy of radioiodine (^{131}I) is significantly enhanced by 2-DG and metformin combination

The effective 2-DG and metformin dose of 20 mM and 2 mM, respectively, was tested prior to Cobalt60 radiation treatment. The similar treatment was applied in combination with ^{131}I radioisotope in NIS expressing cells. Treatment of cells with 100 μCi of ^{131}I alone resulted in high number of cell death (data not shown), making radiosensitization effects obscure, a lowered concentration (50 μCi) of ^{131}I was used for all the experiments. Combination of dual drugs with 50 μCi ^{131}I showed 9.5-fold decrease in survival fraction when compared with the cells treated with ^{131}I alone ($P=0.0019$) and fourfold decrease when compared with cells treated with the two drugs alone ($P=0.0036$) (Figure 5A). The change in cell survival was further confirmed by annexin V staining to measure apoptosis in parallel (Figure 5C). Cells were analyzed for apoptosis after 48 hours of treatment, and it was observed that live cell population was 37.37% in case of ^{131}I therapy, whereas it was only 11% for the combination therapy. When cells were treated with both the drugs together in absence of ^{131}I , 55.54% cells were alive. Thus, the combination of both the drugs with ^{131}I therapy in NIS-expressing cells results in a significant drop in live cell population. Further $\gamma\text{H2A.X}$ foci

immunofluorescence staining showed increase in number and intensity variation of foci in case of combined treatment, while in case of ^{131}I treatment alone, the number and size of foci were much lower (Figure 5B and Figure S4). We also performed cell cycle analysis after 48 hours of the combination treatment and a very slight S-phase elongation was observed, which was not significant (data not shown). Finally, we compared cytotoxic efficacy of 2 Gy Cobalt60 radiation with that of the 50 μCi ^{131}I , which showed a comparable cell survival fraction (Figure 5D). However, in the presence of the radiosensitizers, a significantly lower survival fraction in ^{131}I than in Cobalt60 radiation was noted ($P=0.0201$, CI=95%). This result confirms that the combination of 2-DG and metformin can effectively radiosensitize BC cell and thus can be combined with either Cobalt60 radiation or NIS-targeted radioiodine therapy.

Since in majority of the patients with BC, NIS expression is localized in the cytoplasm, it is important to evaluate the effect of radiosensitizers combined with ^{131}I in plain MCF-7 cells. Our data suggest that treatment with 50 μCi ^{131}I results in 30% cell death (Figure 6A) as opposed to 60% death in case of MCF-7 NIS overexpressing cells. However, addition of radiosensitizers to the ^{131}I therapy results in significant increase in cell death (70%) as compared with ^{131}I alone ($P=0.0110$, CI=95%). Thus, combination therapy of ^{131}I with radio sensitizers in MCF-7 cells is as effective as ^{131}I therapy alone in MCF-7 NIS cells, where NIS is localized at membrane. We also performed $\gamma\text{H2A.X}$ assay to evaluate DNA damage response in plain MCF-7 cells in presence or absence of radiosensitizers

Figure 6 Effect of 2-DG, metformin, and their combination with ^{131}I treatment on survival of plain MCF-7 cells.

(A) MCF-7 cells were treated with 20 mM 2-DG and 4 mM metformin with or without ^{131}I (50 μCi) for 24 hours, and clonogenic assay was performed. Treatment with ^{131}I alone shows only 30% decrease in cell survival; however, its combination with 2-DG and metformin results in significant loss in live cells (30%, PI =0.0110, ^{131}I vs ^{131}I +2-DG + Met F). * $P<0.05$. (B) Immunofluorescence analysis of $\gamma\text{H2A.X}$ foci showing increased expression in combination therapy than various treatment controls as marked. Scale bars represent 10 μm . It is to be noted that the intensity and frequency of foci is lesser in MCF-7 plain cells when compared with MCF-7 NIS clonal population.

Abbreviations: 2-DG, 2-deoxy-D-glucose; Met F, metformin; NIS, sodium-iodide symporter; PI, propidium iodide.

(Figure 6B). Our data showed foci formation post- ^{131}I therapy in plain MCF-7 cells. Quantification of immunofluorescence images showed increased focal intensity after addition of radiosensitizers (Figure S5). As expected, the intensity of foci in MCF-7 plain cells was much less than the intensity in MCF-7 NIS cells undergoing the same treatment.

Discussion

In the present study, we have combined 2-DG and metformin to sensitize radio-ablative effect in BC cell and tested the cytotoxic efficacy in combination with two different sources of radiation, that is, Cobalt60 radiation or ^{131}I in NIS-expressing MCF-7 cells. 2-DG treatment creates a glucose-deprived state preferentially in cancer cells, and output from glycolysis and pentose phosphate pathway is reduced. Metformin is a bi-guanide and anti-diabetic drug, which inhibits gluconeogenesis in liver, and thus helps to control the sugar levels in patients with diabetes. Previously, metformin was also known to inhibit proliferation of a large range of cancer cells by inducing apoptosis.^{23–26} It has also been shown recently to target cancer-initiating cells.²⁷ Individually, the use of 2-DG and metformin has been reported as radiosensitizer,^{19,20,28} whereas the use of both in combination with low-dose radiation (either from Cobalt60 source or NIS gene-targeted radioiodine [^{131}I]), which is shown for first time during this study, indicated much enhanced radio-ablative effect than using either of the drugs alone. To avoid drug toxicity at higher doses, we used very low dose of these drugs as radiosensitizers, as evident from MTT assay (Figure S1). We observed nearly twofold decrease in survival fraction upon 2 mM metformin treatment in combination with 2 Gy radiation as compared with radiation alone, which corroborates with the earlier study by Song et al, where 5 mM metformin showed radiosensitization effect in MCF-7 cells.²⁰ When 20 mM 2-DG was combined with 2 Gy radiation dose, we observed 1.6-fold decrease in cell survival as compared with radiation alone as was reported in a study by Lin et al.¹² The minor variation in our results may be due to differences in accurate drug dosing, timing of 2-DG, and radiation dose. However, when we combine both 2-DG and metformin with 2 Gy radiation dose, we observe a dramatic difference in cell survival, a nearly threefold decrease ($P=0.0094$, $\text{CI}=95\%$) as compared with 2 Gy radiation alone. Combined drug treatment-mediated cell death proved to be a more significantly enhanced radiosensitization effect than any of the treatment.

Further, our study on $\gamma\text{H2A.X}$ foci formation suggests increase in the intensity of foci in case of dual-drug combined treatment than radiation alone, confirming radiosensitization by these agents. It is noteworthy here that cellular heterogeneity for foci formation encountered can possibly be explained

by the presence of cells at different stages of cell cycle. Previous reports suggest the action of metformin and 2-DG is due to lowered ATP content of the cell. Metformin alters mitochondrial oxidative phosphorylation, and 2-DG inhibits ATP synthesis through glycolysis. Therefore, we have also evidenced lowered ATP content of the cell in case of metformin or dual-drug treatment, but not the 2-DG treatment. Further, we looked into cell cycle status under different drug treatment conditions and observed that combined treatment resulted in elongation of S phase. This S-phase elongation was also observed in case of radiation but was relieved after 48 hours of treatment, whereas it remained nearly constant in case of combination therapy even at 48 hours.

Since 20 mM 2-DG and 2 mM metformin showed effective radiosensitization with Cobalt60 radiation, we maintained the similar dose and time points in experiments combining 2-DG and metformin with ^{131}I treatment in MCF-7 NIS and plain MCF-7 cells. The use of 100 μCi ^{131}I treatment alone resulted in complete cell death (data not shown). Therefore, 50% lower dose of ^{131}I (ie, 50 μCi) was attempted, which showed 1.5-fold decrease in survival fraction in MCF-7 NIS cells as compared with untreated cells. Thus, for all the radioiodine therapy experiments, only 50 μCi of ^{131}I was used. Even at this low radiation dose, the dual-drug use for radiosensitization showed significant decrease in cell survival, which is near-equivalent to the twofold higher dose (ie, 100 μCi alone) of ^{131}I . Apoptosis data corroborate with the result of clonogenic assay, showing significant increase in apoptotic population with the addition of dual-drug radiosensitizers. We have also performed cell cycle analysis after 48 hours of treatment but did not observe any significant differences among samples with different treatments (data not shown). Since localization of NIS is crucial for ^{131}I therapy and various reports suggest cytoplasmic localization of NIS in breast tumor tissues, it is important to evaluate the potential of radio sensitizers with ^{131}I therapy in plain MCF-7 cells. Our results indicate that although ^{131}I therapy alone shows minimal effect on cell survival, combination of radioiodine with radiosensitizers results in significant loss of live cell population in plain MCF-7 cells. Thus, the present study demonstrates the potential of metformin and 2-DG combination treatment as radiosensitizer in combination with Cobalt60 or ^{131}I radiation in NIS-expressing cells. The combination can deliver better therapeutic efficacy than using either of these drugs as a single agent. In addition to their drug toxicity, such radiosensitization effect may further boost mass reduction in cancer cases. To note, at present, there are ongoing clinical trials with both these drugs either in combination with chemotherapy or with radiation.^{29,30} Thus,

combination of 2-DG and metformin with Cobalt60 radiation might play an even more important role in sensitizing radioresistant population as was observed previously with combination of metformin and radiation.²⁰ Previous reports suggest that combination of metformin and 2-DG induces p53-dependent apoptosis in prostate cancer cells through AMPK pathway and metformin inhibits 2-DG induced autophagy in these cells.³¹ In future, it will be interesting to understand how these mechanistic pathways are regulated when Cobalt60 radiation or ¹³¹I is combined with these two drug combination. Further, it would also be important to understand the effect of these drugs in vivo in combination with ¹³¹I in NIS-expressing breast tumor in preclinical and clinical settings.

Conclusion

In the present study, we have shown the use of a combination of two radiosensitizers, 2-DG and metformin, to enhance the radio-ablative effect in combination with either Cobalt60 radiation or targeted NIS-mediated radioiodine therapy. The combination of the radiosensitizers improves therapeutic efficacy effectively in BC cell lines than using any one of the radiosensitizers. Thus, in addition to the drug cytotoxicity, the radiosensitization effect may further boost effective mass reduction in pre-surgical cancer cases.

Acknowledgments

Radioactive iodine-related experiments were carried out in the Radiation Medicine Centre, TATA Memorial Hospital campus, Mumbai, India. Research funding from the Indian Council for Medical Research, New Delhi, to AD (reference number 5/13/25/10/NCD-III); and personal support from the Advanced Centre for Treatment, Research and Education in Cancer, Tata Memorial Centre, to SC for graduate study, is also acknowledged.

Disclosure

The authors report no conflicts of interest in this work.

References

- Keston AS, Ball RP, Frantz VK, Palmer WW. Storage of radioactive iodine in a metastasis from thyroid carcinoma. *Science*. 1942; 95(2466):362–363.
- Seidlin SM, Rossman I, Oshry E, Siegel E. Radioiodine therapy of metastases from carcinoma of the thyroid; a 6-year progress report. *J Clin Endocrinol Metab*. 1949;9(11):1122–1137.
- Tazebay UH, Wapnir IL, Levy O, et al. The mammary gland iodide transporter is expressed during lactation and in breast cancer. *Nat Med*. 2000;6(8):871–878.
- Renier C, Vogel H, Offor O, Yao C, Wapnir I. Breast cancer brain metastases express the sodium iodide symporter. *J Neurooncol*. 2010;96(3): 331–336.
- Ryan J, Curran CE, Hennessy E, et al. The sodium iodide symporter (NIS) and potential regulators in normal, benign and malignant human breast tissue. *PLoS One*. 2011;6(1):e16023.
- Upadhyay G, Singh R, Agarwal G, et al. Functional expression of sodium iodide symporter (NIS) in human breast cancer tissue. *Breast Cancer Res Treat*. 2003;77(2):157–165.
- Wapnir IL, Goris M, Yudd A, et al. The Na⁺/I⁻ symporter mediates iodide uptake in breast cancer metastases and can be selectively down-regulated in the thyroid. *Clin Cancer Res*. 2004;10(13):4294–4302.
- Wapnir IL, van de Rijn M, Nowels K, et al. Immunohistochemical profile of the sodium/iodide symporter in thyroid, breast, and other carcinomas using high density tissue microarrays and conventional sections. *J Clin Endocrinol Metab*. 2003;88(4):1880–1888.
- Chatterjee S, Malhotra R, Varghese F, et al. Quantitative immunohistochemical analysis reveals association between sodium iodide symporter and estrogen receptor expression in breast cancer. *PLoS One*. 2013;8(1):e54055.
- Moon DH, Lee SJ, Park KY, et al. Correlation between 99mTc-pertechnetate uptakes and expressions of human sodium iodide symporter gene in breast tumor tissues. *Nucl Med Biol*. 2001;28(7):829–834.
- Kogai T, Taki K, Brent GA. Enhancement of sodium/iodide symporter expression in thyroid and breast cancer. *Endocr Relat Cancer*. 2006; 13(3):797–826.
- Liu Z, Xing M. Induction of sodium/iodide symporter (NIS) expression and radioiodine uptake in non-thyroid cancer cells. *PLoS One*. 2012; 7(2):e31729.
- Kogai T, Schultz JJ, Johnson LS, Huang M, Brent GA. Retinoic acid induces sodium/iodide symporter gene expression and radioiodide uptake in the MCF-7 breast cancer cell line. *Proc Natl Acad Sci U S A*. 2000;97(15):8519–8524.
- Biaglow JE, Clark EP, Epp ER, Morse-Guadio M, Varnes ME, Mitchell JB. Nonprotein thiols and the radiation response of A549 human lung carcinoma cells. *Int J Radiat Biol Relat Stud Phys Chem Med*. 1983;44(5):489–495.
- Mitchell JB, Russo A. The role of glutathione in radiation and drug induced cytotoxicity. *Br J Cancer*. 1987;58(Suppl):96–104.
- Blackburn RV, Spitz DR, Liu X, et al. Metabolic oxidative stress activates signal transduction and gene expression during glucose deprivation in human tumor cells. *Free Radic Biol Med*. 1999;26(3–4):419–430.
- Parniak M, Kalant N. Incorporation of glucose into glycogen in primary cultures of rat hepatocytes. *Can J Biochem Cell Biol*. 1985; 63(5):333–340.
- Spitz DR, Sim JE, Ridnour LA, Galoforo SS, Lee YJ. Glucose deprivation-induced oxidative stress in human tumor cells. A fundamental defect in metabolism? *Ann NY Acad Sci*. 2000;899:349–362.
- Lin X, Zhang F, Bradbury CM, et al. 2-Deoxy-D-glucose-induced cytotoxicity and radiosensitization in tumor cells is mediated via disruptions in thiol metabolism. *Cancer Res*. 2003;63(12):3413–3417.
- Song CW, Lee H, Dings RP, et al. Metformin kills and radiosensitizes cancer cells and preferentially kills cancer stem cells. *Sci Rep*. 2012;2:362.
- Groth AC, Olivares EC, Thyagarajan B, Calos MP. A phage integrase directs efficient site-specific integration in human cells. *Proc Natl Acad Sci U S A*. 2000;97(11):5995–6000.
- Franken NA, Rodermond HM, Stap J, Haveman J, van Bree C. Clonogenic assay of cells in vitro. *Nat Protoc*. 2006;1(5):2315–2319.
- Ben Sahra I, Laurent K, Loubat A, et al. The antidiabetic drug metformin exerts an antitumoral effect in vitro and in vivo through a decrease of cyclin D1 level. *Oncogene*. 2008;27(25):3576–3586.
- Buzzai M, Jones RG, Amaravadi RK, et al. Systemic treatment with the antidiabetic drug metformin selectively impairs p53-deficient tumor cell growth. *Cancer Res*. 2007;67(14):6745–6752.
- Dowling RJ, Goodwin PJ, Stambolic V. Understanding the benefit of metformin use in cancer treatment. *BMC Med*. 2011;9:33.
- Zakikhani M, Dowling R, Fantus IG, Sonenberg N, Pollak M. Metformin is an AMP kinase-dependent growth inhibitor for breast cancer cells. *Cancer Res*. 2006;66(21):10269–10273.

27. Hirsch HA, Iliopoulos D, Tsiichlis PN, Struhl K. Metformin selectively targets cancer stem cells, and acts together with chemotherapy to block tumor growth and prolong remission. *Cancer Res.* 2009; 69(19):7507–7511.
28. Dwarkanath BS, Zolzer F, Chandana S, et al. Heterogeneity in 2-deoxy-D-glucose-induced modifications in energetics and radiation responses of human tumor cell lines. *Int J Radiat Oncol Biol Phys.* 2001;50(4):1051–1061.
29. Cazzaniga M, Bonanni B, Guerrieri-Gonzaga A, Decensi A. Is it time to test metformin in breast cancer clinical trials? *Cancer Epidemiol Biomarkers Prev.* 2009;18(3):701–705.
30. Mohanti BK, Rath GK, Anantha N, et al. Improving cancer radiotherapy with 2-deoxy-D-glucose: phase I/II clinical trials on human cerebral gliomas. *Int J Radiat Oncol Biol Phys.* 1996;35(1):103–111.
31. Ben Sahra I, Tanti JF, Bost F. The combination of metformin and 2-deoxyglucose inhibits autophagy and induces AMPK-dependent apoptosis in prostate cancer cells. *Autophagy.* 2010;6(5):670–671.

Supplementary materials

Figure S1 Cytotoxic effect of 2-DG and metformin in (A) MCF-7 and (B) MDA MB 231 breast cancer cell lines.

Abbreviations: 2-DG, 2-deoxy-D-glucose; Met F, metformin.

Figure S2 Effect of 2-DG, metformin, and their combination with Cobalt60 radiation on MDA MB 231 cells.

Notes: (A and B) MDA MB231 cells were treated with different dose of 2-DG or metformin, with or without radiation exposure for 24 hours, and clonogenic assay was performed. (C) Combination of 5 mM 2-DG and 4 mM metformin with 2 Gy radiation dose showing further decrease in survival fraction. $**P < 0.01$.

Abbreviations: 2-DG, 2-deoxy-D-glucose; Ctrl, control; Gy, grey; Met F, metformin; ns, not significant.

Figure S3 Effect of Cobalt60 radiation and it's combination with 2-DG and Met F on foci formation in MCF-7 cells.

Notes: (A) Foci assay images from cells treated with Cobalt60 radiation and combination with radio sensitizers showing foci staining with dylight 633 secondary antibody in red channel, nucleus stained with DAPI in blue channel, and merged image showing colocalization of foci with nucleus. Scale bars represent 10 μ m. (B) Quantification of foci from the IF images. In each sample, 50 cells were quantified for intensity of staining from various fields of the cover slip.

Abbreviations: 2-DG, 2-Deoxy-D-glucose; DAPI, 4',6-diamidino-2-phenylindole; Gy, grey; IF, immunofluorescence; Met F, metformin; MFI, mean fluorescence intensity.

Figure S4 (A) Foci assay images from Figure 4 (MCF-7 NIS cells treated with ^{131}I and its combination with radio sensitizers) showing foci staining with dylight 633 secondary antibody in red channel, nucleus stained with DAPI in blue channel, and merged image showing colocalization of foci with nucleus. Scale bars represent 10 μm . **(B)** Quantification of foci from the IF images. In each sample, 50 cells were quantified for intensity of staining from various fields of the cover slip.

Abbreviations: 2-DG, 2-Deoxy-D-glucose; DAPI, 4',6-diamidino-2-phenylindole; IF, immunofluorescence; MET, metformin; NIS, sodium-iodide symporter; MFI, mean fluorescence intensity.

Figure S5 (A) Foci assay images from Figure 5 (MCF-7 plain cells treated with ^{131}I and its combination with radio sensitizers) showing foci staining with dylight 633 secondary antibody in red channel, nucleus stained with DAPI in blue channel, and merged image showing colocalization of foci with nucleus. Scale bars represent 10 μm . **(B)** Quantification of foci from the IF images. In each sample, 50 cells were quantified for intensity of staining from various fields of the cover slip.

Abbreviations: 2-DG, 2-deoxy-D-glucose; DAPI, 4',6-diamidino-2-phenylindole; IF, immunofluorescence; MET, metformin; MFI, mean fluorescence intensity.

Breast Cancer: Targets and Therapy

Publish your work in this journal

Breast Cancer: Targets and Therapy is an international, peer-reviewed open access journal focusing on breast cancer research, identification of therapeutic targets and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

Submit your manuscript here: <http://www.dovepress.com/breast-cancer---targets-and-therapy-journal>

View the full aims and scopes of this journal [here](http://www.dovepress.com/breast-cancer---targets-and-therapy-journal). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress